

MACRO FAUNA

OF THE GANGA RIVER

STATUS AND CONSERVATION
— OF SELECT SPECIES —



National Mission
for Clean Ganga



भारतीय वन्यजीव संस्थान
Wildlife Institute of India

MACRO[◉]
FAUNA
OF THE GANGA
RIVER

STATUS AND CONSERVATION
— OF SELECT SPECIES —

Team Leaders

V. B. Mathur and G.S. Rawat

Editors

Syed Ainul Hussain and Ruchi Badola

Text

Niladri Dasgupta, Shivani Barthwal, Michelle Irengbam, Anita Devi, Sayantika Banerjee

Maps and Database

Zeeshan Ali, Aishwarya R. Chandran, Ravindra Nath Tripathi, Debanjan Sarkar, Satakshi Sharma

Editorial Support

Shivani Barthwal, Michelle Irengbam, Niladri Dasgupta

Survey Team

Aftab Usmani, Goura Chandra Das, Arvind Diwedi, Ajit Kumar, Narendra Mohan Katara, Bitupan Boruah, Kritish Dey, Anuja Mittal, Prabhaker Yadav, Animesh Talukdar, Richika Sah, Chetan P.S. Ahada, Saurav Gawan, Ajay Rawat, Megha Shruti, Umar Saeed, Syed Mujahid Ahmad, Arkojyoti Sarkar

Additional Input

Rajeev Kishore, Sandeep Behera, J. Anthony Johnson, Gopi, G.V., Gautam Talukdar, Abhijit Das, Pariva Dobriyal

Published by

Wildlife Institute of India,
Chandrabani
Dehra Dun - 248001
Uttarakhand, India

ISBN: 81-85496-31-5

May 2018

© Wildlife Institute of India
Wildlife Institute of India, Post Box # 18, Dehra Dun - 248 001
Uttarakhand, India
Tel: +91-135-2646100; 2640114-115;
Fax: +91-135-2640117,
Email: wii@wii.gov.in; gacmc@wii.gov.in
<http://www.wii.gov.in/nmcg/national-mission-for-clean-ganga>

Citation

WII-GACMC (2018). Macro Fauna of Ganga River: Status and Conservation of Select Species. Ganga Aqualife Conservation Monitoring Centre, Wildlife Institute of India, Dehra Dun. Uttarakhand, India Pp. 151

Designed and printed by

XPRESSIONS

+91 9219552563

Doc no. XPS1405181396

PREFACE




Freshwater ecosystems around the world account for only 0.01% of the Earth's surface water and cover a mere 0.8 % of the Earth's surface. Yet, these vulnerable 'aquascapes' are home to an astounding 10% of the world's known species. Human civilization has always centred around these ecosystems however; unsustainable anthropogenic activities have led to widespread loss and degradation of these ecosystems as well as their biodiversity. According to UNEP-WCMC, decline in freshwater species diversity is the highest, surpassing losses in marine and terrestrial species. Among freshwater ecosystems, flowing systems, i.e., rivers and stream are the most vulnerable to species loss. Globally 20% of all known freshwater fish, 44% of waterbirds and 42% of amphibian species are threatened with extinction.

On a global scale, the highest loss in freshwater biodiversity has been reported from the Indian subcontinent, more specifically from the Gangetic plains. This may be accounted for by the high human density in the region and the resultant developmental activities leading to the alteration of natural habitats. This alteration, in turn, has taken a huge toll on the hydrology and water quality of the Ganga River, thereby, effecting its biodiversity. In view of this, the Ministry of Water Resources, River Development and Ganga Rejuvenation, through the National Mission for Clean Ganga and the Wildlife Institute of India has taken up the 'Biodiversity Conservation and Ganga Rejuvenation' project to restore the Ganga River and its biodiversity. The project aims to prepare a science-based aquatic species restoration plan for the Ganga River by involving multiple stakeholders.



To achieve this, robust information on the diversity, abundance and distribution of aquatic macro fauna of the Ganga River, their major threats and the various drivers of these threats causing decline in their populations and habitat are crucial. However, information pertaining to the status and distribution of the aqualife of the Ganga River is limited and scattered. This synthesis is a novel attempt to compile the information on select macro fauna of the Ganga River through literature review and rapid biodiversity assessment from Devprayag in Uttarakhand to Nurgpur in West Bengal. The report aims to develop a knowledge base for the priority species of the Ganga River and identify stretches that are crucial for aquatic biodiversity conservation.

Vinod B. Mathur,
Gopal S. Rawat,
Syed Ainul Hussain &
Ruchi Badola



MESSAGE

नितिन गडकरी
NITIN GADKARI



सत्यमेव जयते

मंत्री
जल संसाधन, नदी विकास, गंगा संरक्षण,
सड़क परिवहन, राजमार्ग एवं पोत परिवहन
भारत सरकार
नई दिल्ली- 110001

MINISTER
WATER RESOURCES
RIVER DEVELOPMENT, GANGA REJUVENATION,
ROAD TRANSPORT, HIGHWAYS AND SHIPPING
GOVERNMENT OF INDIA
NEW DELHI- 110001




I am pleased to know that the Wildlife Institute of India, Dehradun, as part of the 'Biodiversity conservation and Ganga rejuvenation' project, entrusted by the National Mission for Clean Ganga, Ministry of Water Resources, River Development and Ganga Rejuvenation, is releasing this Report on the "Status and Conservation of select Aquatic Species of the Ganga River."

The River Ganga, right from its source at Gaumukh to its confluence with Bay of Bengal at Ganga Sagar, nurtures a dense human population which poses serious threat to its aquatic wildlife. The growing urbanisation and economic growth is likely to further intensify these threats. Therefore, rejuvenation of the river ecosystem is critical to restore its ecological integrity and to ensure the survival of its aquatic species such as the iconic Gangetic river dolphin, which is our national aquatic animal.

I appreciate the efforts made by the Wildlife Institute of India and its team in bringing out this valuable report on conservation of River Ganga and its biodiversity including the aspects on *Aviral Dhara* and *Nirmal Dhara* of River Ganga.

(Nitin Gadkari)



MESSAGE

डॉ. सत्य पाल सिंह
Dr. Satya Pal Singh



सत्यमेव जयते

मानव संसाधन विकास; और
जल संसाधन, नदी
विकास एवं गंगा संरक्षण राज्य मंत्री
भारत सरकार
नई दिल्ली- 110001


MINISTER OF STATE FOR
HUMAN RESOURCE DEVELOPMENT;
WATER RESOURCES,
RIVER DEVELOPMENT AND
GANGA REJUVENATION
GOVERNMENT OF INDIA
NEW DELHI- 110001



The Wildlife Institute of India, Dehra Dun has been doing pioneering work in biodiversity related issues over the last three decades, across the country. Its active engagement in the conservation of Ganga's aqualife under the Namami Gange Program is a very positive step towards Ganga rejuvenation. Their recent Report under the project assigned to them by National Mission for Clean Ganga is expected to add new dimensions for further research & development and conservation plans.

I extend hearty congratulations to the Institute & the project team for making valuable contribution to the cause of biodiversity conservation & Ganga rejuvenation and wish them success in future endeavours.

(Dr. Satya Pal Singh)



MESSAGE

सी.के.मिश्रा
C.K. Mishra



सचिव
भारत सरकार
पर्यावरण, वन एवं जलवायु परिवर्तन मंत्रालय
नई दिल्ली-110003

SECRETARY
GOVERNMENT OF INDIA
MINISTRY OF ENVIRONMENT, FOREST & CLIMATE CHANGE
NEW DELHI- 110003




It is a matter of great satisfaction that the report "Macro Fauna of the Ganga River: Status and Conservation of Select Species", has been prepared by the Wildlife Institute of India, Dehradun, under the project funded by the National Mission for Clean Ganga "Biodiversity Conservation and Ganga Rejuvenation".

More than one third of India's population is dependent on the Ganga River and supports a diverse assemblage of aquatic fauna. The river is highly stressed due to unsustainable resource extraction, construction of dams and barrages, and release of effluents and solid wastes. The integrity of the Ganga River is under jeopardy, necessitating interventions at all levels considering both the needs of the local communities and the aquatic wildlife.

The information provided in this report on the past and present status and distribution of select species of conservation importance of the Ganga River, and the various threats faced by them would be crucial for conservation initiatives. This report aptly covers issues pertaining to the river conservation and brings out an important aspect of involving multiple stakeholders for conservation actions, which is a novel approach.

The hard work put in by the Wildlife Institute of India for this report, praiseworthy. It is particularly gratifying to me that this quality work has been done by Wildlife Institute of India, an integral part of the Ministry of Environment, Forest and Climate Change.


(C.K. Mishra)



MESSAGE



यू.पी. सिंह
U.P. Singh



सत्यमेव जयते

सचिव
भारत सरकार
जल संसाधन, नदी विकास एवं
गंगा संरक्षण मंत्रालय
नई दिल्ली - 110001

SECRETARY
GOVERNMENT OF INDIA
MINISTRY OF WATER RESOURCES,
RIVER DEVELOPMENT & GANGA REJUVENATION
NEW DELHI - 110 001




I am pleased to know that the Wildlife Institute of India, Dehradun is successfully carrying out various tasks under the project 'Biodiversity Conservation and Ganga Rejuvenation' entrusted by the National Mission for Clean Ganga, Ministry of Water Resources, River Development and Ganga Rejuvenation, Government of India.

Rivers are the lifelines of an ecosystem, providing direct services such as water and food as well as indirect services such as groundwater recharge. Human civilizations have always centered around river systems. The Ganga River is one of the largest river systems in the world. It harbours important aquatic fauna such as the Gangetic river dolphin, gharial, otters, turtles and several aquatic and terrestrial birds. The aquatic wildlife of the Ganga and their habitats are in peril due to reduction in water flow, pollution and over exploitation of riverine resources.

The restoration of the Ganga River and its key biodiversity, therefore, is one of the most important national agendas of the Government of India. This report on the "Status and Conservation of selected species of Ganga River" provides in-depth information on the biodiversity of the Ganga River. The report is a repository of knowledge on the priority species of the Ganga River, and if the recommendations provided herein for the conservation of the river and its priority aquatic species are implemented thoroughly, it will aid in achieving NMCG's mandate of restoring the biodiversity of the Ganga River. This will also facilitate in accomplishing *Namami Gange's* objectives of *Aviral Dhara* and *Nirmal Dhara*.

I congratulate the Wildlife Institute of India and the project team for the tremendous amount of effort they have made in bringing out this report. I sincerely believe that this will make a significant contribution towards aquatic wildlife conservation in India and assist in making informed policy decisions for the conservation of the Ganga River and its aqualife.

(U.P. Singh)



MESSAGE



राजीव रंजन मिश्रा
Rajiv Ranjan Mishra



सत्यमेव जयते

महानिदेशक
भारत सरकार
जल संसाधन, नदी विकास एवं गंगा संरक्षण मंत्रालय
राष्ट्रीय स्वच्छ गंगा मिशन
नई दिल्ली – 110002
DIRECTOR GENERAL
GOVERNMENT OF INDIA
MINISTRY OF WATER RESOURCES,
RIVER DEVELOPMENT AND GANGA REJUVENATION
NATIONAL MISSION FOR CLEAN GANGA
NEW DELHI - 110002




It gives me immense pleasure to learn that the Wildlife Institute of India, Dehradun has successfully prepared this status report on conservation of select aquatic species of Ganga River, under the aegis of its NMCG-funded 'Biodiversity Conservation and Ganga Rejuvenation' project.

2. Entwined with our culture and tradition, the holy river Ganga provides livelihood to more than a third of the country's population. It is also a habitat to a wide variety of aquatic fauna – whose conservation is globally significant to a large extent.

3. The population load and concomitant development activities have, however, led to degradation of the river regime, thereby detrimental to the fauna it supports. The reduced water availability is playing a major role in the declining population of the aquatic biodiversity of the Ganga River. Restoring this is a daunting task and calls for novel innovations and understanding of science as well as society. In view of this, NMCG has entrusted the Wildlife Institute of India with the project 'Biodiversity Conservation and Ganga Rejuvenation' under the *Namami Gange* programme of the Ministry of Water Resources, River Development and Ganga Rejuvenation. The project aims at a science-based aquatic species restoration plan for the Ganga River and to ensure that threat to the biodiversity of the Ganga River is significantly reduced by 2020.

4. The faunal diversity of the Ganga River has captured the attention of many scholars, and thus a pool of literature is available. However, this information is scattered and needs to be consolidated to enhance its accessibility. This report is an attempt to consolidate this information, and to provide a status on priority aquatic species. It is a collation and synthesis based on an extensive literature review and biodiversity assessment of the Ganga River by the Wildlife Institute of India. I envisage that the information contained in this report will aid in bringing together a strategic conservation plan for these species and would provide direction for the biological restoration process of our national river Ganga.

(Rajiv Ranjan Mishra)



MESSAGE

सिद्धान्त दास
Siddhanta Das



वन महानिदेशक एवं विशेष सचिव
भारत सरकार
पर्यावरण, वन एवं जलवायु परिवर्तन मंत्रालय
नई दिल्ली- 110003

DIRECTOR GENERAL OF FOREST & SPL. SECY
GOVERNMENT OF INDIA
MINISTRY OF ENVIRONMENT, FOREST & CLIMATE CHANGE
NEW DELHI- 110003



I am happy to present the report on the "Status and conservation of select aquatic species of the Ganga River" prepared by the Wildlife Institute of India, Dehra Dun as part of the 'Biodiversity conservation and Ganga rejuvenation' project, entrusted by the National Mission for Clean Ganga, Ministry of Water Resources, River Development and Ganga Rejuvenation.

The Ganga is 2525 km in length and traverses across India and Bangladesh. In India, the river flows through five states viz. Uttarakhand, Uttar Pradesh, Bihar, Jharkhand and West Bengal and covers an area of 861,404 km², representing five bio-geographic zones and about 26% of the country's land mass and is a lifeline for about 43% of the Indian population. It not only adds to the ecological wealth of the India but also symbolizes the rich cultural diversity of the country.

The high dependency of a growing population and developing economy has taken its toll on the ecological health of the Ganga River. Hence, restoration and rejuvenation of the Ganga River is critical to maintain the ecological integrity and, livelihood security and overall wellbeing of the dependent human population. The present report is a significant scientific documentation of the ground realities and information that can be used for the long-term monitoring of the aqualife of Ganga River.

I congratulate the Wildlife Institute of India team for this comprehensive documentation of present species level scenario in Ganga River and I am confident that the information provided will prove to be an excellent base for developing future management and conservation interventions.


(Siddhanta Das)

CONTENTS

ACKNOWLEDGEMENTS

EXECUTIVE SUMMARY

1

INTRODUCTION

01

2

GANGA RIVER

05

- 2.1 Course of the Ganga River
- 2.2 Geology and Geomorphology
- 2.3 Soil Types
- 2.4 Climate
- 2.5 Biogeography, Flora and Fauna
- 2.6 Demography

3

STATUS OF AQUALIFE IN THE GANGA RIVER

11

3.1 Methodological framework

3.2 Mammals

- 3.2.1 Gangetic river dolphin (*Platanista gangetica*)
- 3.2.2 Otters

3.3 Avifauna

- 3.3.1 Black-bellied tern (*Sterna acuticauda*)
- 3.3.2 Indian sarus crane (*Antigone antigone*)
- 3.3.3 Indian skimmer (*Rynchops albicollis*)
- 3.3.4 River tern (*Sterna aurantia*)
- 3.3.5 River lapwing (*Vanellus duvaucelii*)
- 3.3.6 Painted stork (*Mycteria leucocephala*)
- 3.3.7 Great thick-knee (*Esacus recurvirostris*)
- 3.3.8 Little pratincole (*Glareola lactea*)

3.4 Reptiles

3.4.1 Crocodilians

- 3.4.1.1 Gharial (*Gavialis gangeticus*)
- 3.4.1.2 Mugger (*Crocodylus palustris*)
- 3.4.1.3 Estuarine crocodile (*Crocodylus porosus*)

3.4.2 Turtles

- 3.4.2.1 Red-crowned roofed turtle (*Batagur kachuga*)
- 3.4.2.2 Northern river terrapin (*Batagur baska*)
- 3.4.2.3 Three-striped roofed turtle (*Batagur dhongoka*)
- 3.4.2.4 Spotted pond turtle (*Geoclemys hamiltonii*)
- 3.4.2.5 Indian tent turtle (*Pangshura tentoria*)
- 3.4.2.6 Brown roofed turtle (*Pangshura smithii*)
- 3.4.2.7 Indian roofed turtle (*Pangshura tecta*)
- 3.4.2.8 Crowned river turtle (*Hardella thurjii*)
- 3.4.2.9 Indian black turtle (*Melanochelys trijuga*)
- 3.4.2.10 Indian softshell turtle (*Nilssonia gangetica*)
- 3.4.2.11 Indian peacock softshell turtle (*Nilssonia hurum*)
- 3.4.2.12 Indian narrow-headed softshell turtle (*Chitra indica*)
- 3.4.2.13 Indian flapshell turtle (*Lissemys punctata*)

3.5 Amphibians

- 3.5.1 Himalaya paa frog (*Nanorana vicina*)
- 3.5.2 Annandale's paa frog (*Nanorana annandalii*)
- 3.5.3 Nepal paa frog (*Nanorana minica*)
- 3.5.4 Cascade frog (*Amolops formosus*)
- 3.5.5 Marbled toad (*Duttaphrynus stomaticus*)
- 3.5.6 Tytler's pond frog (*Hylarana tytleri*)
- 3.5.7 Indian bullfrog (*Hoplobatrachus tigerinus*)
- 3.5.8 Jerdon's bullfrog (*Hoplobatrachus crassus*)
- 3.5.9 Dudhwa tree frog (*Chiromantis dudhwaensis*)

3.6 Fish

- 3.6.1 Golden mahaseer (*Tor putitora*)
- 3.6.2 Bronze featherback (*Notopterus notopterus*)
- 3.6.3 Freshwater shark (*Wallago attu*)
- 3.6.4 Minor carp (*Labeo bata*)
- 3.6.5 Spotted barb (*Puntius sophore*)

3.7 River Stretches with High Biodiversity Value

4

THREATS TO THE BIODIVERSITY OF GANGA RIVER

83

- 4.1 Upper Ganga
- 4.2 Middle Ganga
- 4.3 Lower Ganga
- 4.4 Conclusions

5

CONSERVATION IMPLICATIONS

93

REFERENCES

97

ANNEXURES

111





ACKNOWLEDGEMENTS

The Wildlife Institute of India would like to acknowledge the support provided by the National Mission for Clean Ganga (NMCG) and the Ministry of Water Resources, River Development and Ganga Rejuvenation (MoWR, RD&GR) for the implementation of the 'Biodiversity Conservation and Ganga Rejuvenation' project. We acknowledge the support provided by the forest departments of the five Ganga River states in carrying out the rapid biodiversity assessment along the river. We would also like to acknowledge the support provided by the Nuclear Power Corporation of India Limited (NPCIL) and the Narora Atomic Power Station (NAPS) for the establishment of a rescue and rehabilitation centre for distressed aqualife. We acknowledge the support provided by the field staff and the local people during the rapid biodiversity assessment of the Ganga River. We would like to thank the entire Ganga team at WII for their help and fruitful discussions.

EXECUTIVE SUMMARY



Freshwater ecosystems harbour a rich diversity of species and provide habitats for at least 9.5% of all animal species. However, freshwater diversity is declining at an alarming rate due to a multitude of factors, most of which can be attributed to human-induced habitat loss or degradation and alteration. It is being realized globally that the loss of aquatic ecosystems often have repercussions on the economic and social well-being. Thus, conservation planners are turning to solutions for ecological restoration of rivers and the processes therein.

The Ganga River is one of the largest and most diverse river systems of Asia supporting one of the highest human densities in its basin. Varied geomorphological features along the Ganga River, coupled with climatic variations, result in a continuous gradient of habitats for various life forms, right from the headwaters to its mouth. Encompassing three ecologically diverse biogeographic zones, the river also harbours important faunal species such as the Gangetic river dolphin, gharial, otters, turtles and several aquatic and terrestrial birds. However, the aquatic diversity of the Ganga River and their habitats are in peril due to a multitude of factors arising from unsustainable resource use practices.

The restoration of the Ganga River and its key biodiversity, therefore, is one of the most important items on the national agenda of the Government of India. Science-based aquatic species restoration of the river by involving multiple stakeholders is thus planned through the National Mission for Clean Ganga's project 'Biodiversity Conservation and Ganga Rejuvenation' entrusted to the Wildlife Institute of India. This synthesis is an attempt to compile the available information, consolidate the baseline data on the historical and present distribution of its aquatic species, identify areas with high biodiversity values and threats to the ecological integrity of river. The species that are most vulnerable to habitat alterations, endemic species and species listed in the IUCN Red List and/or the Indian Wild Life (Protection) Act, 1972 (IWPA) were selected as priority species for the synthesis and thus for future conservation action.



An extensive literature review and the rapid biodiversity assessment of the Ganga River revealed that the distribution range of priority aquatic species such as the Gangetic river dolphin, gharial and mugger have reduced significantly. Loss of longitudinal and lateral connectivity and limited availability of suitable habitat conditions have restricted the populations of these species to smaller stretches of the river. A reduced availability of pools deeper than 4 m was found to be the most limiting factor, hampering the movement and distribution of priority species, namely the Gangetic river dolphin and gharial. Only 38.7% of the river had a depth of 4 m or more during the summer season. Structural changes in the river morphology due to dams, barrages, bank alteration, agriculture and sand mining are other major factors that have affected the distribution of priority species and threaten their survival. The rapid biodiversity assessment revealed that the populations of the priority aquatic species were mostly restricted to the relatively undisturbed areas, including the Protected Areas (PAs). These PAs account for around 10% of the total length of the river, including parts of the Sundarban Tiger Reserve. The priority species were found to be distributed along six high biodiversity stretches along the river and these stretches together formed about 50% of the surveyed stretch. These stretches are Devprayag to Rishikesh (61 km), Makdumpur to Narora (147 km), Bhitaura to Ghazipur (454 km), Chhapra to Kahalgaon (296 km), Sahibganj to Rajmahal (34 km) and Baharampur to Barrackpore (246 km). The geomorphology of these stretches and the issues affecting their conservation values are varied. Therefore, restoration attempts should primarily focus on these high-biodiversity stretches, and site specific conservation planning should be initiated in consultation with various stakeholders.

The differential resource use scenarios along the river have given rise to diverse issues and threats. Hence, measures to address these threats should be strategic and stretch-specific. The upper stretch has undergone immense structural changes, and any further changes would undermine the ecological integrity of the river along with its religious and recreational values. Agricultural intensification along the river banks and on the river islands, and sand mining in the middle stretch of the river have altered the habitats of island-nesting birds and turtles. The lower stretch of the Ganga River is also highly modified by the Farakka Barrage. Flow alterations and changes in salinity regimes in the estuaries has resulted in changes in the species assemblage in this stretch. Impacts of climate change are exacerbating these threats in the form of an unpredictable flow regime and changes in both the upstream and downstream distribution ranges of species.

The water quality in the middle and lower stretches of the Ganga River has been degraded due to the influx of wastewater from domestic and industrial sources and agricultural run-off.

The altered water quality is aggravated by the low flow conditions in the Ganga River. Indiscriminate use of synthetic chemicals in agriculture and healthcare releases persistent pollutants such as heavy metals and other endocrine-disrupting chemicals, which interfere with the hormonal and immune systems of the aquatic species through bioaccumulation and biomagnification along the food chain. Evidence of bioaccumulation and biomagnification has been documented in the food web of the Gangetic river dolphin.

Critical areas in the middle stretch with high susceptibility to these threats include stretches from Farrukabad to Kanpur and from Allahabad to Varanasi. The stretches from Sitab Diara to Ajimganj/Jiaganj and from Barrackpore to Falta were the critical areas in the lower stretch.

However, planning restoration for the Ganga River is not only a river conservation issue but also has significant social dimension owing to the high dependence of the dense human population living along the river. In this regard, efforts should be made to raise awareness among stakeholders and to garner societal support. Site-specific strategies need to be developed by aligning the local people's livelihoods with conservation priorities. Stakeholders, including the local communities and policy makers need to be made aware of the role of an ecologically intact Ganga River in enhancing the quality of life. To maintain a balance between societal and ecological needs, sustainable and efficient water use practices in norms with modern abstraction techniques should be incorporated in the water policy and allocation guidelines. Prioritizing efficient water usage in consonance with the needs of aquatic species and agricultural demand would help augment flows, enhancing habitat suitability for aquatic fauna and retain sufficient water in the river to dilute the pollution load. Strengthening of the PA network, particularly creation of conservation reserves in the biodiversity rich stretches of the river through strategic planning will restore ecological values in select stretches, which will assist in natural recolonization of areas devoid of species of conservation significance.

INTRODUCTION

Freshwater ecosystems harbour a rich diversity of species and habitats, and although it still remains incompletely studied, it is conservatively estimated that freshwater ecosystems provide suitable habitats for at least 9.5% of all animal species (Balian et al., 2008; Collen et al., 2014), nearly 6% of all described species (Dudgeon et al., 2006) and one-third of all vertebrate species (Strayer, 2010).

The patterns and processes of interactions amongst these species and between species and their habitats account for numerous ecosystem services that are critical to the survival of all life forms, including humans (Carrizo et al., 2013). However, freshwater diversity is declining at an alarming rate due to a multitude of factors (Revenga et al., 2005; Collen et al., 2009; Strayer & Dudgeon, 2010; Darwall et al., 2011). On an average freshwater populations declined by 81% between 1970 and 2012 with an average annual decline of 3.9% (WWF, 2016), most of which can be attributed to human-induced habitat loss or degradation (Collen et al., 2014).

Ecological studies have proved that there is a correlation between species diversity and ecosystem stability as well as productivity (Odum, 1953; MacArthur, 1955; Elton, 1958; Gardner & Ashby, 1970; May, 1973). In effect, ecosystem stability depends on the ability of communities to contain species, or functional groups, that are capable of differential response, and the greater the species diversity of a biological community, the lesser the risk of ecological collapse (Palmer et al., 2010; Mellin et al., 2014). Additionally, ecoregional species richness and riverine productivity have also been found to be positively correlated (Duffy et al., 2007; McIntyre et al., 2016). Unless appropriate actions are taken to maintain this diversity, ecosystem processes will be disrupted (Pressey et al., 2007). Declines in species diversity will lead to reduced ecological stability and productivity and, ultimately, to ecological collapse



(Mellin et al., 2014), severely affecting human well-being (Feld et al., 2011).

With increasing evidences of the ecological, economic and social losses associated with loss of aquatic systems, especially rivers (Vorosmarty et al., 2010), conservation planners are turning to solutions for ecological restoration of rivers (Lake et al., 2007; Palmer et al., 2004). There are many theories and physical measures (single species focus, channel re-configuration and in-stream habitat improvement) related to restoration. However, the concept of the restoration of an entire stream or river ecosystem by restoring its biodiversity is widely accepted (Lake et al., 2007; Palmer et al., 2010).

The aim of ecological restoration is to enhance the biodiversity and the resultant ecosystem services (Palmer et al., 2007). Maintenance of habitat connectivity, seasonal flow variability, augmenting habitat suitability and addressing issues related to unsustainable resource use are some of the parameters for successful ecological restoration of a river or a stream and are seldom seen in the restoration process (Lake et al., 2007).

The Aichi targets of the Convention on Biodiversity have set priorities for threatened freshwater taxa occurring outside the present global PA network (Raghavan et al., 2016). However, success would rely on science-based systematic conservation efforts at the national, state and local levels. Systematic planning requires prioritization of selected stretches of a river on the basis of their biodiversity and habitat suitability through participatory planning and collaborative implementation of strategies,



decisions and actions that secure the long-term survival and favourable conservation status of biodiversity (Kukkala & Moilanen, 2013). However, owing to the incomplete biological information available for any riverscape, setting conservation priorities is a challenging task (Scott et al., 1993; Noss et al., 1997). Nevertheless, the fragmented state of riverscapes and the high degree of habitat loss and fragmentation aggravates the urgency to conserve remaining habitats, the refuges that can act as repositories of source populations.

To address these problems, conservation planners have developed alternative methods to identify and prioritize conservation areas. A common approach used to prioritize conservation areas involves identification of umbrella species and their habitat requirements. Conservation areas are selected on the basis of the existing and potential habitat of the umbrella species and the connectivity among those habitat patches; which in turn would provide sufficient habitat variability to confer protection to a large number of naturally co-occurring species of conservation interest (Launer & Murphy, 1994; Berger, 1997; Martikainen et al., 1998).

Most of the freshwater conservation literature available focuses on systematic, strategic and landscape-level planning through the declaration of freshwater PAs (Saunders et al., 2002; Kingsford et al., 2005; Abell et al., 2007), which in principle should take into account the catchment characteristics of the river basin (Abell et al., 2007). This approach, although coherent, is not practical for river basins with high human densities where social and economic constraints exist, and it is not feasible to protect all areas (Nel et al., 2009). Therefore, the question arises as, how to conserve the freshwater ecosystems in these densely populated areas, where the majority of the population lives in perpetual poverty and is highly dependent on the resources of the river.



In view of the foregoing discussion, a synthesis on the status and conservation of the macro fauna of the Ganga River was carried out. A comprehensive review was carried out to identify the species assemblage of the river, which is provided in Annexures I to VII. This synthesis critically examines the historical and current status of the priority aquatic species and the factors threatening their survival. It aims to identify the most strategic and feasible approach to conserve human-dominated river basins in the wake of the expanding population of the country and the increasing threats to rivers and their biodiversity. The report comprises a comprehensive strategy to restore the biodiversity values of the Ganga River and is a result of extensive primary and secondary studies carried out under the project 'Biodiversity Conservation and Ganga Rejuvenation', funded by the National Mission for Clean Ganga (NMCG). The project aims to prepare a science-based aquatic species restoration plan for the Ganga River by involving multiple stakeholders. The Ganga Aqualife Conservation Monitoring Centre (GACMC), established under the mandate of the project, intends to build a scientific knowledge base on the Ganga River, develop national guidelines for consideration of aquatic fauna in the planning and execution of water infrastructure development and to disseminate this knowledge to the general public as well as the scientific community. The species most vulnerable to habitat alterations, endemic species and species listed as Critically Endangered, Endangered, Threatened or Vulnerable in the IUCN Red List or listed in the Indian Wild Life (Protection) Act, 1972 (IWPA) were selected as priority species for the synthesis.



GANGA RIVER



The Ganga River is among the largest rivers in Asia. It flows through India and Bangladesh and has been a cradle of human civilization since time immemorial (Trivedi, 2010; Kumar, 2017a). It is considered one of the most sacred rivers in the world. In Hindu mythology it is worshipped as “Goddess Ganga” and is believed to have been brought to earth from heaven by King Bhagirath (Kumar, 2017a). Historically, several provincial or imperial capitals were located along its banks, and the Ganga basin was the centre of power from the Mauryan period to the Mughal era (Wink, 2002; Kumar, 2017a). The Ganga River was declared the “National River” on 4 November 2008 (Sanghi & Kaushal, 2014). The Ganga River basin is spread across 11 states and covers 26.3% of India's total geographical area (Sanghi & Kaushal, 2014). The mainstem Ganga River flows through the five Indian states, viz., Uttarakhand, Uttar Pradesh, Bihar, Jharkhand and West Bengal. The remaining six states in which the basin lies are Delhi, Haryana, Himachal Pradesh, Madhya Pradesh, Chhattisgarh and Rajasthan (NGRBA, 2011).



Course of the Ganga River

2.1

The mainstem Ganga River (Goumukh to the Bay of Bengal) is roughly 2525 km in length (Tandon & Sinha, 2018), with 1450 km falling in Uttarakhand and Uttar Pradesh, 445 km in Bihar and Jharkhand and 520 km in West Bengal (Sanghi & Kaushal, 2014). The Bhagirathi River is considered the source stream of the Ganga River. It originates from the Gangotri glacier at Goumukh, in the Western Himalaya, and merges with the Alaknanda River at Devprayag, where the “Ganga River” is formally formed. After this, the Ganga River flows through narrow Himalayan valleys and reaches the plains at Haridwar (NRCD, 2009). At Haridwar, the Bhimgoda Barrage diverts water to the Upper Ganga Canal for irrigation. Further down, two barrages at Bijnor and Narora divert the water into the Madhya Ganga Canal (only during the monsoon) and Lower Ganga Canal, respectively. Thereafter the river is joined by the Ramganga River at Kannauj and the Yamuna River at Allahabad (NGRBA, 2011). Downstream of Allahabad, the river is met by the Tons, Son, Punpun, Gomti, Ghagra, Gandak and Kosi rivers, flowing through Varanasi, Patna and Bhagalpur (Tandon & Sinha, 2018). At Pakur, the Farakka Barrage regulates the flow of the Ganga River by diverting water into a feeder canal linking the Hooghly River. Downstream of this barrage, the Ganga River bifurcates to form the Hooghly River, which flows through West Bengal before merging into the Bay of Bengal, and the main branch, which enters Bangladesh as the Padma River (NRCD, 2009).



The main stem of the Ganga River may be divided into three stretches: the upper stretch (from Goumukh to Haridwar), the middle stretch (from Haridwar to Varanasi) and the lower stretch (from Varanasi to Ganga Sagar). These divisions reflect variations in geology, geomorphology, soil type, climate, flora and fauna and social and economic issues.

Geology and Geomorphology

2.2

Structurally, the Ganga River basin is made of three large divisions of the Indian subcontinent, namely the Himalayan Fold Mountains and the Central Indian Highlands; the Peninsular Shield; and the Gangetic Plains (NRCD, 2009).

The Ganga River basin occupies the Himalayan foredeep in a northeast–southwest-oriented elongate depression (Tandon & Sinha, 2018), formed in response to the uplift of the Himalaya after the collision of the Indian and Asian plates (Singh et al., 1990; Sinha et al., 2005). The average gradient of the basin is 20 m/km (Valdiya, 2016). The Gangetic Plain is made of two discrete hinterlands—the Himalaya in the north and the cratons to the south (Tandon & Sinha, 2018). It evolved as a result of the filling up of the foredeep basin with a thick deposition of sediments derived chiefly from the Himalaya and partly from the hills of northern peninsular India (Das, 2014; Valdiya, 2016). The terrain along the Ganga River is divided into two parts, the northern part, composed of alluvial terrain of Quaternary age, and the southern part, composed of consolidated crystalline and sedimentary formations of Archaean-to-Palaeozoic age (Wadia, 1961). Two distinct physiographic and chronological units are recognized in this plain: the Older Alluvium, comprising coarser sedimentary units of the Banda, Varanasi and Bhangar, and the Newer Alluvium, consisting of the Khadar, Bhur and Bhabhar (Valdiya, 2016).

The Ganga River, in its upper stretch, from Gangotri to Haridwar, flows through steep, mountainous valleys of the Himalaya, consisting of a partly confined floodplain and braided channels (Sinha et al., 2017;

Tandon & Sinha, 2018). The middle stretch, from Haridwar to Varanasi, winds through the Gangetic Plain region and consists of the piedmont, craton margin and valley interfluve landscape; a partly confined floodplain; and braided and sinuous channels (Sinha et al., 2017).

The lower stretch of the Ganga River, from Varanasi downstream, passes through the Gangetic Plain and before merging with the Bay of Bengal forms the distributary-delta system south and east of Farakka (Tandon & Sinha, 2018). It is made of craton margin and valley interfluve landscape settings; a partly confined and confined floodplain; and sinuous, braided channels and anabranches (Sinha et al., 2017). The Sundarban delta was formed in the Tertiary period as a result of tectonic movements in northwestern Punjab and a southeastern flow of the Ganga River, resulting in the deposition of sediments in the Bengal basin (Gopal & Chauhan, 2006; Wadia, 1961). The delta complex formed as a result of combined contributions of the Ganga and the Brahmaputra rivers is the world's largest delta, covering an area of 60,000 km² (Valdiya, 2016).

Soil Types

2.3

The Ganga River basin is characterized by a wide variety of soils. The upper stretch of the basin consists of mountain and sub-montane soils covering around 1.59% and 4.28% of the area, respectively, and are highly erodible. The middle and lower stretches of the Ganga River basin are made of mainly alluvial soil (covering 52.44% of the area), with very high erodibility. Red soil covers 12% of the area, and is also highly erodible. Red and yellow soils and mixed red-and-black soils cover 8% of the area, and they are moderately erodible. Deep black soils and medium black soils cover 14% of the area, and have low erodibility. The remaining 6% of the area has shallow black soils and lateritic soils. The erodibility of these

soils is very low (NGRBA, 2011; NRCD, 2009). It can be concluded that the Ganga River basin is highly erodible (NRCD, 2009; Sanghi & Kaushal, 2014), as a result of which the floodplains and banks are vulnerable to erosion through run-off.

Climate

2.4

Due to significant variations in the geographical extent of the Ganga River, including altitudinal variations, specifically at the source the temperature along the river basin varies between 5°C and 45°C (Misra, 2011; Jain et al., 2016). The climate is positively influenced by the southwest monsoon (Ghosh, 1991), which extends from June to October. During the monsoon, the basin receives almost 85% of the total annual rainfall. The annual average rainfall varies between 39 cm and 200 cm, with an average of 110 cm (NRCD, 2009). The average annual temperatures are high throughout the basin except the high-elevation areas of the upper Ganga River basin (Jain et al., 2016). They rise sharply from March to May (summer season), from 30°C to 45°C. The average temperature of the basin during winter season (from November to February) is between 5°C and 30°C (Misra, 2011).

The upper stretch of the river, close to the snow line, receives snowfall, and 100-250 cm of annual rainfall (number of rainy days being 100-150). The annual average temperature varies from <10°C to 20°C (Nautiyal, 2010).



The middle stretch and most of the lower stretch of the river flow through the Gangetic Plains. The average annual rainfall in the plains varies between 60 cm and 160 cm, with the western parts receiving lesser rainfall (from 60 cm to 140 cm) in comparison with the eastern parts (from 90 cm to >160 cm). The northern part of the plains receives higher rainfall compared with the southern part. The temperature varies from 5°C to 25°C in winter and from 20°C to more than 40°C during summer in the Gangetic Plain (Sinha et al., 2005).

The climate of the Sundarban delta region of the lower stretch is characterized by relatively high temperatures and humidity (>80%) throughout the year. The average annual rainfall is 166.1 cm and decreases from 180.5 cm near Sagar Islands, in the south, towards Kolkata, in the north. The daily minimum temperature during winter varies between 2°C and 4°C, and rises to a maximum of about 43°C in March and may exceed 32°C during the monsoon. Thunderstorms in the afternoon are common during summer and are known as Kalbaisakhi (disastrous winds of Baisakh) (Gopal & Chauhan, 2006).

Biogeography, Flora and Fauna

2.5

The Ganga River harbours a unique assemblage of biodiversity due to its passage through three distinct biogeographic zones, viz., the Himalayas (West Himalaya - 2B), Gangetic Plains (Upper Gangetic Plain - 7A and Lower Gangetic Plain - 7B) and Coasts (East Coast - 8B). A smaller stretch of the river, in Jharkhand state, also borders the Deccan Peninsula (Chhota Nagpur - 6B) (Rodgers & Panwar, 1988).

The upper stretch of the Ganga River falls under the Himalaya biogeographic zone (West Himalaya province - 2B) and is one of the richest areas in terms of habitat and species diversity (Rodgers & Panwar, 1988). This stretch comprises five forest types, viz., Alpine scrub (*Juniperus* sp., *Rhododendron campanulatum*); Sub-alpine forest (*Betula utilis*, *Abies spectabilis*, *Abies pindrow*); Himalayan Dry temperate forest (*Pinus gerardiana*, *Cedrus deodara*); Himalayan Moist temperate forest (*Quercus floribunda*, *Quercus leucotricophora*); and Sub-tropical pine forest (*Pinus roxburghii*) (Champion & Seth, 1968; Bahuguna et al., 2016). The snow leopard (*Panthera uncia*), Himalayan tahr (*Hemitragus jemlahicus*), musk deer (*Moschus* sp.), Himalayan black bear (*Ursus thibetanus laniger*) and Himalayan monal (*Lophophorus impejanus*) (Rodgers & Panwar, 1988) are some representative terrestrial species of this stretch. Representative aquatic faunal species of this biogeographic zone include the golden

mahaseer (*Tor putitora*), snow trout (*Schizothorax richardsonii*), mahaseer (*Tor tor*) (Bilgrami, 1991; Rao, 2001), Eurasian otter (*Lutra lutra*) and smooth-coated otter (*Lutrogale perspicillata*) (Hussain, 2002; Nawab & Hussain, 2012).

The middle stretch and most of the lower stretch of the Ganga River wind through the Gangetic Plain, which is one of the largest biogeographic zones in India, extending from the Yamuna River eastward across Uttar Pradesh, Bihar, Jharkhand and West Bengal (Sharma, 2005). This biogeographic zone is divided into the Upper Gangetic Plain (7A) and the Lower Gangetic Plain (7B) provinces (Rodgers et al., 2000). This biogeographic zone is considered one of the most fertile areas in the world, and most of the original vegetation has been converted into cropland. The region harbours two major forest types, Tropical Dry deciduous forest (*Dalbergia sissoo*, *Acacia nilotica*, *Madhuca indica*) and Tropical Moist deciduous forest (*Shorea robusta*, *Tectona grandis*, *Terminalia arjuna*) (Champion & Seth, 1968). Representative terrestrial species of this zone include the chital (*Axis axis*) and sambar (*Rusa unicolor*) (Rodgers & Panwar, 1988), and among the representative aquatic species are the Gangetic river dolphin (*Platanista gangetica*), smooth-coated otter (*Lutrogale perspicillata*), gharial (*Gavialis gangeticus*) and mugger (*Crocodylus palustris*) (Rodgers & Panwar, 1988). Examples of birds from these stretches of the Ganga River are





Sarus crane (*Antigone antigone*) and Indian skimmer (*Rynchops albicollis*) (Sundar et al., 2000; Sundar, 2004).

Before merging with the Bay of Bengal, the Ganga River flows through the East Coast (8B), mostly the Sundarbans delta (Rodgers & Panwar, 1988). The vegetation of this area falls under the swamp-and-littoral-forest type (Champion & Seth, 1968; Bahuguna et al., 2016) and is represented by species such as the sundari (*Heritiera fomes*), mangrove palm (*Nypa fruticans*), and tall-stilt mangrove (*Rhizophora apiculata*) (Champion & Seth, 1968; Gopal & Chauhan, 2006). The royal Bengal tiger (*Panthera tigris tigris*), barking deer (*Muntiacus muntjak*), leopard cat (*Prionailurus bengalensis*), fishing cat (*Prionailurus viverrina*) and lesser adjutant stork (*Leptotilos javanicus*) are the representative terrestrial fauna. Representative aquatic faunal species include the northern river terrapin (*Batagur baska*), estuarine crocodile (*Crocodylus porosus*), Indian flapshelled turtle (*Lissemys punctata*) and Indian softshelled turtle (*Trionyx gangeticus*) (Gopal & Chauhan, 2006).

Demography

2.6

The Ganga basin is one of the most densely populated river basins in the world, with a population of more than 500 million people (Misra, 2011). It supports around 43% of the Indian population, an average population density of over 700 individuals/km² and the density near the deltaic zone being over 1000 individuals/km² (Table 2.1) (GOI, 2011; Sanghi & Kaushal, 2014). There are 30 cities, 70 towns and thousands of villages located along the banks of the Ganga River (Behera et al., 2013). The major cities along the upper stretch of the Ganga River are Srinagar, Rishikesh and Haridwar (Uttarakhand). Major cities along the middle stretch within the state of Uttar Pradesh are Bijnor, Narora, Kanauj, Kanpur, Allahabad, and Varanasi. The lower stretch passing through the states of Uttar Pradesh (U.P), Bihar, Jharkhand and West Bengal has Mirzapur (U.P), Patna, Bhagalpur (Bihar), Sahibganj (Jharkhand), Farakka, Bahrapur, Serampore, Howrah and Kolkata (West Bengal) as major cities (CPCB, 2013).

Stretch	State	Persons	Density (persons/km ²)
Upper	Uttarakhand	10,116,752	189
Middle	Uttar Pradesh	199,581,477	829
Lower	Bihar	103,804,637	1102
	Jharkhand	1,150,038	414
	West Bengal	91,347,736	1029
Total		437,816,840	712.6

Table 2.1
Human density
along the Ganga
River (Source:
GOI, 2011)

3

STATUS OF AQUALIFE IN THE GANGA RIVER

Methodological Framework

3.1

A multistage approach was used to assess the status of the priority species (Table 3.1) in the Ganga River. The first stage consisted of a comprehensive literature review on the distribution, abundance and threats to the priority species. Information was collected and compiled using Google Scholar, 'Harzing's Publish or Perish and other online search engines. Mimeo reports, Fauna of British India, Fauna of India and other available literature from the British era were consulted. In the second stage, historical and current occurrence, abundance and distribution of the selected species were studied from the literature gathered. In the third stage, the results of the rapid biodiversity assessment of the mainstem Ganga River, carried out during pre-monsoon (April to June 2017) and post-monsoon (October and November 2017) periods, covering a distance of 2200 km, from Devprayag, in Uttarakhand, to Nurgur, in West Bengal, were incorporated in this synthesis. The rapid biodiversity assessment generated biological information, threat and depth profile of the Ganga River. This information on the aquatic diversity, their distribution pattern, threats, areas of concentration of threats and depth profile was collated and areas of higher conservation significance were identified by integrating these with the spatial data.



Class	Representative species	IUCN status	CITES	IWPA status
Himalaya (West Himalaya - 2B)				
Mammal	Asian small-clawed otter	Vulnerable	Appendix II	Schedule I
	Eurasian otter	Near Threatened	Appendix I	Schedule I
Reptile	Gharial	Critically Endangered	Appendix I	Schedule I
Fish	Golden mahaseer	Endangered	Not listed	Not listed
Gangetic Plain (Upper Gangetic Plain - 7A & Lower Gangetic Plain - 7B)				
Mammal	Gangetic river dolphin	Endangered	Appendix I	Schedule I
	Smooth-coated otter	Vulnerable	Appendix II	Schedule II
Reptile	Gharial	Critically Endangered	Appendix I	Schedule I
Bird	Sarus crane	Vulnerable	Appendix II	Schedule IV
Fish	Bata fish	Least Concern	Not listed	Not listed
Coast (East Coast - 8B)				
Mammal	Gangetic river dolphin	Endangered	Appendix I	Schedule I
Bird	Lesser adjutant	Vulnerable	Not listed	Not listed
Reptile	Northern river terrapin	Critically Endangered	Appendix II	Schedule I
	Estuarine crocodile	Least Concern	Appendix I	Schedule I
Fish	Hilsa	Least Concern	Not listed	Not listed

Table 3.1
Representative
aquatic fauna
from three
biogeographic
zones of the
Ganga River

The earliest historical records of the species of the Ganga River can be found in the fifth pillar edict of Emperor Asoka (Annandale, 1923; Dhammika, 1993), which prohibits killing or hunting of animal species, including the Gangetic river dolphin, turtles (possibly *Batagur dhongoka*) and bird species found in the rice fields in the Ganga River basin. The endangered Gangetic stingray was first described by Hamilton (1822) and thereafter by Annandale (1909), Bilgrami (1991) and Lakra et al. (2010a). Annexure I to VII provides an account of the species of the Ganga River on the basis of previous studies and the present study. Some of the species that were not reported from the Ganga River in the recent past, such as Siebold's smooth scaled water snake (*Enhydris sieboldii*) (Annexure V), and presence of Indian skimmer's nesting colonies between Kachhla and Varanasi, were recorded for the first time during this study. Gangetic river dolphins, birds, crocodilians and fishes are some of the most studied taxa of the Ganga River. The following sections describe the historical and present-day status of these and other priority aquatic species in the Ganga River.

Mammals

3.2

The major aquatic and semi-aquatic mammalian species found in the Ganga River include the Gangetic river dolphin and three species of otter, viz. smooth-coated otter, Asian small-clawed otter and Eurasian otter (Annexure I). In the Sundarban part of the lower Ganga there are frequent sightings of Irrawady dolphins (*Orcaella brevirostris*). These species are habitat specialists and have special adaptive features for living in aquatic systems. Being at the top of the food chain, they rapidly move between habitats, transport nutrients and energy between systems and stabilize the ecosystem. They also exert different top-down regulations upon trophic interactions and the coupling of ecosystems and habitats.





3.2.1



Gangetic river dolphin (*Platanista gangetica*)

The Gangetic river dolphin is one of the four exclusive freshwater dolphins in the world, whose distribution is restricted to the Indian subcontinent. It is an indicator of a healthy river ecosystem.

Historical Distribution

Prior to the initiation of water resource development activities in the Ganga River during the 19th century, the Gangetic river dolphin was distributed between 77°E and 88°E, throughout the Ganga, Brahmaputra/Meghna and Karnaphuli rivers and their tributaries in India, Nepal and Bangladesh (Sinha et al., 2000). In the Ganga River, their distribution ranged from Haridwar to the Sundarbans, and in the Yamuna River, the species was reported up to Delhi (Anderson, 1878; Sinha et al., 2010) (Figure 3.1). During the late 19th century, about 10,000 Gangetic river dolphins were estimated to be thriving in the Ganga River and its tributaries (Anderson, 1878); however, Sinha and Kannan (2014) estimated their population to be 3526 individuals during the early 2000s (Box 1).

Gangetic river dolphins are now restricted to the mainstem Ganga River and its large tributaries viz. the Ramganga, Yamuna, Gomti, Ghaghara, Rapti, Son, Gandak and Kosi (Sinha et al., 2000) (Box 2). They have disappeared from the Haridwar stretch of the Ganga River, and the Bijnor Barrage, 100 km downstream of Haridwar, now restricts their population (Sinha et al., 2000). The population in the Yamuna River is now only restricted to the confluence of the Chambal and Yamuna rivers, near Etawah, almost 490 km downstream from Delhi (Sinha & Sharma, 2003) (Figure 3.1). It has also become locally extinct in most of the small tributaries or is sighted only during the rainy season (Sinha & Sharma, 2003). During the past 20 years, many scholars have studied the Gangetic river dolphin, but most of the studies were done in patches and at different temporal scales. Table 3.2 provides information on the encounter rate of the Gangetic river dolphin on the basis of earlier studies. According to these studies, encounter rate was highest at Vikramshila Gangetic Dolphin Sanctuary (1.8/km) and lowest between Bijnor and Narora (0.36/km). While in the present study, encounter rate was highest between Allahabad to Buxar, followed by the Vikramshila Gangetic Dolphin Sanctuary. The abundance of the Gangetic river dolphin was noted to be 179 and 270 in the mid and peak dry seasons, respectively (Kelkar et al., 2010), in the Vikramshila Gangetic Dolphin Sanctuary. The loss of habitat connectivity, hydrology alteration, overexploitation of the prey-base and hunting moved the species from the Vulnerable category to the Endangered category of the IUCN Red List in 2004 (IUCN, 2016).

Figure 3.1
Historic and
current
distribution of
Gangetic river
dolphin in the
Ganga River





Table 3.2
Encounter rate of
Gangetic river
dolphin in the
main stem of the
Ganga River

River stretch	Encounter rate (individuals/km)		
	Previous studies		Present study (2017)
Bijnor and Narora	0.36	Behera (1995)	0.21
Narora and Kanpur	0	Sinha (1999)	0.03
Kanpur and Allahabad	0.49	IIT (2012)	0.24
Allahabad to Buxar	0.48	Sinha (1999)	0.77
Buxar to Maniharighat	1.62	Sinha et al. (2010)	0.36
Vikramshila Gangetic Dolphin Sanctuary	1.8	Choudhary et al. (2006)	0.65
Maniharighat to Farraka	1.64	Sinha (1999)	0.22
Farakka Feeder Canal	0.55	Sinha et al. (2000)	0.10

Present Distribution

During the rapid biodiversity assessment, a total of 648 sightings of the Gangetic river dolphins were made from Bijnor (Uttar Pradesh) to Nurpur (West Bengal). Gangetic river dolphins were most abundant from Brijghat to Narora, upstream of Kanpur, from Bhitura to Ghazipur and between Chappra and Kahalgaon. Smaller and scattered groups were observed between Sahibganj and Rajmahal and from Baharampur to Barrackpore (Figure 3.2). Groups of Gangetic river dolphin mostly congregated at pools with a depth of >4 m and at meanders with steep clayey banks.


Figure 3.2
Distribution of
Gangetic river
dolphin in the
Ganga River, on
the basis of the
rapid biodiversity
assessment



Loss of habitat connectivity, reduced flow regime, reduced prey base and water pollution restricts homogeneous distribution of the species and, as a result, this species is now restricted to deep pools, particularly from downstream of Kanpur. Optimal habitats are present in stretches from Brijghat to Narora, downstream of Kanpur from Allahabad to Bhagalpur, from Gopal Ghat to Palasi, from Ram Nagar to Nabadwip and downstream of Nabadwip to Howrah-Kolkata. It was observed that fishing using gill nets, overfishing and a reduced flow regime are major threats to the species.

From the literature review and rapid biodiversity assessment, it was inferred that the populations of the Gangetic river dolphin in the Ganga River are highly fragmented and therefore undergoing loss of genetic diversity. The population is concentrated in the middle stretch of the river, where dispersal constraints such as dams and barrages as well as shallow areas are minimal.

BOX 1



Decline in the population of the Gangetic river dolphin from 1878 to 2010 on the basis of estimates provided by Anderson (1878); Jones (1982); and Sinha and Kannan (2014)

BOX 2

River	River stretch	Encounter rate (individuals/km)	Reference
Chambal	Batesura and the confluence of the Yamuna River	0.15	Singh and Sharma (1985)
Son	Between Bicchi, in Madhya Pradesh, and Banjari	0.076	Sinha et al. (2000)
Kosi	Between Kosi Barrage and Kursela	0.42	Sinha and Sharma (2003)
Gandak	Gandak Barrage to Gandak-Ganges confluence at Patna	0.8	Choudhary et al. (2012)
Ken	Confluence with Yamuna to Sindhan Kalan village	0.26	Sinha et al. (2000)
Betwa	Confluence with Yamuna to Orai	0.07	Sinha et al. (2000)
Sind	Confluence with Yamuna	0.045	Sinha et al. (2000)

Encounter rates of the Gangetic river dolphin in the tributaries of the Ganga River



3.2.2



Otters

Otters form a well-marked group, representing the family Mustelidae. They are mammals adapted for a semi-aquatic life. They are the principal predators of aquatic environments and indicators of the health of wetland or riverine ecosystems as they are sensitive to degradation along the food chain and habitat (Erlinge, 1972; Kruuk et al., 1993; Mason & MacDonald, 1987). Otters are the top predators of aquatic ecosystems and shape the faunal species composition of the ecosystems they inhabit.

Of the 13 species of otters worldwide, five species occur in Asia, and three of them are found in India; including the Ganga River basin. Otters usually live in low densities, are elusive and are largely nocturnal, thus making direct sightings difficult (Hussain, 2002). Hence, most of the studies on otter distributions are based on occasional sightings supplemented by indirect evidence of presence.

Historical Distribution

The earliest authentic reports of the occurrence of otters in the tributaries of the Ganga River in the Terai, Garhwal and Kumaun areas (lower Himalaya) were given by Atkinson (1882), Blanford (1881) and Hinton and Fry (1923). Historically, the smooth-coated otter (*Lutrogale perspicillata*) was distributed from the Himalaya southward to Karnataka and eastward to Burma (Pocock, 1941). It was also reported from the Terai region during the late 19th century; from the Yamuna River; and from Salt Lake and the Sundarbans, in West Bengal (Atkinson, 1882, 1974) (Table 3.3) (Figure 3.3).

In the last decade the smooth-coated otter has been reported from the Upper Ganga (Alaknanda), Ramganga, Mandal, Sonanadi and Palain rivers in the Upper Gangetic Plain (Nawab & Hussain, 2012); Hastinapur Wildlife Sanctuary (Khan et al., 2014); the Bijnor–Narora stretch of the Ganga River (Bashir et al., 2012); and Vikramshila Gangetic Dolphin Sanctuary (Choudhary et al., 2006).

The Eurasian otter (*Lutra lutra*) was historically distributed north of the Ganga River, extending throughout the Himalaya and in the Upper and Lower Gangetic plains up to Kolkata (Sclater, 1891). It was also distributed in the eastern coastal region of India up to Sri Lanka (Figure 3.4). Atkinson (1882) reported that the species was present throughout the Terai and in all the larger streams along the foothills of the Himalaya, ascending around 60 km. He also noted groups of 5 to 12 individuals in the Ramganga River in the Patli Dun and around 20 individuals in the Suswa River in Dehra Dun. In the recent past, Hussain (2002) reported signs of *L. lutra* from the Ramganga River, in the upper Gangetic basin.

Historically, the distribution of the third species of otter, viz. the Asian small-clawed otter (*Aonyx cinereus*), has been discontinuous (Blanford, 1881). It was reported from the smaller streams occurring in the lower elevations of the Himalayas and lower Bengal, being common in the Sundarbans, Assam and Burma. Historically, the species was observed in the Sarda River at Barmdeo and major tributaries of the upper Ganga, Pindar rivers near Karnaprayag; in the Nandakini, above Nandprayag; and in Bhilang (Jerdon, 1874) (Figure 3.5). These otters are diurnal animals occurring in small groups and are found in streams free of human disturbance. In recent years, its presence has been confirmed from the Sundarbans and the Eastern ghats of Odisha (Mohapatra et al., 2014).

A survey conducted during 1993–1995 from Rishikesh to Kanpur reported absence of otters in the Ganga River (Rao, 2001). In more recent studies, their presence has only been reported from the lower Himalayan and Terai regions at smaller tributaries of the Ganga River in the Alaknanda valley, in PAs (Hussain, 2002). It is evident from the literature that otters were present throughout the length of the Ganga River. These studies also provided insights into the habitat and behaviour of the otters. However, these studies were conducted at different time scales and were limited in their geographical extent and were not specific to the Ganga River. Hence the population trends of otter species cannot be deduced from these studies. Extensive studies are therefore needed to assess the population trends of otter species in the Ganga River and its basin.

Species	Scientific name	IUCN status	Uttarakhand	Uttar Pradesh	Bihar	Jharkhand	W. Bengal
Smooth-coated otter	<i>Lutrogale perspicillata</i>	VU	H + P	H + P	H + P	H + P	H + P
Eurasian otter	<i>Lutra lutra</i>	NT	H + P	-	-	-	-
Asian small-clawed otter	<i>Aonyx cinereus</i>	VU	H + P	-	-	-	H

H - Historically distributed; P - Presently distributed

Table 3.3
Historical and current distribution of otters in the Ganga River states

Figure 3.3
Distribution of
smooth-coated
otter (*Lutrogale
perspicillata*),
adapted from
IUCN




Figure 3.4
Distribution of
Eurasian otter
(*Lutra lutra*),
adapted from
IUCN






Figure 3.6
Distribution of
smooth-coated
otter in the
Ganga River, on
the basis of the
rapid biodiversity
assessment



Avifauna


3.3

India is one of the global hotspots for birds, with over 1340 bird species (13% of the world's species) recorded from the country (Manakadan & Pittie, 2001). Falling within the Central Asian Flyway, the area is utilized by 307 species of migratory waterbirds, most of them coming from Central and North Asia, at the beginning of the winter season. Of the 307 migratory species, two are Critically Endangered, five are Endangered and 13 species are Vulnerable. Out of the remaining 287 species, 10 species are Near Threatened and 277 species are Least Concern as per the IUCN Red List.

Waterbirds maintain the diversity of other organisms, control pests, are effective bioindicators of ecological conditions, and act as sentinels of potential disease outbreaks. They also provide important provisioning and cultural services to humans.


The Ganga River basin has several seasonally flooded wetlands, permanent lakes and marshes, which serve as major migratory waterbird habitats. Some of these have been declared as PAs, for e.g., Jhilmil Conservation Reserve in Uttarakhand, Surhataal Wildlife Sanctuary in Uttar Pradesh, Kusheshwar Asthan Bird Sanctuary in Bihar and Udhwa Bird Sanctuary in Jharkhand. The Ganga River, with its mosaic of habitats, supports 140 birds, including 129 waterbirds and obligate bird species (Annexure II). Apart from resident and migratory species, some iconic and globally threatened birds such as the black-bellied tern (*Sterna acuticauda*), Indian skimmer (*Rynchops albigollis*), Sarus crane (*Antigone antigone*) and river lapwing (*Vanellus duvaucelii*) also breed on the islands, sandbars and banks of the Ganga River. The literature suggests that the greatest species richness of the Ganga River is in Uttar Pradesh (Figure 3.7), which has 135 bird species. The least number of species is reported from Jharkhand, in the lower stretch. Very few systematic studies have been carried out on the distribution of waterbirds along the Ganga River. For example, Rao (2001) reported 46 species from Rishikesh to Kanpur, covering the upper and middle stretches. In the middle stretch, Rahmani (1981) recorded 120 species at the Narora Barrage and Bashir et al. (2012) recorded 55 species from Bijnor to Narora. In the lower stretch, Bilgrami (1991) recorded 23 bird species from Munger, in Bihar, to Farakka, in West Bengal.

Figure 3.7
Riverine and wetland avifauna richness in the Ganga River states



During the assessment of waterbird diversity, 87 species of waterbirds and five wetland obligate species belonging to 22 families were recorded from six reservoirs along the Ganga River (Figure 3.8). Among these six reservoirs, 65 bird species were recorded from Bijnor barrage followed by 62 species from Narora, 41 species from Rishikesh, 37 species from Kanpur, 30 species from Farakka and 27 species from Bhimgoda barrage in Haridwar (Figure 3.9). Of the total species observed in the reservoirs, the members of the Family Anatidae (ducks and geese) dominated (18 species) followed by Scolopacidae (waders) and Ardeidae (egrets and herons) (Figure 3.10). Of the total 92 species recorded during the assessment along the reservoirs, 17 species belonged to Rare, Endangered and Threatened (RET) categories of IUCN, of which one species each were Critically Endangered and Endangered, six were Vulnerable and 10 species were Near Threatened.

Figure 3.8
Migratory birds monitoring sites






Figure 3.9
Number of waterbird species observed at six reservoirs along the Ganga River




Figure 3.10
Number of species belonging to different Families of birds seen at six reservoirs along the Ganga River

Focused studies were also carried out in select stretches of the Ganga River as part of the rapid biodiversity assessment to supplement the available information on the distribution and abundance of waterbirds along the Ganga River. For example, Kumar (2017b) carried out an intensive study in the middle stretch of the Ganga River near Allahabad, covering 25 km upstream and downstream, and encountered 7153 waterbirds belonging to 41 species and 12 families. Among them, the river lapwing (*Vanellus duvaucelii*), ruddy shelduck (*Tadorna ferruginea*), Temminck's stint (*Calidris temminckii*), black-headed gull (*Larus ridibundus*), brown-headed gull (*Larus brunnicephalus*), pied avocet (*Recurvirostra avosetta*) and spotted redshank (*Tringa erythropus*) were the most abundant species due to the habitat characteristics of the Ganga River in the study area. In another study conducted by Vasudeva (2017) at Narora, a Ramsar site, 83 bird species were encountered, which represented 43 families. Passeriformes was the dominant order, with 15 families, followed by Charadriiformes (six families) and Coraciiformes (five families). During the rapid biodiversity assessment, 104 species of waterbirds and wetland obligate species belonging to 26 families were recorded of which 62% were resident and 38% were migratory species. Of the total species observed, the members of the Family Anatidae (ducks and geese) dominated (26 species) followed by Scolopacidae (waders) and Ardeidae (egrets and herons). Of the 104 species recorded during the assessment, 12 species belonged to Rare, Endangered and Threatened (RET) categories, of which six species were Critically Endangered, three species were Endangered, three were Vulnerable and 11 species were Near Threatened. The migratory waterbirds use sandy islands and sand banks of the Ganga River for roosting and the resident waterbirds such as Indian skimmer, little pratincole and river terns for breeding.



3.3.1




Black-bellied tern (*Sterna acuticauda*)

The black-bellied tern is a small, riverine, colonial island-nesting species of the family Laridae and is a resident breeder of the Middle and Lower Gangetic plains (7A & 7B). This species has been recorded from India, Pakistan, Nepal, Bangladesh, Myanmar, Thailand, Laos, Cambodia, Vietnam and southern China (Figure 3.11). It is listed as Endangered in the IUCN Red List, as it is almost extinct or rapidly declining all across its range owing to destruction of its breeding habitat, collection of eggs for food, overfishing and the flooding of nests, often caused by dams.

Historical records of the species (1990–2016) are presented in Table 3.4. Behera (1995) reported the presence of the species from Bijnor to Narora during 1995 in Uttar Pradesh. A continuous survey conducted by Dey et al. (2014) during 1999–2014 found the species regularly at Vikramshila Gangetic Dolphin Sanctuary, in Bihar. The species has not been reported from Uttarakhand and Jharkhand. In 1994, the species was reported from the Farakka Barrage (Jha, 2006). Extensive studies are needed to assess the population trend of this species in the Ganga River and its basin.

During the rapid biodiversity assessment, a total of 198 individuals were recorded from Bijnor to Farakka at 65 locations. The greatest abundance was found between Narora and Kannauj, followed by the stretch from Sahibganj to Rajmahal and upstream of Ghazipur (Figure 3.19).

Figure 3.11
Distribution of
black-bellied tern
(*Sterna*
acuticauda)





Indian sarus crane (*Antigone antigone*)



3.3.2

The largest crane in the Indian subcontinent, the sarus is a resident breeder of the Gangetic floodplains. It inhabits the natural wetlands, small seasonal marshes, floodplains, fallow and cultivated lands and paddy fields. The global population of the species is showing a declining trend and is listed as Vulnerable in the IUCN Red List. The Indian population is also suspected to have decreased owing to the loss and degradation of wetlands and ingestion of pesticides used in agricultural fields (Sundar et al., 2000; BirdLife International, 2016).


Historically this species inhabited northern central and eastern India, including West Bengal, Nepal and Pakistan (Figure 3.12). The number of individuals in India, Nepal and Pakistan is estimated to be 8000–10,000. In India, the species was distributed from Jammu and Kashmir, in the north, to Chandrapur, in Maharashtra, in the south, and from Gujarat, in the west, to West Bengal and Assam, in the east (Choudhury, 1998; Kaur et al., 2002).

More recent studies reveal that Uttar Pradesh still remains the stronghold of the species, with a population estimated at over 6000 individuals (Sundar, 2008). Bashir et al. (2012) reported encounter rates of 0.285 ± 0.065 during boat surveys and 0.192 ± 0.076 during bank searches in 2007 along the Bijnor–Narora stretch of the Ganga River. Forty-five individuals of the species were reported (2006–2007) from Farrukhabad, Shahjahanpur and Hardoi districts, of Uttar Pradesh, along the Ganga River (Maheswaran et al., 2010). A state-wide survey conducted in Uttar Pradesh during 2010 (Jha & McKinley, 2014) found 6927 individual along the Gangetic Plain.

The sarus does not seem to be a common bird in the Lower Gangetic Plain of West Bengal any more, and its population in West Bengal and Bihar remains unknown (Sundar et al., 2000). The species has not been reported from Uttarakhand. Extensive studies are needed to assess the population trend of this species in the Ganga River and its basin. During the rapid biodiversity assessment, 1362 sarus cranes were observed between Brijghat and Farakka, with the greatest abundance near Bhitura, followed by Kanpur, Kannauj and Farrukhabad (Figure 3.16). In the site-focused study conducted by Vasudeva (2017), 12 individuals of the species were encountered from Karnwas to Seensai, near Narora, during January–May 2017.



Figure 3.12.
Distribution of
Indian sarus
crane (*Antigone
antigone*)



Indian skimmer
(*Rynchops albicollis*)

The Indian skimmer is one of the three skimmer species of the world and the only one that is found in the Indian subcontinent. This predatory fish-eating bird has a unique adaptive feature: uneven bills. The lower mandible is longer than the upper.

This species has been reported from India, Pakistan, Bangladesh and Myanmar and is a rare visitor to Nepal (Figure 3.13). It is a globally threatened bird listed as Vulnerable in the IUCN Red List, with an estimated Indian population of 2500 individuals (BirdLife International, 2017). Its range has become increasingly fragmented in recent decades. It is found in Uttar Pradesh, Bihar, Jharkhand and West Bengal along the Ganga River basin and in the Indus River in Kashmir. The species has a widespread distribution in winter and is found in Karwar, on the western coast, and in Chennai and Pondicherry, on the eastern coast. Breeding colonies are known from the Chambal River. Historical records of the species (1990–2016) are presented in Table 3.5. The species has not been reported from Uttarakhand and Jharkhand. Early observations from 1991 were obtained from the Lower Gangetic Plain of West Bengal (Jha, 2006), and recent sightings from Uttar Pradesh have been reported by Mishra et al. (2016).

During the rapid biodiversity assessment, 411 individuals of Indian skimmers were recorded. The Indian skimmer were most frequently encountered between Bijnor and Ghazipur. Seven nesting colonies of the Indian skimmer were recorded between Kachhla and Varanasi. Maximum number of nesting colonies were encountered between Allahabad and Varanasi. The species was most abundant near Varanasi, followed by Allahabad, Mirzapur and Narora. (Figure 3.16). In earlier site-specific studies, Kumar (2017b) encountered 83 birds at a nesting island near Allahabad and 76 individuals of the species were encountered from Karnwas to Seensai, near Narora, during January–May 2017 (Vasudeva, 2017).




Figure 3.13
Distribution of
Indian skimmer
(*Rynchops*
albicollis)



3.3.4




River tern (*Sterna aurantia*)

A resident breeder in India, the river tern is a common riverine species and an island-nesting bird. This species occur across a wide range in southern Asia and southeast Asia, including Pakistan, Nepal, Bhutan, Bangladesh, Myanmar, Thailand, Laos, Cambodia, Vietnam and southern China (del Hoyo et al., 1996) (Figure 3.14). Globally, the species is showing a declining trend in its population and is listed as Near Threatened in the IUCN Red List. The population in India is estimated to be >50,000 individuals (Delany & Scott, 2006). The nesting areas are vulnerable to flooding, predation and disturbance (del Hoyo et al., 1996), and abrupt declines have occurred in parts of southeast Asia.

Bashir et al. (2012) reported an encounter rate of 0.272 ± 0.059 for the species during a boat survey conducted in 2007, along the Bijnor–Narora stretch of the Ganga River. In a continuous survey conducted by Dey et al. (2014) from 1999 to 2014, 10–20 individuals were regularly sighted at Vikramshila Gangetic Dolphin Sanctuary. The species is reported occasionally from the Lower Gangetic Plain and the Coastal Zone of West Bengal. No literature pertaining to the species is available for Uttarakhand and Jharkhand.

During the rapid biodiversity assessment, 197 individuals of the species were encountered between Bijnor and Farakka. Major congregations were observed near Farukhabad and Bijnor and at Brijghat (Figure 3.19). In a site-focused study conducted by Vasudeva (2017), 72 individuals of the species were encountered from Karnwas to Seensai, near Narora, during January–May 2017.

Figure 3.14
Distribution of
river tern (*Sterna*
aurantia)





River lapwing (*Vanellus duvaucelii*)



3.3.5

The river lapwing is a resident breeding riverine species in India. The species occurs in parts of the Indian subcontinent, including Nepal, Bhutan, Bangladesh, Myanmar, Thailand, Cambodia, Laos and Vietnam; southern China; and southeast Asia (Chandler, 2009) (Figure 3.15). Listed as Near Threatened in the IUCN Red List, it is showing moderately rapid overall population decline owing to the impacts of human pressures on riverine ecosystems and multiple dam construction projects (BirdLife International, 2016).

Few systematic surveys have been conducted to assess the population of the species in the Gangetic Plains of Uttar Pradesh, Jharkhand and West Bengal in the past decades, and the species is only mentioned in checklists from these areas (State Fauna Series, Zoological Survey of India). During a continuous survey conducted by Dey et al. (2014) from 1999 to 2014, the species was found at Vikramshila Gangetic Dolphin Sanctuary. Kumar (2017b) encountered 568 individuals of the species along a 50 km stretch near Allahabad from December 2016 to January 2017.

River lapwings were very common and widely distributed throughout the stretch of the Ganga River that was surveyed. A total of 2923 individuals of this species were observed during the assessment. The greatest abundance was found near Kanpur, followed by Bhitura, Farukhabad, Kannauj, Manikpur, Ghazipur, Kachla, Mirzapur and Bijnor (Figure 3.19). In a site-focused study conducted by Vasudeva (2017), 210 individuals of the species were encountered from Karnwas to Seensai, near Narora, during January–May 2017.




Figure 3.15
Distribution of
river lapwing
(*Vanellus*
duvaucelii)



3.3.6





Painted stork (*Mycteria leucocephala*)

This threatened species occurs widely in the Indian subcontinent; however, it has become rare in Myanmar, Thailand, Laos and Vietnam and is possibly extinct in China (Figure 3.16). There are an estimated 15,000–25000 individuals in south Asia (Wetlands International, 2012) and fewer than 10,000 in southeast Asia (Perennou et al., 1994), with populations declining throughout the range owing to hunting, wetland drainage and pollution. Therefore, the species is listed as Near Threatened in the IUCN Red List.

The species has been recorded in the Himalayan zone of Pauri-Garhwal District and Rajaji National Park, in Uttarakhand (Islam & Rahmani, 2008). During a continuous survey conducted by Dey et al. (2014) from 1999 to 2014, the species was found at Vikramshila Gangetic Dolphin Sanctuary (Table 3.6). Bashir et al. (2012) recorded the species from the Bijnor–Narora stretch of the Ganga River. A large flock was observed by Mishra et al. (2016) at Rae Bareilly, Uttar Pradesh. The species is occasionally reported from the Lower Gangetic Plain of West Bengal. However, apart from a few checklists, there is no literature on the abundance and distribution of the species.

Figure 3.16
Distribution of
painted stork
(*Mycteria*
leucocephala)





Great thick-knee (*Esacus recurvirostris*)



3.3.7

The species is distributed across India, Pakistan, Sri Lanka, Nepal, Bhutan, Bangladesh, Myanmar, Thailand, Iran, Laos, Cambodia, Vietnam and southern China (Figure 3.17). Precipitous declines have occurred in parts of Southeast Asia. India probably holds the bulk of the population now. Increasing numbers of dams and impoundments on large rivers in the range of the species threaten to alter flow regimes and inundate nesting habitats downstream. This has resulted in declines in the population of the species and therefore is listed as Near Threatened in the IUCN Red List.

According to checklist, the species is present in Uttar Pradesh, Bihar and West Bengal. Seven individuals of this species were reported from mud flats and sandbanks of the Sundarbans (Zöckler et al., 2005). During the rapid biodiversity assessment, 197 individuals were recorded. Maximum sightings were from Kachhla – Farrukhabad stretch, followed by Brijghat – Narora, and Narora – Kachhla stretches (Figure 3.19).




Figure 3.17
Distribution of
great thick-knee
(*Esacus
recurvirostris*)



3.3.8



Little pratincole (*Glareola lactea*)

This species has an extremely large range, distributed in India, Afghanistan, Bangladesh, Bhutan, Cambodia, China, India, Laos, Myanmar, Nepal, Oman, Pakistan, Sri Lanka and Thailand (Figure 3.18). The global population size and trend are currently unknown (BirdLife International, 2016) but the species is listed as Least Concern in the IUCN Red List. According to checklists, the species is present in Uttar Pradesh, Bihar and West Bengal. Apart from this, there is very little in the literature on the abundance and distribution of this species with respect to the Ganga River and its floodplain.

A total of 3339 individuals of this species were encountered from Makdumpur to Nabadwip during the rapid biodiversity assessment. The species was most abundant between Narora and Kanpur, followed by the stretch upstream of Bhagalpur and from Bhitura to Ghazipur (Figure 3.19). In site specific studies, Kumar (2017b) encountered 440 individuals of this species along a 50 km stretch near Allahabad district from December 2016 to January 2017 and a total of 166 individuals were encountered from Karnwas to Seensai, near Narora, during January–May 2017 (Vasudeva, 2017).

Figure 3.18
Distribution of
little pratincole
(*Glareola*
lactea)




Table 3.4
Records of black-bellied tern in the Ganga River and its floodplain

Year	Ganga River state								
	Uttar Pradesh			Bihar			West Bengal		
	No.	Region	Reference	No.	Region	Reference	No.	Region	Reference
1994	No literature available			No literature available			23	Farakka Barrage	Jha (2006)
1995	NA	Bijnor-Narora	Behera (1995)						
1999 to 2013	No literature available			15-20	50-65 km of Vikramshila Gangetic Dolphin Sanctuary	Dey et al. (2014)			
2014				30-40					

Table 3.5
Records of Indian skimmer in the Ganga River and its floodplain

Year	Uttar Pradesh			Bihar			West Bengal		
	No.	Region	Reference	No.	Region	Reference	No.	Region	Reference
1991	No literature available			No literature available			7	Manikchak Ghat	
1995							4	Near Gopalpur	
1997							9	Near Farakka Barrage	
1998							23		
1999							20		Jha (2006)
2004 2005				Breeding	Barari, Bhagalpur	Dey et al. (2014)	No literature available		
2007				No literature available			9	Chittaranjan Bara Dam, Burdwan District	Sharma (2008)
2008				Frequently observed	Bhagalpur	Dey et al. (2014)	6	Nidoya and Rajar Char, Purbasthali-Ganges Islets, Burdwan District	
2011				2			No literature available		
2015 2016	148	Gegaso, Dalmau, Unchahar, Raebareilly District	Mishra et al. (2016)	No literature available					

Table 3.6
Records of
painted stork in
the Ganga River
and its floodplain

Year	Ganga River States					
	Uttar Pradesh			Bihar		
	No.	Region	Reference	No.	Region	Reference
1999-2006	No literature available			40-50	Lodipur and Bhagalpur	Dey et al. (2014)
2007	0.063±0.011, boat survey 0.041±0.014 boat survey	Bijnor-Narora	Bashir et al. (2012)			
2008-2013	No literature available					
2014	No literature available			14	Bhagalpur	
2015 2016	187	Rae Bareilly	Mishra et al. (2016)	No literature available		

A total of 104 waterbirds and wetland obligate species, belonging to 26 families were sighted during the rapid biodiversity assessment. Of the total species observed, the members of the Family Anatidae (ducks and geese) dominated (26 species) followed by Scolopacidae (waders) and Ardeidae (egrets and herons). The Indian skimmer, sarus crane, black-bellied tern, river tern, river lapwing, great thick-knee and little pratincole are some of the resident breeding birds of the Ganga River that were most frequently encountered during the rapid biodiversity assessment (Figure 3.19). Indian skimmers were most frequently encountered between Bijnor and Ghazipur. Nesting colonies of the Indian skimmer were recorded between Kachhla and Farrukhabad, at Bithura Ghat, at Allahabad and from Mirzapur to Varanasi. The species was most abundant near Varanasi, followed by Allahabad, Mirzapur and Narora. Sarus cranes were observed between Brijghat and Mirzapur, with maximum abundance near Bhitura, followed by Kanpur, Kannauj and Farukhabad. River terns were recorded between Bijnor and Varanasi. Major congregations were observed near Farukhabad and Bijnor, followed by Brijghat. River lapwings were very common and widely distributed throughout the stretch of the Ganga River that was surveyed. The greatest abundance was found near Kanpur, followed by Bhitura, Farukhabad, Kannauj, Manikpur, Ghazipur, Kachla, Mirzapur and Bijnor. The great thick-knee was encountered from Bijnor to Kannauj. The greatest abundance was recorded near Farukhabad, followed by Kachla, Kannauj, Narora and Brijghat. Black-bellied terns were sighted from Bijnor to Kannauj. The greatest abundance was found between Narora and Kannauj, followed by the stretch from Sahibganj to Rajmahal and upstream of Ghazipur. Little pratincoles were encountered from Makdumpur to Nabadwip. The species was most abundant between Narora and Kanpur, followed by the stretch upstream of Bhagalpur and Bhitura to Ghazipur.

However, the population trends of these birds could not be estimated due to the lack of temporal-scale information. More extensive studies on the distribution and ecology of these birds are essential for planning their conservation.



Figure 3.19
Distribution of
resident breeding
birds, along the
Ganga River, on
the basis of the
rapid biodiversity
assessment





Reptiles

3.4

Crocodylians

3.4.1

Crocodylians are represented by 24 living species (Martin, 2008). In India, mugger (*Crocodylus palustris*) and estuarine crocodile (*C. porosus*) represent the family Crocodylidae, and gharial (*Gavialis gangeticus*) represents the family Gavialidae. These species are distributed in the Ganga River and its major tributaries (Annexure III).

Presence of crocodiles increases yield of fish and improves health of fish stock, since crocodiles eat ailing fish. Fishes common are also eaten by crocodiles and thus they maintain balance in fish species assemblage (van der Ploeg et al., 2011).



Gharial (*Gavialis gangeticus*)



3.4.1.1

Historically, gharials were restricted to the perennial Himalayan river systems in the northern part of the Indian subcontinent (Boulenger, 1890; Smith, 1931; Wright, 1982). Indian rivers that formerly supported large populations of the gharial include the Ramganga, Ghagra, Gandak, Kosi and Girwa (Whitaker, 1979). The species were on the verge of extinction in India by the mid-1970s, and the population had plummeted to fewer than 200 individuals. To conserve this species a captive breeding programme was initiated in India (Bustard, 1980, 1984) during 1975. As part of this programme captive reared crocodiles were reintroduced to augment wild populations, especially in the Chambal River, at the National Chambal Sanctuary, in the Girwa River at Katerniaghat Wildlife Sanctuary (Hussain, 2009). The restocking programme, along with legal protection, increased the breeding population in the Chambal River in the Ganga River basin.

Recently, gharials have been reported from the main stem Ganga River between Bhagalpur and Sultanganj (Nawab et al., 2016) and from its tributaries viz., Ramganga at Corbett Tiger Reserve (Chowfin & Leslie, 2013); Son River at Son Gharial Sanctuary; Girwa River at Katerniaghat Wildlife Sanctuary; and Chambal River, at National Chambal Sanctuary (Figure 3.20). The Chambal River has the largest subpopulation of breeding gharials in the wild with around 48% of the total population (IUCN, 2009).




Figure 3.20
Distribution of
gharials (*Gavialis*
gangeticus)



3.4.1.2




Mugger (*Crocodylus palustris*)

The mugger is one of the most widely distributed species owing to its highly adaptable nature. Throughout India, this species survives in stagnant water, jheels, reservoirs, tanks and rivers with placid currents (Boulenger, 1890). In India, it is present in 15 states including much of the Ganga River drainage. Significant populations occur in the lower Ganga (Bihar and Jharkhand), Chambal River (Rajasthan and Madhya Pradesh) (Sharma & Singh, 2015) and Gujarat (Figure 3.21). In Uttarakhand, the species has been reported from the Ramganga River and the reservoir area in Corbett Tiger Reserve (Nawab, 2007), Rajaji National Park and parts of Haridwar and Lansdowne forest divisions (Joshi, 2013).

In the Ganga River basin, this species has been reported from Hastinapur Wildlife Sanctuary and the Ramsar site between Garhmukteshwar and Narora (Nawab et al., 2016), from the Banas and Tapi rivers (McCann, 1940) and from the states of Madhya Pradesh, Uttar Pradesh, Bihar and West Bengal (Whitaker and Daniel, 1980). Table 3.7 provides a comparative account of the gharial and mugger populations in the Ganga River and its tributaries on the basis of literature. It is evident that these species have been studied mostly in the tributaries of the Ganga River, rather than the mainstem.

Figure 3.21
Distribution of
mugger
(*Crocodylus*
palustris)






Estuarine crocodile (*Crocodylus porosus*)



3.4.1.3

Historically, the estuarine crocodile (*Crocodylus porosus*) was distributed along the coast of south Asia (Figure 3.22). The rarity of the species in India was apparent by the late 1960s (Daniel, 1970) as the population became extinct in the states of Kerala, Tamil Nadu and Andhra Pradesh (de Vos, 1984; FAO, 1974). A small population remained in the Brahmani-Baitarani Delta in Odisha, the Gangetic Delta of the Sundarbans in West Bengal, and Andaman Islands (FAO, 1974). The species has become locally extinct over large parts of its range, with viable populations only occurring in PAs (Santiapillai & Silva, 2001).

Figure 3.22
Distribution of
estuarine
crocodile
(*Crocodylus*
porosus)



As part of the Crocodile Conservation Project (GOI, 1975), *C. porosus* were released in Bhitarkanika Wildlife Sanctuary in 1975 and in the Sundarbans until 1982 to rejuvenate the existing population. Bustard and Choudhury (1981) and later Kar and Bustard (1986) reported a decline in the population of the Sundarbans. During 1998, the total Indian population of *C. porosus* was 170–330 (Sharma, 1998). Recently, Saha et al. (2011) reported the presence of a good population of *C. porosus* in the Sundarban estuary. Extensive studies are needed to assess the population trend of this species in the Ganga River and its basin.

Present Distribution

During the rapid biodiversity assessment, individual gharials were sighted in Rajaji National Park downstream of the Bhimgoda Barrage, and groups of gharials were sighted between Bijnor and Brijghat. Individual muggers were sighted near Bijnor and at Malda, upstream of the Farakka Barrage (Figure 3.23). The gharials sighted between Bijnor and Brijghat were part of the released population restocked in Hastinapur Wildlife Sanctuary. Suitable habitats in terms of deep-water stretches, lateral connectivity, nesting sandbars and islands were found at certain locations in the Ganga River, viz. from Bijnor to Narora and from Farrukhabad to Varanasi.

Figure 3.23
Distribution of
gharials and
muggers in the
Ganga River, on
the basis of the
rapid biodiversity
assessment



Crocodylians are highly vulnerable to anthropogenic habitat alteration and are therefore conservation-dependent species. However, most of the crocodilian habitats along the Ganga River are not protected. The absence of habitat features such as deep waters, a good prey-base and basking and nesting beaches hinders the survival of gharials and muggers. The population trend of these species in the Ganga River cannot be inferred from a literature review due to a lack of consolidated information.



Table 3.7
Past and present
population status
of gharials and
muggers along
the Ganga River
and its tributaries

Species	River	Past		Present		Change	Reference
		No.	Year	No.	Year		
Gharial	Ganga	-	-	<200	2006	-	Andrews (2006); Whitaker (2007)
	Chambal	107	1979	1065	1992	895% increase	Hussain (1999)
		226	1997	78	2006	65% decrease	Andrews (2006); Sharma and Basu (2004); Whitaker (2007)
	Chambal (Tasod to Chakar)	311-368	2004 - 2008	720	2015	95% increase	Sharma (2004, 2005, 2006, 2007); Anon (2015)
	Chambal (stretch between Pali and Khirkan)			122	2009	10% increase	Katdare et al. (2011)
	Girwa (Katarniaghat)	30	1997	26 (20 F/6 M)	2006	13% decrease	Andrews (2006); Sharma and Basu (2004); Whitaker (2007)
	Son	10	1997	3	2006	70 % decrease	
	Corbett Tiger Reserve (confluence of Ramganga River and Kalagarh Reservoir)	18	2008	32	2013	77% increase	Chowfin and Leslie (2013)
	Bhagalpur and Sultanganj				Sighted		Bharati et al. (2016)
	Gandak			25	2012		
Mugger	Patna			4	2007		
	Corbett Tiger Reserve (Ramganga)			129	2007		Nawab (2007)
	Chambal	15	1998				Nair (2009)
	Chambal (Pali-Rajghat-Ch. Nagar)	33	1984-1985	356	2014	Increase	Sharma and Singh (2015)
	Pali-Rajghat	23	1984-1985	147	2014	Increase	
	Rajghat-Ch. Nagar	10	1984-1985	209	2014	Increase	



Turtles

3.4.2

India has one of the most diverse turtle fauna, with 24 species of freshwater turtle and four species of tortoise and ranks among the top five countries in terms of importance for turtle conservation in Asia (Stuart & Thorbiarnarson, 2003) and the world (Rhodin, 2006). Freshwater turtles were first described by Gunther (1864) in 'The Reptiles of British India'. The Ganga River provides habitats for 14 testudines (13 turtle and 1 tortoise species) in the middle and lower stretches up to the deltaic region (ZSI, 1991). The turtles in Ganga are represented by nine hardshell species, viz. *Batagur kachuga*, *B. dhongoka*, *B. baska*, *Hardella thurjii*, *Pangshura tecta*, *P. smithii*, *Geoclemys hamiltonii*, *Melanochelys trijuga* and *P. tentoria* (two subspecies, *P. t. tentoria* and *P. t. circumdata*) and four softshell species, viz. *Nilssonia gangetica*, *N. hurum*, *Chitra indica*, and *Lissemys punctata* (two subspecies, *L. p. punctata* and *L. p. andersoni*) (Annexure IV).

Turtles play a critical ecological role by controlling aquatic vegetation, serve as scavengers and help maintain the health of rivers and lakes. Unfortunately, these remarkable reptiles are now facing extinction due to habitat degradation and poaching. Ample literature is available on the turtles of the Indian subcontinent, however, extensive ecological studies are needed to assess the population trend of the turtle species in the Ganga River and its basin.



Red-crowned roofed turtle (*Batagur kachuga*)



3.4.2.1

The turtles belonging to the genus *Batagur* are herbivorous and aquatic and grow to a large size. With the males exhibiting striking seasonal breeding colours, they are also some of the most attractive and rarest turtles in the world. The red-crowned roofed turtle (*Batagur kachuga*) is endemic to South Asia. This rare turtle, previously categorized as Endangered in the IUCN Red list, was updated to Critically Endangered in 2000 owing to its alarming disappearance from its range (IUCN, 2016).

Little is known about the historic distribution of the red-crowned roofed turtle. Anderson (1871) observed the species in the Yamuna River in Agra. Native to south Asia, it had an extensive historic range in northern India, Nepal and Bangladesh (IUCN, 2016; Gray, 2017) (Figure 3.24). It was historically reported from the Ganga, Brahmaputra and Chambal river systems, of northern India (IUCN, 2016).

The last known stronghold of this large river turtle is the Chambal River, in central India. However, small isolated populations still exist in the Ganga and Brahmaputra river basins (IUCN, 2016). The species has been decimated by extensive hunting and intense habitat degradation, including pollution and large-scale water extraction projects for agriculture and drinking purposes. The main anthropogenic threats to the remaining population are accidental drowning of adults in illegal fishing nets, sand-mining, agricultural cultivation on sandbanks and sandbars, water diversion and irregular flows from upstream dams.




Figure 3.24
Distribution of
red-crowned
roofed turtle
(*Batagur
kachuga*)




Northern river terrapin (*Batagur baska*)


The northern river terrapin (*Batagur baska*) is a Critically Endangered turtle species owing to its small and declining wild population and regional extinctions in most of its range (IUCN, 2016).

Historically, the species was distributed throughout south Asia from the Brahmani-Baitarani Delta (Odisha) and the Sundarban region through the mouth of the Irrawaddy River in Myanmar and the Malay Peninsula (southern Thailand, Malaysia) to Sumatra, Cambodia and southern Vietnam (Praschag et al., 2008; Moll et al., 2009) (Figure 3.25). Museum specimens of the species from Salt Lake in Kolkata, and the Indus Delta of southern Pakistan, suggest that this species was distributed in these areas during the mid-19th century (Moll et al., 2009). Das (1987) reported that the species was found in tidal rivers, channels and estuaries and along the coasts of Odisha and the Sundarbans of West Bengal.

Recent studies indicate that a huge population of *B. baska* was extirpated from the river deltas and estuaries of Odisha and West Bengal in India, and the Ayeyarwady Delta in Myanmar, during the 19th and early 20th centuries (Moll et al., 2009; Sinha et al., 2014). No viable population and wild nesting sites of this species are currently known (Hudson, 2016). During 2015–2016, a few remnant individuals were recorded from village ponds in the Sundarbans, where the local fishermen maintain the turtles as source of eggs, as there are no longer any known nesting areas (unpublished data). The species was extirpated from the Hooghly River due to overharvesting of both adults and eggs for human consumption.

Figure 3.25
Distribution of
northern river
terrapin
(*Batagur baska*)





Three-striped roofed turtle (*Batagur dhongoka*)

3.4.2.3

The three-striped roofed turtle is endemic to the rivers of Bangladesh and India (Gunther, 1864). The status of the species was updated from Lower Risk/Near Threatened in 1996 to Endangered in 2000 in the IUCN Red List, and it is listed in Schedule I of the IWPA (1972). Historically the species was distributed in the Gangetic Plain, in Uttar Pradesh, Uttarakhand, Madhya Pradesh, West Bengal and Bihar (Gunther, 1864; Theobald, 1876). The species was recorded from Corbett National Park in Uttarakhand (Moll, 1986; Hanfee, 1999), Dudhwa Tiger Reserve in Uttar Pradesh, and National Chambal Sanctuary in Madhya Pradesh (Hanfee, 1999) (Figure 3.26). The species is currently distributed in the drainages of North India in the Ganga and Chambal rivers (Devaprakash, 2015) and is among the most exploited turtles of southern Asia due to its predictable nesting habits and site fidelity.




Figure 3.26
Distribution of
three-striped
roofed turtle
(*Batagur
dhongoka*)



3.4.2.4




Spotted pond turtle (*Geoclemys hamiltonii*)

The spotted pond turtle (*Geoclemys hamiltonii*), also known as the black spotted turtle or the black pond turtle is endemic to south Asia. It is classified as Vulnerable (VU) according to the IUCN Red List, and it is listed in Schedule I of the IWPA (1972). The species is carnivorous.

The spotted pond turtle is a relatively rare species, found only in the Indus and Ganga river drainages in Pakistan, northern India, Bangladesh and Nepal. Historically, the turtle was abundant in the Lower Ganga and was reported from the Gandak Dam, of Bihar, and its vicinity (Gunther, 1864; Moll & Vijaya, 1986; Theobald, 1876). In the 19th century, this turtle was widely distributed in the deep water systems of the northern and northeastern parts of the Indian subcontinent (Smith, 1931; Vijaya, 1983) (Figure 3.27). Bhaduria et al. (1995) reported the occurrence of the species in the Gomti and Girwa rivers, of Uttar Pradesh. More recent studies indicate the presence of the species in PAs along the Ganga River and its tributaries and in West Bengal (Daniel, 2002; Sinha et al., 2014). The most significant threat to the black pond turtle is uncontrolled trade of freshwater turtles in Asia.

Figure 3.27
Distribution of
spotted pond
turtle (*Geoclemys*
hamiltonii)





Indian tent turtle (*Pangshura tentoria*)



3.4.2.5

The Indian tent turtle is similar to the Indian roofed turtle. The species is listed under the Least Concern category in the IUCN Red List and Schedule I of the IWPA (1972). It is mainly omnivorous and consumes a wide variety of aquatic plants and animals in the wild.

The Indian tent turtle (*Pangshura tentoria*) is found in central and northern India and Bangladesh (Theobald, 1876). In the recent past, the species has been reported from Kolkata, Bihar (Moll, 1987) and Katarniaghat Wildlife Sanctuary and Dudhwa Tiger Reserve in Uttar Pradesh (Hanfee, 1999). The Indian tent turtle is restricted to the upper and central rivers of India i.e. the rivers flowing into the Bay of Bengal (Moll, 1987) (Figure 3.28). Rao (1993) reported the species from the Chambal River. Baruah et al. (2016) mentioned the presence of the species in the Yamuna River and at the Kalindri–Ganga confluence, in Uttar Pradesh, and at Farrukhabad (Uttar Pradesh), Deogarh (Madhya Pradesh), Katarniaghat, Gorakhpur (Madhya Pradesh) and West Bengal.

During the rapid biodiversity assessment, evidence of the presence of *Pangshura* was found all along the mainstem of the river, especially in the middle stretches, such as at Bijnor.




Figure 3.28
Distribution of
Indian tent turtle
(*Pangshura*
tentoria)



3.4.2.6



Brown roofed turtle (*Pangshura smithii*)

The species is listed as Near Threatened (NT) in the IUCN Red List and in Schedule IV of the IWPA (1972). It is a rapid swimmer and is omnivorous in nature (Das, 1991; Sial et al., 2016).

The species inhabits medium to large rivers with vegetation. It is distributed in the Upper and Middle Ganga (Moll, 1987; Hanfee, 1999) and in West Bengal in the Lower Ganga (Daniel, 1970). The main anthropogenic factors affecting the species include habitat degradation by the construction of dams and barrages, cultivation along river banks and pollution of aquatic habitats (Bista & Shah, 2010).



3.4.2.7



Indian roofed turtle (*Pangshura tecta*)

This unique crawling reptile is listed as Least Concern (LC) in the IUCN Red List and in Schedule I of the IWPA (1972) and has an omnivorous diet.

The Indian roofed turtle has been recorded from lentic habitats of the Ganga River and is commonly seen basking on logs or river banks (Moll, 1987). Historically, *P. tecta* was distributed in northern and peninsular India (Smith, 1931; Edds, 1998). The single greatest threat to *P. tecta* is commercial collection of the species for the pet trade. The species has been observed in food markets in south China (Lau & Shi, 2000).



Crowned river turtle
(*Hardella thurjii*)



3.4.2.8

This large freshwater turtle is classified as Vulnerable (VU) in the IUCN Red List and is listed in Schedule IV of the IWPA (1972). The crowned river turtle has an omnivorous diet.

Historically the species was distributed in the Middle Ganga and the Brahmaputra River (Chaudhuri, 1912). In recent studies, this species was found in the middle and lower reaches of the northern river systems, comprising the Indus, Ganga and Brahmaputra and their tributaries (Das & Bhupathy, 2009b). This species occurs in slow-moving rivers (Daniel, 1970). This hardshell species is commonly sold in eastern Indian markets because its body is large. Some studies indicate that hatchlings of the species are exported as pet (Rashid & Khan, 2000). The other threats to the species include pollution and conversion of wetlands.



Indian black turtle
(*Melanochelys trijuga*)



3.4.2.9

This medium-sized turtle is listed as Near Threatened (NT) in the IUCN Red List and in Schedule IV of the IWPA (1972). The species is omnivorous.

M. trijuga is a relatively common species. It is the most widespread of Indian terrapins. Historically, the species was distributed in the hill ranges of the Western Ghats and south Gujarat (Daniel & Shull, 1964). It has also been recorded from Uttarakhand, in the northern part of India, and Meghalaya, in northeastern India (Das, 1991). The species is exploited commercially for food in northeastern India (Das, 1991; Pawar & Choudhury, 2000), West Bengal and Kerala (Das & Bhupathy, 2009a).



3.4.2.10




Indian softshell turtle (*Nilssonia gangetica*)

The Indian softshell turtle inhabits deep rivers, streams, large canals, lakes and ponds with a bed of mud or sand and prefers areas with turbid waters (Ernst et al., 1997). This peculiar-looking freshwater reptile is listed as Vulnerable in the IUCN Red List and in Schedule I of the IWPA (1972). It feeds mostly on fish, amphibians, carrion and other animal matter, but it also feeds on aquatic plants.

The species inhabits multiple river systems, including the Ganga, Indus and the Mahanadi systems. Historically, the Indian softshell turtle was distributed in the Ganga River and its tributaries. It was also recorded from Pakistan, Bangladesh and southern Nepal (Figure 3.29) (Annandale, 1912; Gunther, 1864). Annandale (1912) reported the species from different places in the Gangetic Delta, especially from Khulna and markets of West Bengal. Das et al. (2010) reported nests of *N. gangetica* from the Chambal River.

The main threats to the Indian soft-shelled turtle is the trade in this species in east Asia for its meat and fishing as this river-dwelling turtle becomes trapped in fishermen's nets. Pollution is also a significant problem in the rivers inhabited by these turtles. Extensive studies are needed to assess the population trend of this species in the Ganga River and its basin. During the rapid biodiversity assessment, the species was abundant all along the mainstem of the Ganga River.

Figure 3.29
Distribution of
Indian softshell
turtle (*Nilssonia*
gangetica)





Indian peacock softshell turtle (*Nilssonia hurum*)



3.4.2.11

The Indian peacock softshell turtle is similar to the Indian softshell turtle, but it is smaller in size. The species is listed as Vulnerable in the IUCN Red List and in Schedule I of the IWPA (1972). The Indian peacock softshell turtle (*Nilssonia hurum*) is a relatively abundant, large riverine species and widespread in the northern and central parts of India, Pakistan, Bangladesh and Nepal.

Annandale (1912) reported the species from the lower reaches of the Ganga River. Later, Smith (1931) and Moll and Vijaya (1986) reported the species from the Ganga River as well as from its tributaries. Das (1988) reported the species from Bhopal and Madhya Pradesh. The species is widespread in the northern and central parts of the Indian subcontinent, but the southern and eastern limits of its distribution remain unclear (Figure 3.30). Das et al. (2010) reported *N. hurum* nests from the Chambal River (these nests were less than 2% of *N. gangetica* nests) and reported the species from the Saryu, Kane and Rapti rivers and a few oxbow lakes in the area. Sinha et al. (2014) reported the species from Bihar, Madhya Pradesh, Rajasthan, Uttar Pradesh and West Bengal. *N. hurum* appears to be the most common large softshell turtle in the lower parts of the Ganga River system; however, information on the population sizes is still lacking (Das et al., 2010). The species is extensively hunted in all of northern and eastern India for its meat and calipee (Das et al., 2010).




Figure 3.30
Distribution of
Indian peacock
softshell turtle
(*Nilssonia hurum*)




3.4.2.12



Indian narrow-headed softshell turtle (*Chitra indica*)

The Indian narrow-headed softshell turtle is an extremely large aquatic species. The species is listed as Endangered in the IUCN Red List and in Schedule II of the IWPA (1972). The species is widespread in South Asia, and its distribution in India includes the Indus, Ganga, Godavari, Coleroon and Mahanadi rivers (Das, 1991; Moll, 1986) (Figure 3.31). The distribution of the species is extensive but apparently localized and patchy. Gunther (1864) and Theobald (1876) reported the species from the Ganga River and its tributaries. Annandale (1912) reported that the species was common in the Gangetic Delta and that it was commercially exploited in West Bengal. Inglis et al. (1919) recorded *C. indica* in Jalpaiguri District, of West Bengal. Smith (1931) reported it from Fatehgarh on the Ganga River. Iverson (1992) reported the species from Barrackpore, and Sharma (1998) reported it from Uttar Pradesh, Bihar and West Bengal. Hanfee (1999) and later Das and Singh (2009) found the species in the Ghagra and Chambal rivers. The species is now hunted intensively for its calipee and is threatened by the modification of the riverine habitat (Das & Singh, 2009).

Figure 3.31
Distribution of
Indian narrow-
headed softshell
turtle (*Chitra*
indica)





Indian flapshell turtle (*Lissemys punctata*)



3.4.2.13

This relatively small softshell turtle is listed as Least Concern (LC) in the IUCN Red List and in Schedule I of the IWPA (1972). The species is an opportunistic omnivore. Das (1995) noted a seasonal shift in the diet of this turtle in northern India. It is normally carnivorous.


Lissemys punctata is restricted to the Indian sub-region and has been found in the states of Assam, Bihar, Haryana, Jammu & Kashmir, Madhya Pradesh, Meghalaya, Rajasthan, Sikkim, Uttar Pradesh and West Bengal. It is also found in the southern part of peninsular India in the states of Kerala and Tamil Nadu. *L. punctata* and its eggs are rather heavily exploited for food, and it is one of the species observed most frequently in markets (Whitaker, 1997).

During the rapid biodiversity assessment, a total of 2788 turtles were sighted. These were most frequently encountered between Makdumpur and Kanpur Barrage (Figure 3.32). Of these, 2693 were hardshell turtles, with *Pangshura* spp. being the most abundant, followed by *Batagur dhongoka* and a few individuals of *Geoclemys hamiltonii*, and *Hardella thurjii*. Between the two species of softshell turtles sighted viz., *Nilssonia gangetica* and *Lissemys punctata*, sightings of *N. gangetica* was more frequent.

Highest numbers of individuals were encountered between Farukhabad and Kanpur, where few individuals of *G. hamiltonii*, *H. thurjii* and *B. dhongoka* were recorded. This was followed by the stretch from Makdumpur to Narora, where *Pangshura* spp. was most abundant, and a few individuals of *B. dhongoka* and *N. gangetica* were also encountered. The least number of individuals were recorded between Kachla and Farukhabad, where *Pangshura* spp. was the most abundant, followed by *L. punctata*, *N. gangetica*, *G. hamiltonii* and *B. dhongoka*. Scattered populations of *Pangshura* spp. and *N. gangetica* were sighted near Manikpur, Allahabad and Baharampur.

The river stretch with the highest number of turtle sightings is dynamic with shallow water and numerous sandy islands, which are preferred by turtles for basking. The riverbanks between Bijnor and Narora were less disturbed as human habitations were far from the banks. Hardshell turtle species such as *Pangshura* spp. and *Batagur* sp. were observed basking on the loamy shore, which is characterized by soil bolus with little vegetation. Individuals of these species were also observed basking on submerged logs and bushes. Individuals of softshell turtle such as *N. gangetica* were mostly observed basking along the sandy islands.

Figure 3.32
Distribution of
turtles in the
Ganga River on
the basis of rapid
biodiversity
assessment



The population trends of these turtle species in the Ganga River cannot be inferred from a literature review due to a lack of information. Habitat destruction and poaching are the major threats to the survival of these species. Species such as *Batagur baska* have been extirpated from the lower stretches due to poaching and overexploitation. Sightings of recruitment class (hatchlings and juveniles) between Bijnor and Narora indicates a recovering population.



Amphibians

3.5

Amphibians play a pivotal role in both aquatic and terrestrial ecosystems as secondary consumers in several food chains. From the ecological perspective, amphibians are regarded as good ecological indicators. Their responses have been used to indicate habitat fragmentation, ecosystem stress, impacts of pesticides, etc. Adult amphibians are one of the best biological pest controllers.

Amphibian populations are currently undergoing a rapid decline due to habitat loss, overutilization and fungal diseases. Amphibians are semi-aquatic vertebrates, and their life cycle is primarily dependent on freshwater resources. The eggs are laid in water or other moist habitats and the larvae grow in water bodies, where they metamorphose into adults. The characteristic feature of this group is the presence of semi-permeable skin, used for cutaneous gas exchange. This feature and the dependence of these animals on freshwater ecosystems make them acutely sensitive and susceptible to slight changes in temperature, humidity and air or water quality, thereby making them highly vulnerable to climate change.

The Ganga River nurtures about 90 species of amphibian in its basin, from as high as 3000 m asl through the plains to the delta. The endemic amphibian *Amolops chakrataensis* (Chakrata stream frog) is known only from the hill streams from Chakrata, in Uttarakhand. Other frogs such as the Indian bullfrog (*Haplobatrachus tigrinus*), Himalaya paa frog (*Nanorana vicina*) and Nepal paa frog (*Nanorana micina*) are reported from various stretches of the river. According to Bilgrami (1991), the genus *Rana* is distributed widely across the Ganga River. Bilgrami (1991) recorded *R. limnocharis*, *R. tigrina* and *R. cynophlyctes* from the middle and lower stretches of the Ganga River. Amphibians are the least studied group and hence information pertaining to them is either unavailable or scanty. Annexure VI provides a comprehensive list of amphibian species of the Ganga River. On the basis of the available literature, descriptions of a few species are provided in the following pages. The literature pertaining to amphibians is limited. Therefore, extensive studies are needed to assess the population trends of these species in the Ganga River and its basin. The presence and distribution of these species along the Ganga River can only be defined after an extensive survey, which is currently underway.



3.5.1


Himalaya paa frog (*Nanorana vicina*)

The species is a medium-sized frog (58 mm), with olive or brown coloured body. It is listed as Least Concern in the IUCN Red List.

The Himalaya paa frog is associated with high-altitude streams, springs, fountains and other running water within open forests and grassland habitats. The threats faced by this species are currently not known, but habitat destruction is presumed to be one of them.

Figure 3.33
Distribution of
Himalaya paa
frog (*Nanorana*
vicina)

It is mainly reported from the upper stretches of the Ganga River, from elevations of 2000–3000 m asl (Figure 3.33).






Annandale's paa frog (*Nanorana annandalii*)



3.5.2

The species is listed as Near Threatened in the IUCN Red List. It is a medium-sized species (55 mm). It is olive coloured, with marbling and a white-coloured belly. It is distributed in the upper stretches of the Ganga River between 1500 and 2000 m asl (Figure 3.34). This frog is found in rocky streams and brooks in montane forests and in pools in forest clearings. Stream modification due to boulder collection and dams are major threats.

Figure 3.34
Distribution of
Annandale's paa
frog (*Nanorana*
annandalii)






3.5.3


Nepal paa frog (*Nanorana minica*)

The species has a relatively small-sized body (28—41 mm), with a brownish colour, with black spots and small warts at the back of the body. This frog is found only in montane sub-tropical forests and streams.


The distribution of the species is highly fragmented and is restricted to western Nepal, and northern India. It is listed as Vulnerable in the IUCN Red List. Habitat loss through dam construction and pollution are the major threats to the species. In the Ganga River, it is reported from the upper stretch, from 1000 to 2400 m asl (Figure 3.35).

Figure 3.35
Distribution of
Nepal paa frog
(*Nanorana minica*)





Cascade frog (*Amolops formosus*)




3.5.4

The cascade frogs is a large-sized frog (75 mm), with a green coloured body with dark brown or black spots. It is listed as Least Concern in the IUCN Red List.

Cascade frogs are obligate stream dwelling species, associated with streams and riparian vegetation and breed along fast-flowing waters. Their tadpoles have ventral suckers, using which they attach themselves to rocks in fast-flowing streams. Changes in river hydrology caused by dams and deforestation are major threats.

The species occurs in the upper stretch of the Ganga River between 1000 and 2500 mm asl (Figure 3.36).

Figure 3.36
Distribution of
cascade frog
(*Amolops*
formosus)






3.5.5



Marbled toad (*Duttaphrynus stomaticus*)

This species is moderately large bodied (76 mm). It is light brown in colour, with numerous warts and marblings. A large parotid gland is present behind the eye. The marbled toad is found in open plains, grasslands, scrubland, forests, agricultural land and human habitations. It breeds in both permanent and seasonal pools, seasonal streams and slow-flowing streams. It is listed as Least Concern in the IUCN Red List. Threats to the species include habitat loss and water pollution due to pesticide and herbicide use in agricultural lands. It is distributed in the upper, middle and lower stretches of the Ganga River (Figure 3.37).

Figure 3.37
Distribution of
marbled toad
(*Duttaphrynus*
stomaticus)






Tytler's pond frog (*Hylarana tytleri*)



3.5.6

This is a lowland species found up to 300 m asl in the middle stretch of the Ganga River, in pools, lakes and marshes (Figure 3.38). The species generally breeds in stagnant water bodies. It is listed as Least Concern in the IUCN Red List. The species is threatened by water pollution due to the use of agrochemicals.

Figure 3.38
Distribution of
Tytler's pond frog
(*Hylarana tytleri*)





3.5.7




Indian bullfrog (*Hoplobatrachus tigerinus*)

The Indian bullfrog is a large-sized frog (134 mm), greenish, olive or brown in colour, with a mid-dorsal yellow line. It is a common frog with a wide distribution along the Ganga River (Figure 3.39) and is listed as Least Concern in the IUCN Red List and in Schedule IV of the IWPA (1972).

This frog is found in freshwater wetlands, especially in paddy fields and ponds. It breeds during the monsoon and feeds on invertebrates, small mammals and birds. The species is reported from the middle and lower stretches of the Ganga River. The major threats to the species include loss of wetland habitats, water pollution due to agrochemical use, road mortality and consumption as food.

Figure 3.39
Distribution of
Indian bullfrog
(*Hoplobatrachus*
tigerinus)





Jerdon's bullfrog (*Hoplobatrachus crassus*)




3.5.8

This is a large-sized frog (121 mm). Its dorsal body is brown with dark spots. Irregular glandular folds are present on the dorsal side. Jerdon's bullfrog is found in seasonally flooded grasslands, open plains and cultivated areas and around human settlements. It is listed as Least Concern in the IUCN Red List and in Schedule IV of the IWPA (1972).

The species breeds in various types of water body, but it prefers large rivers. It has been reported from the middle and lower stretches of the Ganga River (Figure 3.40). Habitat loss and collection for food are major threats.

Figure 3.40
Distribution of
Jerdon's bullfrog
(*Hoplobatrachus*
crassus)





3.5.9




Dudhwa tree frog (*Chiromantis dudhwaensis*)

The Dudhwa tree frog is a small-sized frog, brownish yellow in colour. The species is currently known only from the type locality, Dudhwa National Park, in Uttar Pradesh, India, where it was observed and collected below 100 m asl (Figure 3.41). However, it is believed that the species occurs more widely. The arboreal species is mostly associated with scrub forests, grasslands and rural areas. The species is Data Deficient in the IUCN Red List. Although there are currently no known threats, this frog may be vulnerable to habitat loss and climatic shifts.

Figure 3.41
Distribution of
Dudhwa tree frog
(*Chiromantis*
dudhwaensis)

The population trends as well as distribution of these amphibian species cannot be inferred from the literature as information pertaining to it is lacking. Detailed studies are required to identify these trends and their vulnerability to environmental and anthropogenic changes.





Fish


3.6

Fishes regulate trophic structure and influence temporal availability of nutrients, as they mineralize nitrogen and phosphorus, making these nutrients available for primary production. They also transport nutrients across different ecosystems. The unique hydrology and geo-climatic conditions of the upper, middle and lower stretches of the Ganga River support a distinctive fish fauna. Though most of the fishes presently found in the Ganga River belong to the class Actinopterygii, there are historical records of fishes of the class Elasmobranchii. Two species of stingray were found in the Ganga River up to Kanpur by Hamilton during his “Statistical and Economic Survey”, conducted between 1807 and 1814 (Chaudhuri, 1911). Later, Day (1878) and Chaudhuri (1911) also recorded these stingray species from the Ganga River specifically at Bhagalpur. Another elasmobranch, the Gangetic shark (*Glyphis gangeticus*), was reported by Muller and Henle (1839) and later by Bal and Rao (1984) in the Ganga River beyond the tidal range (ZSI, 1992). The present status of these species of the class Elasmobranchii is mostly unknown.

The earliest documented records of fish species from the Ganga River go back to 1822, when Hamilton recorded 272 freshwater fish species (Hamilton, 1822). Later, Bilgrami (1991) reported 97 species along three stretches of the Ganga River, viz. Kalankar to Phaphamau, Buxar to Barh and Munger to Farakka. Talwar (1991) carried out a comprehensive study of the entire Ganga River and recorded a total of 375 species, including both freshwater and brackish water species. Later, Payne et al. (2004) reported 161 species from the river. Sarkar et al. (2012) reported 143 species belonging to 32 families. A recent report by the Central Inland Fisheries Research Institute (ICAR-CIFRI, 2017) lists a total of 158 species from Tehri (Uttarakhand) to Fraserganj (West Bengal) in the mainstem of the Ganga River (Figure 3.42). Large discrepancies are seen in the number of fish species reported from the Ganga River, in various survey reports, most of which can be accounted for by the variations in sampling stretches and seasons. The report produced by ICAR-CIFRI (2017) is the most comprehensive one relating to the fish species diversity of the entire mainstem Ganga River. Literature review revealed that 236 fish species representing 60 families are found in the Ganga River (Annexure VII).

The most common fishes reported by Bilgrami (1991) in the upper stretch of the Ganga River are represented by *Schizothorax* spp., *Tor* spp., *Wallago attu* and *Notopterus* sp. In the middle stretch, the most common species included the Indian major carps, *Bagarius* sp., *Mystus* sp., *Tenualosa ilisha* and *Wallago attu*. Bilgrami (1991) expressed concern that the dominance of carnivorous species in the lower stretch of the river from Berhampur to Katwa could adversely affect the natural food chain. Sarkar et al. (2012) reported 10 exotic fish species, viz. *Pterygoplichthys anisitsi*, *Oncorhynchus mykiss*, *Oreochromis mossambicus*, *O. niloticus niloticus*, *Hypophthalmichthys molitrix*, *H. nobilis*, *Ctenopharyngodon idella*, *Cyprinus carpio*, *C. carpio* (var. *specularis*), and *Clarias gariepinus*. ICAR-CIFRI (2017) also reported these exotic species, with an additional exotic *Salmo trutta fario* from the Ganga River.

Figure 3.42
Fish species
diversity in the
Ganga River






Golden mahaseer (*Tor putitora*)




3.6.1

Found in streams and rivers of montane and sub-montane regions, *Tor putitora* is a major tor species of the mid-hill stretches of the Himalayan region. The species is listed as Endangered in the IUCN Red List and has a declining population trend. It inhabits rapid streams with rocky bottoms, riverine wetlands and lakes. Threats to the species include overfishing, the use of modern fishing techniques and alterations of habitats. The species was once abundant in Uttarakhand (Stevens, 1905) and was also reported from the Song–Ganga confluence (Dhu, 1923) and from the Sharda River and its tributaries (Macdonald, 1935). The species has also been reported from south Asia and southeast Asia (Jha & Rayamajhi, 2010). In India, it is restricted to the upper stretches of the Ganga River (Figure 3.43). During the rapid biodiversity assessment, *T. putitora* was recorded from the upper stretch of the Ganga River.

Figure 3.43
Distribution of
golden mahaseer
(*Tor putitora*),
adapted from
IUCN





3.6.2




Bronze featherback (*Notopterus notopterus*)

The species is found in clear streams and brackish waters. The adults are found in standing and sluggish waters of lakes, floodplains, canals and ponds (Froese & Pauly, 2017). The bronze featherback was described by Pallas (1769) as *Gymnotus notopterus* and as occurring in Indian rivers and estuaries. The species is distributed widely in southeast Asia, Bangladesh, Cambodia, India, Laos, Burma, Nepal and Pakistan. Its status is not clear because its taxonomy is in an undetermined state at present. It has been suggested that there is a species complex (Ng, 2010). Population trend of the bronze featherback in the wild is currently unknown and it is listed as Least Concern in the IUCN Red List.

In India, this fish is widely distributed in the Indus and Ganga–Brahmaputra basins as well as in the Mahanadi, Krishna, Cauvery and other river basins in peninsular India (Talwar & Jhingran, 1991) (Figure 3.44). The species was reported from the Mahanadi River near the Hirakud Dam (Jayaram & Majumdar, 1976). Specimens of the species were collected from the Gomti River near Daliganj and from Khadra, at Lucknow (Srivastava et al., 2012). The fish has been categorized as an important commercial fish by FAO (Casavas et al., 1996) for its use as food as well as the ornamental trade in the species.

Figure 3.44
Distribution of
Bronze
featherback
(*Notopterus
notopterus*)





Freshwater shark (*Wallago attu*)



3.6.3

Wallago attu is widely distributed in the Ganga River and its tributaries. Declared as Near Threatened in the IUCN Red List, there is a declining trend of the species in its range. *W. attu* has been reported from India, Pakistan, Bangladesh, Srilanka, Thailand, Burma, Java, Sumatra, Borneo and Afghanistan (Giri et al., 2002; Talwar & Jhingran, 1991) (Figure 3.45). In earlier records, CIFRI (1964) reported 49.5 t landing of *W. Attu* in eight landing sites near the Ganga River in 1963. Ray (1998) reported that the landings at Patna decreased from 3.89 t to 1.22 t during 1988–1989. A similar decreasing trend in the landings was noted at Bhagalpur, from 12.04 t in 1988 to 0.06 t in 2006 (Montana et al., 2011; Ray, 1998).

Apart from fish landing data, primary catch data are also required to assess the population trends of fishes in the river. In such a study, Dwivedi et al. (2016) reported that the catch of this species was 4.78% of the total catch, from August 2012 to July 2013 at landing sites at Kanpur, Allahabad and Varanasi. However, more such quantitative studies are needed to assess their population in the Ganga River and its basin.




Figure 3.45
Distribution of
Freshwater shark
(*Wallago attu*)




3.6.4



Minor carp (*Labeo bata*)

L. bata is a benthopelagic and potamodromous species, inhabiting ponds, rivers and rivulets (Bhuiyan, 1964). It is an herbivorous column feeder. It is distributed throughout the Indian subcontinent, including Bangladesh, India, Nepal, Myanmar and Pakistan (Talwar & Jhingran, 1991) (Figure 3.46). With the population trend currently unknown, the species is listed as Least Concern in the IUCN Red List. It is commercially exploited in the Ganga River and its tributaries (Dwivedi, 2013; Dwivedi et al., 2004, 2007; Tripathi et al., 2013). Size-selective harvesting of the species is a very common practice in riverine habitats (Dwivedi & Nautiyal, 2010; Fenberg & Roy, 2008).

Figure 3.46
Distribution of
Minor carp
(*Labeo bata*)



The abundance of the major carps including *L. bata* was greater in the middle stretches of the Ganga River. These species accounted for 53% of the catch at Agra, 45% at Kanpur and 38% at Allahabad, but they were less abundant in the lower stretches (19–26%) at Patna (Jhingran, 1991; Payne et al., 2004). In landings near Patna, the major carps represented 26.5% of the total catch in 1958–1966, and by 1993–1994, the proportion reduced to 4.0% of the total catch (Montana et al., 2011). In a survey conducted during 2001–2007, the percentage of major carps (including *L. bata*) in the total annual catch varied between 9% and 15%, reflecting a small percentage compared with the historical catches from the region (Montana et al., 2011).



Spotted barb (*Puntius sophore*)



3.6.5

P. sophore is a small-sized cyprinid, highly adapted to confined environments (Collins et al., 2012). It is widely distributed throughout India, Pakistan, China and some parts of the southeast Asian countries (Petr, 1999; Collins et al., 2012) (Figure 3.47) and is listed as Least Concern in the IUCN Red List. Historically, the species was native to Bengal and common throughout the Gangetic provinces (Hamilton, 1822). It was reported from the Ramganga River (Atkore, 2005; Atkore et al., 2011), the Paisuni River, a tributary of the Yamuna River (Dwivedi, 2009), the Betwa River (Lakra et al., 2010b) and the upper, middle and lower stretches of the Ganga River (Das et al., 2013), including Kanpur, Allahabad and Varanasi (Dwivedi et al., 2016). In India, populations of *P. sophore* are reported to be declining owing to overexploitation (Latif et al., 2017).

From the literature review, it can be inferred that the populations of species such as the golden mahaseer, bronze featherback and spotted barb are declining due to habitat degradation and overexploitation. An overall decline in fish landings is also evident throughout the stretch.




Figure 3.47
Distribution of
Spotted barb
(*Puntius sophore*)

During the rapid biodiversity assessment, 114 species belonging to 60 families were recorded, including the abovementioned species in their respective stretches along the Ganga River (Annexure VII). Golden Mahaseer, mahaseer and snow trouts were only encountered in the upper stretch of the Ganga River. *W. attu* was frequently observed in the middle and lower stretches of the river. *Tenulosa ilisha* was encountered in the lower stretch downstream of Farakka barrage in the mainstem Ganga River and in the Feeder Canal.


River Stretches with High Biodiversity Value

3.7

Identification of areas with high biodiversity values that are relatively intact and prioritizing them for ecological restoration is based on the 'refuge approach' to restoration. This approach provides resilience and sources for future restored sites (Beechie et al., 2008) and also provides a measure against the likelihood of local extirpation of species. Thus, for ecological restoration of the Ganga River, stretches with high biodiversity values were identified using information obtained from the rapid biodiversity assessment. Habitat-defining variables such as water depth were also determined along the entire length of the Ganga River.

Assessment of hydrology of the river revealed that channel depth was a major limiting factor for species distribution. The thematic diagram shows the depth profile of the Ganga River from Bijnor to Nurpur during summer months (Figure 3.48).





According to the depth profile, depth categories were identified that answered to the preference of umbrella species, the Gangetic river dolphin and the gharial (Figure 3.48). These species showed a preference for depths above 4 m, which were available in scattered pools in approximately 38.7% of the river stretch assessed (Figures 3.48 and 3.49). Shallow areas (depth < 4 m) prevail throughout the stretch. These shallow areas act as a barrier to the lateral movement of these large-bodied aquatic animals.




Figure 3.49
Depth class distribution of the Ganga River during summer and optimal accessible stretch for umbrella species

The distributions of priority species such as the Gangetic river dolphin, otters, island-nesting birds such as the Indian skimmer, the gharial and the mugger were overlaid in the GIS domain (Figure 3.37). These were integrated with habitat characteristics and the assemblage of benthic macro invertebrates, periphyton and plankton along the Ganga River. Thus, six stretches covering about 50% of the river's length were identified as high biodiversity stretches (Figure 3.50). Table 3.8, summarizes these stretches identified along the Ganga River, in terms of their biogeographical location, river characteristics, species observed and the specific threats.




Figure 3.50
High biodiversity
stretches and
distribution of
priority species
along the Ganga
River

Stretch I: Devprayag to Rishikesh, Uttarakhand

This is a 61 km stretch between Devprayag and Rishikesh in Uttarakhand (Figure 3.50). This stretch falls under the West Himalaya (2B) biogeographic province and is characterized by rapids with rocky bottoms, deep gorges and gentle slopes. The local communities in this region are dependent on the Ganga River for domestic water usage, their livelihoods from religious and adventure tourism and fishing.

Biodiversity value and ecological significance

- This stretch nurtures about 56 species of fish including the Endangered golden mahaseer. Semi-aquatic mammals such as the smooth-coated and Eurasian otters have been reported from this stretch. There are also reports of Asian small-clawed otter from the narrow streams joining Ganga River. 41 waterbird species were recorded at Rishikesh barrage during migratory waterbird survey. About 93 phytoplankton, 76 periphyton and 19 zoobenthic species that form important components of the food web have been recorded from this zone.
- This stretch along with the tributary Nayar River provides breeding grounds for the Golden mahaseer.
- This stretch forms the headwaters for the lower stretches of the Ganga River.
- Riffles and pools provide habitat heterogeneity for the benthic flora and periphyton and forms headwaters for the lower stretches of the Ganga River.

Stretch II: Makdumpur to Narora, Uttar Pradesh

This is a 147 km stretch from Makdumpur, in Jyotibha Phule Nagar District, to Narora, in Bulandshahar District, Uttar Pradesh (Figure 3.50). This stretch of the river falling under the Upper Gangetic Plain (7A), is mostly meandering with extensive alluvium, sandbars and mid-river islands. The flow regime of this

stretch is highly altered by the Bijnor Barrage upstream and the Narora Barrage, at the end of the stretch. During summer months, the river depth ranged from 1.6 m to 5.2 m, with a mean value of 2.3 m. There are major human settlements in Garhmukteshwar, Anupshahar and Narora. The urban population and the local communities in this region are dependent on the Ganga River for daily ablutions, drinking water, agriculture and irrigation.

Biodiversity value and ecological significance

- a. During the rapid assessment a total of 27 gharials were encountered in this stretch. Besides, around 41 sightings of Gangetic river dolphins were also made in this stretch. More than 800 turtles, including 176 individuals of *Batagur* sp., 50 *Nilssonia gangetica* and 619 individuals of *Pangshura* spp., were encountered within this stretch. Around 80 fish and 62 waterbird species were recorded from this stretch. About 35 phytoplankton and 31 zooplankton species have been reported from this stretch.
- b. About 110 km of this stretch is protected as Hastinapur Wildlife Sanctuary and stretch between Brijghat to Narora, is recognized as wetland of international importance.
- c. Presence of sandbars, mid-river islands, backwaters and floodplain wetlands within this stretch provide habitat mosaics for different aquatic biota.

Stretch III: Bhitaura to Ghazipur, Uttar Pradesh

This is a 454 km stretch from Bhitaura, Fatehpur District, to Ghazipur, Ghazipur District, Uttar Pradesh (Figure 3.50). This stretch under the Upper Gangetic Plain (7A) is highly braided and anastomosed and passes through an extensive alluvial plain. The flow regime of this stretch is altered by the Kanpur Barrage upstream, and hence the river depth reaches a value as low as 1.1 m. However, the Yamuna River, a major tributary, joins it near Allahabad, and the flow regime improves significantly, resulting in an increased river depth of up to 20.4 m. This stretch has an average depth of 3.7 m in summer months (range 1.1–20.4 m). There are dense human settlements at Fatehpur, Mirzapur, Mughalsarai, Allahabad and Varanasi. The urban population and rural communities in this region are dependent on the Ganga River for daily ablutions, municipal and industrial water extraction, religious tourism, fishing and irrigation.

Biodiversity value and ecological significance

- a. A total sightings of 269 Gangetic river dolphins were made in this stretch. Around 193 Indian skimmers were also seen. Besides, sightings of *Batagur* sp. and *Nilssonia gangetica* were also made in this stretch. A total of 50 fish species were encountered during fish sampling. The stretch is also known to harbour about 357 phytoplankton, 42 zooplankton, 19 periphyton and 45 zoobenthic species.
- b. Seven km length of this stretch from Rajghat to Ramnagar, in Varansai, is protected as Kachhua Sanctuary.
- c. Sandbars, meander belts and ox-bow lakes within this stretch provide a mosaic of habitats for different aquatic biota.

Table 3.8. Description of the six high biodiversity stretches along the Ganga River

Stretch	Location	Length	Biogeographic Provinces	River characteristics	River depth (m) (average)
I	Devprayag to Rishikesh	61 km	West Himalaya (2B)	Rapids with rocky bottoms, deep gorges and gentle slopes	NA
II	Makdumpur to Narora	147 km	Upper Gangetic Plain (7A)	Meandering channels with extensive alluvium, sandbars and mid-river	1.6 to 5.2 (2.3 m)
III	Bhithaura to Ghazipur	454 km	Upper Gangetic Plain (7A)	Highly braided and anastomosed, with sandbars, meander belts and ox-bow lakes	1.1 to 20.4 (3.7)
IV	Chhapra to Kahalgaon	296 km	Lower Gangetic Plain (7B)	Very wide, highly braided with large, stable islands, meanders and ox-bow lakes	1.6 to 33.2 (4.7 m)
V	Sahibganj to Rajmahal	34 km	Lower Gangetic Plain (7B)	Very wide, highly braided and anastomosing with multichannel formation on an alluvial plain	1.5 to 17 (5.8 m)
VI	Baharampur to Barrackpore	246 km	Lower Gangetic Plain (7B)	Highly meandering, with convex sandbars and a few mid-river islands	2.9 to 31 (8.1 m)

Presence of fauna of conservation significance					Threats
Gangetic river dolphin	Otter species	Avian species	Reptilian species	Fish species	
Nil	3 species	41 waterbird species	-	56 species, including the Golden Mahaseer	Structural changes in river morphology and loss of connectivity due to hydroelectric projects Collection of sand and boulders
41	-	62 waterbirds	2 Crocodilians; 3 Turtle species	80 species	Cultivation on river islands and sand bars, leading to altered river bank Excessive water abstraction leading to reduced flow and water depth Unsustainable resource extraction, such as excessive fishing and sand mining Poaching of turtles
269	-	140, including 193 individuals of Indian skimmer and 6 nesting colonies	2 turtle species	50	Cultivation on river islands and sand bars, leading to altered river bank Excessive water abstraction leading to reduced flow and water depth Unsustainable resource extraction such as excessive fishing and sand mining Industrial and domestic discharge into the river Poaching of turtles
141	1 species	-	-	61	Cultivation on river islands and sand bars, leading to altered river bank Excessive water abstraction leading to reduced flow and water depth Unsustainable resource extraction such as fishing Poaching of turtles and Gangetic river dolphin Industrial and domestic discharge into the river
7	1 species	-	-	89	Alteration of the river bank due to agriculture, sand mining. Unsustainable resource extraction such as fishing Poaching of turtles and Gangetic river dolphin Industrial and domestic discharge into the river
49	-	-	2 turtle species	25 species	Altered flow regime due to Farakka barrage Destructive fishing practices River bank alteration Industrial and domestic discharge into the river

Stretch IV: Chhapra to Kahalgaon, Bihar

This is a 296 km stretch from Chhapra, Saran District, to Kahalgaon, Bhagalpur District, in the state of Bihar (Figure 3.50). Falling within the Lower Gangetic Plain (7B), the Ganga River in this stretch is very wide, highly braided and meandering and passes through an extensive alluvial plain. The flow regime of this stretch is enhanced by substantial inflows from the Ghaghra and Son rivers near Chhapra and from the Gandak River at Patna. During summer months the depth ranged from 1.6 m to 33.2 m, with a mean value of 4.7 m. There are major human settlements at Chhapra, Patna, Munger and Bhagalpur. The urban population and rural communities in this region are dependent on the Ganga River for their daily ablutions, drinking water, irrigation, fishing, municipal and industrial water, sand mining and daily communte.

Biodiversity value and ecological significance

- a. A total of 141 sightings of Gangetic river dolphins and two smooth-coated otters were seen during the rapid assessment within this stretch. About 61 fish species and more than 80 phytoplankton and zooplankton species have been recorded from this stretch.
- b. About 65 km of this stretch from Sultanganj to Kahalgaon, is protected as Vikramshila Gangetic Dolphin Sanctuary.
- c. Large, stable islands, meanders and ox-bow lakes within this stretch provide a mosaic of habitats for the aquatic fauna.

Stretch V: Sahibganj to Rajmahal, Jharkhand

This is a 34 km stretch from Sahibganj to Rajmahal, Sahibganj District, in Jharkhand (Figure 3.50). Falling within the Lower Gangetic Plain (7B), the Ganga River in this stretch is very wide, highly braided and anastomosing with multichannel formation on an alluvial plain. Channel splitting takes place and convex sandbars are formed due to the low-energy channel flow and poor lateral stability. The flow regime of this stretch is enhanced by substantial inflows from the Kosi River near Katihar District, Bihar. During summer months the depth ranged from 1.5 m to 17.0 m, with a mean value of 5.8 m. There are major human settlements at Sahibganj and Rajmahal. The urban population and rural communities in this region are dependent on the Ganga River for their daily ablutions, religious activities and drinking water and for fishing and irrigation.

Biodiversity value and ecological significance

- a. Around seven sightings of Gangetic river dolphins were made in this stretch. Besides, a family of smooth-coated otter were also encountered. About 89 species of fish, 182 phytoplankton and 40 zooplankton species have been recorded from this stretch.
- b. The habitat mosaic of this stretch includes deep pools, sandbars and mid-river islands, which are potential habitats for freshwater turtles, island-nesting birds and dolphins.

Stretch VI: Baharampur to Barrackpore, West Bengal

This is a 246 km stretch from Baharampur, Murshidabad District to Barrackpore, North 24 Parganas District, West Bengal (Figure 3.50). This stretch falls within the Lower Gangetic Plain (7B), and the river in this stretch is highly meandering with convex sandbars and a few mid-river islands. The flow regime of this zone is enhanced by the feeder canal originating from Farakka Barrage and meeting the main channel, Hooghly at Ahiran, Murshidabad District. The Ganga River receives substantial inflows from the Mayurakshi and Ajay rivers at Bardhaman District. During summer months, the depth of the river ranged from 2.9 m to 31.0 m, with a mean value of 8.1 m. There are major human settlements at Baharampur, Katwa, Kalyani and Barrackpore. The urban population and rural communities in this region are dependent on the Ganga River for daily ablutions, religious activities and drinking water and for industrial water and irrigation.

Biodiversity value and ecological significance

- a. Around 49 sightings of Gangetic river dolphins were made in this stretch. A total of 25 fish, 44 phytoplankton and 21 zooplankton species have been recorded from this stretch.
- b. Meanders with convex sandbars and mid-river islands within this stretch provide a mosaic of habitats for the aquatic fauna.



THREATS TO THE BIODIVERSITY OF GANGA RIVER

The Ganga River, along with the Brahmaputra River, forms one of the largest and most diverse river systems of the world, spanning 10 biomes (Nilsson et al., 2005). The varied geomorphological features along the Ganga River, coupled with climatic variations, result in a continuous gradient of habitats for various life forms, right from the headwaters to its mouth.

This continuous yet varied gradient of habitats and life forms, results in differential availability of resources, which has been critical in shaping the local economy along the river. The Ganga River basin also hosts a dense population of diverse stakeholders having varied resource uses and are linked culturally, religiously, socially and economically to the river. The differential resource use by different stakeholders has resulted in varied impacts and threats along the length of the River. Therefore, for successful ecological restoration of the river, identification of site-specific threats and ways of minimizing or eliminating them is a prerequisite (Pressey et al., 2007; Palmer et al., 2010). Given this, the following sections describe the threats specific to the upper, middle and lower stretches of the Ganga River.



Upper Ganga


4.1

Located amidst the Himalayan mountain range, this stretch of the river is highly sensitive and ecologically fragile. The stretch is high in biodiversity value hosting habitats for endangered species such as otters, snow trout and golden mahaseer. The human population density is low (190 persons/km²) compared with the middle and lower stretches. The following are the specific threats faced by this stretch:

- Significant structural changes have occurred in this stretch due to 16 hydroelectric projects in the Bhagirathi and Alaknanda basins. The physical habitat of the river is further threatened by 14 projects under construction and 39 proposed projects in the Bhagirathi and Alaknanda basins.
- 70.7% of the Bhagirathi River and 48.0% of the Alaknanda River are affected by inundation and diversion, which have disrupted the longitudinal connectivity and water flow. As a result, 28.6% and 35.2% of the Bhagirathi and Alaknanda river channels have turned into ecological deserts (AHEC, 2011). A decline in the population of the golden mahaseer due to the Tehri Dam acting as a migration barrier has been noted upstream of the Bhagirathi River (Sharma, 2003).
- The flow regime in the headwaters is vulnerable to the retreat of the Gangotri glacier (Singh et al., 2017). The glacial melt is expected to increase summer flows for some years until the disappearance of the glacier, which would be followed by a reduction in the flow (Mall et al., 2006).

- The altered flows due to structural and climatic changes cause disruptions in the life history strategies of many in-stream organisms and affect the food web (Nautiyal, 2010).
- The climate and hydrological changes have transformed the thermal gradient in this stretch of the river, leading to a shrunken distribution range of cold water fish species such as *Schizothorax* sp. and upstream range extensions of several fish species such as *Cyprinus carpio* (Sarkar et al., 2012).

Figure 4.1
pH level in the
upper, middle
and lower
stretches of the
Ganga River



- Water quality of the this stretch as assessed by CPCB at 10 locations between 2003 and 2014 reveals that the pH ranged from 6.6 to 8.5. Over the study period, the DO showed decreasing trend, from 9.2 mg/L to 8.0 mg/L, and the BOD showed an increasing trend, from 1.3 mg/L to 1.9 mg/L. The BOD increased above the permissible limit (3 mg/L) near Satyanarayan temple, downstream of Raiwala and Haridwar, and the DO dropped below the critical point (5.4 mg/L) at Haridwar. During the present study, the pH was found to be in the alkaline range (8.05–8.79) between Uttarkashi and the Bhimgoda Barrage (Figure 4.1). The average DO concentration was 9.3 ± 0.3 mg/L (Figure 4.2), and the nitrate concentration was 2.2 ± 0.2 mg/L (Figure 4.3). Two active drains were observed downstream of Rishikesh Barrage and at Haridwar (Figure 4.4; Table 4. 1).

Critical locations

In this stretch, the nitrate concentration was highest (3.7 mg/L) Haridwar, downstream of Bhimgoda Barrage (Figure 4.4). It is evident that the organic pollutant load is higher at Haridwar, which could deplete the DO beyond the critical level (5 mg/L) and affect the aquatic biota.

Figure 4.2
DO (mg/L) in the
upper, middle
and lower
stretches of the
Ganga River




Figure 4.3
Nitrate concentration in the upper, middle and lower stretches of the Ganga River




Figure 4.4
Spatial distribution of threats to the biodiversity in the upper stretch of the Ganga River




Table 4.1
Active wastewater
inflow points along
the Ganga River

Sl. no.	Place	Geographic location		Type of wastewater
		Latitude	Longitude	
1	Anupshahr	28.35149	78.27200	Sewage treatment plant outlet
2	Ghatia Ghat, Farrukhabad	27.41547	79.62725	Sewage
3	Mehndi Ghat	27.01294	79.98461	Small drains
4	Bithur, Kanpur	26.61535	80.27442	Sewage
5	Kanpur city	26.50056	80.32013	Industrial effluent
6	Kanpur city	26.49638	80.32536	Sewage
7	Kanpur city	26.49147	80.33330	Sewage
8	Kanpur city	26.48378	80.35062	Sewage
9	Kanpur city	26.47454	80.36568	Sewage
10	Kanpur city	26.46869	80.37277	Sewage
11	Kanpur city	26.46567	80.37600	Sewage
12	Kanpur city	26.44514	80.39501	Industrial effluents
13	Kanpur city	26.43141	80.41152	Industrial effluents
14	Allahabad	25.50416	81.86130	Sewage
15	Allahabad	25.50338	81.87253	Sewage
16	Mirzapur	25.15864	82.58467	Sewage
17	Purana Setu, Varanasi	25.26774	83.01821	Sewage
18	St. Ravidas Park, Varanasi	25.28371	83.01180	Sewage
19	Harischandra Ghat, Varanasi	25.29786	83.00814	Sewage
20	Downstream of Rajghat Bridge, Varanasi	25.32575	83.03923	Sewage
21	Varuna confluence, Saray Mohana	25.32864	83.04503	Sewage
22	Jhandatar, Ghazipur	25.58135	83.59493	Sewage
23	Near Ghazipur Bridge	25.58508	83.60267	Sewage
24	Buxar	25.58526	83.98525	Sewage
25	Danapur	25.64593	85.05993	Sewage
26	Downstream of Danapur	25.64706	85.11176	Sewage
27	Patna	25.62233	85.16379	Sewage
28	Barh	25.48718	85.70846	Sewage
29	NTPC, Barh	25.50471	85.75761	NTPC outlet
30	NTPC, Barauni	25.36854	86.00635	NTPC outlet
31	Jamalpur	25.37152	86.54046	Sewage
32	Kahalgaon	25.26631	87.22543	Sewage
33	Farakka	24.75905	87.90965	NTPC outlet

Sl. no.	Place	Geographic location		Type of wastewater
		Latitude	Longitude	
34	Azimganj-Jiaganj	24.24178	88.25825	Sewage
35	Berhampore	24.11547	88.24617	Sewage
36	Kashiganj, Katwa	23.64273	88.14476	Sewage
37	Nabadwip	23.41174	88.37898	Sewage
38	Tribeni NTPC outlet	22.99142	88.40703	NTPC outlet
39	Tribeni	22.97104	88.40680	Sewage
40	Kalyani	22.92951	88.40748	Sewage
41	Downstream of Kalyani	22.91978	88.40249	Sewage
42	Chandan Nagar	22.85857	88.37097	Sewage
43	Bandel	22.84007	88.37442	Sewage
44	Bandel	22.82117	88.35438	Sewage
45	Downstream of Bandel	22.78862	88.34006	Sewage
46	NTPC, Barrackpore	22.72688	88.35859	NTPC outlet
47	Uttarpara	22.67203	88.35517	Sewage
48	Near Bally Bridge	22.66410	88.35332	Sewage
49	Bally	22.63855	88.35870	Sewage
50	Downstream of Bally	22.61497	88.36162	Sewage
51	Bag Bazar	22.60837	88.35798	Sewage
52	Bata Nagar	22.51383	88.21169	Sewage

Middle Ganga

4.2

This stretch is characterized by large, fertile floodplains that are extensively used for agriculture. The original vegetation has been replaced by crops. The human density is about 820 persons/km². This stretch hosts a significant populations of the Gangetic river dolphin, gharial, mugger, turtles and island-nesting birds. Threats specific to this stretch include the following.

- Water abstraction from the Upper Ganga Canal at Bhimgoda, Haridwar, Middle Ganga Canal at Bijnor and Lower Ganga Canal and Parallel Lower Ganga Canal at Narora has reduced the flow of the Ganga River to 10% of the natural flow regime, resulting in disruption of the horizontal and longitudinal connectivity between habitats for the priority species.
- Bank feature alteration due to agriculture, construction and sand mining has disrupted the lateral connectivity of the river. Extensive sandbar cultivation has rendered the habitat unsuitable for use as nesting sites by turtles and island-nesting birds.
- Unsustainable biological resource extraction using destructive methods is ubiquitous in this stretch of the Ganga River.
- Analysis of the water quality parameters assessed by CPCB between 2003 and 2014 for this stretch at 20 locations reveals that the pH ranged from 7.3 to 8.8. The DO showed an overall increasing trend from 7.4 mg/L to 7.9 mg/L. Subsequently the BOD showed decreasing trend from 4.31 mg/L to 3.2 mg/L, which is higher than the permissible limit (3 mg/L). The BOD was well above the permissible limit at Narora, Kannauj, Kanpur, Dalmau, Kala Kankar, Allahabad and Varanasi. During the present study

the pH ranged from 7.4 mg/L to 8.8 mg/L between Bijnor and Varanasi (Figure 4.1). The mean nitrate concentration was low, at 1.6 ± 0.2 mg/L (Figure 4.3), which resulted in an optimal mean DO concentration of 7.5 ± 0.2 mg/L (Figure 4.2). Within this stretch, there were 21 active wastewater inflow points (mainly raw sewage and industrial discharge) (Figure 4.5; Table 4.1).

Figure 4.5
Spatial
distribution of
threats to the
biodiversity in the
middle stretch of
the Ganga River



Kanpur city accounted for the greatest number of discharge points (9) (Figure 4.5). Indiscriminate use of synthetic chemicals in the agricultural and healthcare sectors releases persistent pollutants (PPs) such as heavy metals, polychlorinated bi-phenyls (PCBs) and perfluorinated compounds (PFCs). Both PCBs and PFCs are endocrine disrupting chemicals (EDCs) as these interfere with the hormonal and immune systems of aquatic species through bioaccumulation and biomagnification along the food chain (Bergman et al., 2012; Senthilkumar et al., 1999) as has been reported in the food web of the Gangetic river dolphin (Yeung et al., 2009).

Critical locations

- The DO was critically low (1.1 mg/L) at the confluence of the Varuna sewage drain and the Ganga River, near Varanasi (Figure 4.2). The nitrate concentration was higher (5.3 mg/L), near Raona village, in Varanasi, where the output of the Mughal Sarai sewage treatment plant runs into the Ganga River (Figure 4.3). The mixing of sewage at this location might have created eutrophic conditions and elevated the DO to 10.1 mg/L.
- In this stretch of the river, area between Farrukabad and Kanpur and between Allahabad and Varanasi have high concentrations of all the threats to the aquatic diversity (Figure 4.5).

Lower Ganga

4.3


The floodplains and the natural vegetation of this stretch of the river have also been extensively replaced by croplands. This is the most densely populated of the three stretches (950 persons/km²). Representative animal species of this stretch include the Gangetic river dolphin, otters, the estuarine crocodile and turtles, particularly *Batagur baska*. The following are the threats specific to this stretch:

- The Farakka Barrage has resulted in structural changes and modified flow regimes, leading to sediment and nutrient accretion. Increased freshwater inflow from the Farakka Barrage into the Hooghly River has significantly changed the salinity regime, water transparency, suspended sediments and nutrient load and altered the freshwater fish assemblage. The freshwater fishes (such as *Apocryptes bato*, *Goniolosa manminna* and *Labeo* sp.) that were found only from Nabadwip to Nawabganj in West Bengal during the pre-Farakka Barrage period are now available in the Godakhali-Uluberia region, signifying a shift of at least 60 km of the freshwater zone towards the mouth of the estuary (Manna et al., 2013).
- Unsustainable biological resource extraction is also widespread in this stretch of the river. Destructive fishing techniques and intentional capture of Gangetic river dolphins, turtles and storks (Choudhary & Mishra, 2006) have led to mortality and reduced populations of these species. *Batagur baska* and *Glyptis gangeticus* have also extirpated from this stretch due to indiscriminate killing (Compagno, 2007; Moll et al., 2009).
- Bank feature alteration due to land-use changes such as agriculture, construction activities and sand mining has disrupted the lateral connectivity of the river in the floodplain. Sandbar cultivation in the states of Bihar and Jharkhand has rendered the habitat unsuitable for use as nesting sites by turtles and island-nesting birds.
- A comparison of water quality parameters from 2003 to 2014 as assessed by CPCB in this stretch at 35 locations reveal that the pH ranged from 7.3 to 8.8. The DO showed an increasing trend from 7.2 mg/L to 7.5 mg/L, and subsequently the BOD showed decreasing trend from 2.4 mg/L to 2.8 mg/L. Ghazipur, in Uttar Pradesh, and Baharampur, Ghoshpara, Serampore, Dakshineswar, Shivpur, Garden Reach, Uluberia, Palta and Diamond Harbour in West Bengal were critical locations where the BOD was well above the permissible limit. The water quality parameters assessed during the present study from Ghazipur to Diamond Harbour reveal that the pH was in the alkaline range (7.4–8.5) (Figure 4.1). The average nitrate concentration was high (4.3 ± 0.5 mg/L) (Figure 4.3), which lowered the average DO to 6.2 ± 0.1 mg/L (Figure 4.2). 31 active wastewater inflow points (raw sewage and industrial discharge) were observed. (Figure 4.6; Table 4.1).
- The city of Kolkata produces 5114 tonnes/day of solid waste, of which 10% is plastic (Das & Bhattacharya, 2013; Singh & Francis, 2017). This untreated plastic waste disrupts the biodiversity (Dey et al., 2014) and reaches the Bay of Bengal at a rate of 0.12 million tonnes per year (Lebreton et al., 2017).

Critical locations

- In the Lower Ganga, the first critical stretch is Sitab Diara to Ajimganj/Jiaganj, which includes critical locations, viz. Munger, Kahalgaon, Sahibganj, Revelganj and Ajimganj/Jiaganj, where the nitrate level was > 10 mg/L (Figures 4.3 and 4.6). The second critical stretch falls between Barrackpore and Falta (Figure 4.6), where the major wastewater inflow points elevated the nitrate level and lowered the DO concentration below 5 mg/L at Sujapur, Batanagar, Shibpur, Garden Reach, Uluberia and Falta.

Figure 4.6
Spatial
distribution of
threats to the
biodiversity in the
lower stretch of
the Ganga River



Conclusions

4.4

It is evident from the plethora of literature that the ecological integrity of the Ganga River is threatened throughout its course. Each stretch has its unique set of ecological assemblage and threats. The upper stretch of the river is threatened by structural changes in its morphology due to altered flow regime and climate change. Bank alteration, unsustainable resource extraction and changes in water quality are dominant threats in the middle and lower stretches of the river. Thus, it can be deduced that alteration of the structural morphology of the river is the key threat, leading to habitat degradation and biodiversity loss. These existing threats are further heightened by the impacts of climate change and altered water quality, questioning the survival of the aquatic species.

In order to maintain the ecological integrity of the river, it is crucial to address the stretch-specific threats, keeping in view the current global river conservation scenario. In the upper stretch of the Ganga River, the focus should be on maintaining the natural structural features of the river, and micro-hydel power projects should be considered instead of larger dams in this stretch. In the middle and lower stretches of the river, restoration of the hydrology regime through flow augmentation to a near-natural condition would be the key to bringing about ecological restoration in the river ecosystem. A holistic approach should be adopted for the assessment of 'flow-ecology' relationships of designated aquatic umbrella species and then integrate the eco-hydrological requirements of these species during the conception of nature-oriented and ecologically relevant e-flow targets (Hussain et al., 2011). This flow restoration would, in turn, enhance the suitability of the habitat for the aquatic biota, improve the flow regime and retain enough water in the Ganga River to dilute the heavy pollution load.

The concentration of nitrate in natural rivers is typically low (< 1 mg/L). The preliminary assessment of the water quality of the Ganga River reveals that the wastewater inflow has degraded the water quality at specific locations and elevated the nitrate and BOD concentrations above critical thresholds. An excess organic load, especially nitrate, can rapidly deplete the dissolved oxygen concentration and become toxic to aquatic organisms, mainly fishes, at > 10 mg/L or a higher level. Hence, the critical locations identified with respect to deteriorated water quality should be the focus of intervention efforts.



CONSERVATION IMPLICATIONS

The Ganga River is a repository of various life forms, resulting from the variations in its geomorphological and fluvial processes. This diversity of species or life forms is crucial for reducing the risk of ecological collapse and maintaining ecosystem services, which are essential for human well-being. The current state of the Ganga River as evident from the rapid biodiversity assessment, coupled with an extensive review of secondary literature, highlights the need for its ecological restoration, whilst considering the varied geomorphological, climatic and ecological features and societal aspects. An effective ecological restoration plan essentially needs to consider the biotic and abiotic components of the ecosystem, viz. species composition and abundance, the characteristics of the habitat and its use by species, an assessment of the threats faced by important taxa and distinct spatial zones requiring differential treatment (McDonald et al., 2016). In view of the foregoing discussion, the following points have been inferred for the conservation of aquatic diversity aiming at rejuvenation of the Ganga River.



Stretch-specific implications

- The differential use of resources in the different stretches of the river have given rise to diverse threats, and hence measures to address these threats should be stretch-specific. Flow maintenance, preservation of umbrella species and identification of the priority areas are key concepts that can be applied to the assessment of stretch-specific requirements.
- The upper stretch of the Ganga River has undergone structural changes and is unlikely to revert to its natural hydrological patterns. Construction of further hydroelectric projects in this stretch will compromise the ecological, religious and recreational values of the river. This can be averted by incorporating sustainable and efficient water use practices in norms with modern abstraction techniques, which should be reflected in the water policy and allocation guidelines.
- The middle and lower stretches of the Ganga River are facing severe changes in water quality due to the wastewater from domestic and industrial sources and agricultural run-off. Altered water quality is aggravated by the low flow condition. Intensive agriculture along the banks of the river and on the river islands and sand mining in these stretches has altered the habitat for island-nesting species of bird and turtle. The lower stretch of the Ganga River is also highly modified by the Farakka Barrage, leading to flow alterations and changes in the salinity regimes in the estuarine areas, causing changes in the species assemblage. Addressing these problems is not only a river conservation issue but also a social one owing to the high dependence of the dense populations of the local communities living along the banks of the river.

River hydrology

- Assessment of the hydrology of the Ganga River revealed that the channel depth is a major limiting factor for the distribution of the species. Only 38.7% of the river had a suitable depth for the umbrella species, viz. the Gangetic river dolphin and gharial, during summer months.
- As the natural flow cannot be restored due to the highly fragmented state of the river, restoration of the catchment area and periodic reservoir releases should be carried out.
- Prioritizing efficient water usage in consonance with the needs of aquatic species through an ecologically sound and economically just water budget and policy would augment flows.
- The flow restoration would enhance the habitat suitability for the aquatic fauna and retain sufficient water in the river to dilute the pollution load.

Species restoration

- It is evident from literature that the populations of the umbrella species, such as the Gangetic river dolphin, otters, gharial and mugger in the Ganga River have become highly fragmented over time. Historically, these species had wider distributions in the Ganga River and its tributaries. However, due to the loss of longitudinal and lateral connectivity and unsuitable habitat conditions, they are now restricted to small stretches of the river where suitable habitat persists.
- The rapid biodiversity assessment revealed significant distribution of several of these umbrella species and their habitats, albeit in particular stretches of the Ganga River, including the PAs. Based on the abundance and distribution of these select aquatic species, areas with higher biodiversity values were identified. These areas have been recommended for in-depth studies and planning ecological restoration of the Ganga River.
- Crocodilians and turtles are highly sensitive to anthropogenic habitat alterations and are therefore conservation-dependent species. However, their habitats along the Ganga River are not protected and these species are susceptible to regular human disturbances. Bank alteration, sandbar cultivation and sand mining have resulted in competitive use of sandy banks and loss of basking and nesting beaches, threatening the survival of these taxa. Efforts to conserve these species should emphasize science-based restoration with the support of the local communities. Therefore, aligning

Figure 5.1
Protected areas
along the Ganga
River



conservation initiatives with local livelihoods is a prerequisite of the restoration process and is currently being carried out under the community-based conservation component of the project. Awareness programmes and rescue programmes for species in distress are needed to protect them.

- Amphibians are the least studied group along the Ganga River, hence, information pertaining to them is either unavailable or scanty. Therefore, extensive studies are needed to assess the population trends of these species in the Ganga River and its basin.
- The Ganga basin has several seasonally flooded wetlands, permanent lakes and marshes, which serve as wintering grounds for migratory waterbirds. Some of these areas have been declared as PAs, for e.g., Jhilmil Conservation Reserve in Uttarakhand, Surhataal Wildlife Sanctuary in Uttar Pradesh, Kusheshwar Asthan Bird Sanctuary in Bihar and Udhwa Bird Sanctuary in Jharkhand. These wetlands are kidneys of the landscapes and need restoration measures. The migratory waterbirds use sandy islands and sand banks of the Ganga River for roosting and the resident water birds such as Indian skimmer, Little pratincole and River terns for breeding, hence these habitats need to be restored and protected.

Emergent situations

- Incidental mortality of the priority species by entanglement in fishing nets, direct hit by propellers of the vessels and boats, poaching and a lack of concerted efforts to address these issues are major concerns.
- In order to be prepared for such situations in the future, rescue protocols involving stakeholders as well as a network of rescue & rehabilitation centres need to be established along the river. Work has already been initiated in this regard by the rescue & rehabilitation component of the project.
- For the long-term success of this operation, it is important to build the capacity of the frontline staff of the forest department, local communities and other stakeholders of the five Ganga River states so that they are able to successfully carry out rescue operations of priority aquatic species. These activities are currently being carried out under the training component of the NMCG-WII project.

Way forward

- Planning for ecological restoration of an ecosystem starts with exploring its historical and present day conditions, for which a literature review and rapid biodiversity assessment were carried out. These also helped identify areas with high biodiversity values.
- The PAs along the Ganga River account for around 10% of the total river stretch including parts of Sunderban Tiger Reserve (Figure 5.1). These areas provide relatively undisturbed habitats along the river and are repositories of aquatic umbrella species such as Gangetic river dolphin, otters and crocodilians. Increasing the PA network by covering biodiversity-rich stretches of the Ganga River through strategic planning and with the involvement and support of local communities should be incorporated in the conservation process.
- The water quality of the Ganga River has been severely degraded due to domestic, industrial and agricultural sediment accretion. Evidence of bioaccumulation and biomagnification has also been documented in the food web of major aquatic species. Further aquatic ecotoxicological research is needed to assess the implications of altered water quality for the long-term survival of the species.
- Impacts of climate change are exacerbating threats in the form of unpredictable flow regime and changes in both the upstream and downstream distribution range of species. The restoration process should thus take into account the externalities caused by climate change, using future scenario modelling techniques to make climate change projections and evaluate their impacts on species and habitats.
- Planning river restoration is not only a conservation issue but also a social one owing to the high dependence of the dense human population living along the banks of the Ganga River. Thus, efforts are under way to create awareness among stakeholders and to garner local support for conservation. Site-specific strategies are being developed by aligning the livelihoods of the local people with conservation priorities. These local communities are also being made aware about the role of an ecologically intact Ganga River in enhancing the quality of life.

REFERENCES

- Abell, R., Allan, J.D., & Lehner, B. (2007). Unlocking the potential of protected areas for freshwaters. *Biological Conservation*, 134(1), 48-63.
- AHEC (Alternate Hydro Energy Center), IIT (Indian Institute of Technology) Roorkee (2011). Assessment of Cumulative Impacts of Hydroelectric Projects on Aquatic and Terrestrial Biodiversity in Alaknanda and Bhagirathi Basins, Uttarakhand, India. Report submitted to Ministry of Environment, Forest & Climate Change, Government of India.
- Anderson, J. (1871). On some Indian reptiles. In *Proceedings of the Zoological Society of London*, 11, 149-211.
- Anderson, J. (1878). *Anatomical and Zoological Researches: Comprising an Account of Zoological Results of the Two Expeditions to Western Yunnan in 1868 and 1875; and a Monograph of the Two Cetacean Genera, Platanista and Orcella*. London: Bernard Quaritch.
- Andrews, H. (2006). Status of the Indian Gharial (*Gavialis gangeticus*), Conservation Action and Assessment of Key Locations in North India. Report to the Madras Crocodile Bank Trust.
- Annandale, N. (1909). Reports on the fishes taken by the Bengal fisheries steamer "Golden Crown". Part I, Batoidei. *Indian Museum Memoirs*, 2(1), 1-60.
- Annandale, N. (1912). The Indian mud turtles (Trionychidae). *Records of the Indian Museum*, 7(2), 151-180.
- Annandale, N. (1923). Animal life of the Ganges. *Journal of the Bombay Natural History Society*, 29, 633-642.
- Anon (2015). Annual Survey Report on Gharial and Mugger in the National Chambal Sanctuary, Madhya Pradesh. Report submitted to Conservator of Forest, Gwalior, Rahim, V., Pateria, A., & Sharma, R.K. (Eds.).
- Atkinson, E.T. (1882). *The Himalayan Districts of the North Western Province of India*. Allahabad. Reprinted in 1996 as the Himalayan Gazetteer, Vol. 2, Part I. Dehradun: Natraj Publishers.
- Atkinson, E.T. (1974). *Fauna of the Himalayas, Containing Species of Kumaon, Garhwal, Nepal and Tibet*. New Delhi: Cosmo Publications.
- Atkore, V.M. (2005). Conservation status of fishes in the tributaries of Ramganga with special reference to golden mahseer (*Tor putitora*) Hamilton. M.Sc. dissertation, Saurashtra University, Rajkot, 67 pp.
- Atkore, V.M., Sivakumar, K., & Johnsingh, A.J.T. (2011). Patterns of diversity and conservation status of freshwater fishes in the tributaries of River Ramganga in the Shiwaliks of the Western Himalaya. *Current Science*, 731-736.
- Bahuguna, V.K., Swaminath, M.H., Tripathi, S., Singh, T.P., Rawat, V.R.S., & Rawat, R.S. (2016). Revisiting forest types of India. *International Forestry Review*, 18(2), 135-145.
- Bal, D.V., & Rao, K.V. (1984). *Marine Fisheries*. New Delhi: Tata McGraw-Hill.
- Balian, E.V., Segers, H., Lévêque, C., & Martens, K. (2008). The freshwater animal diversity assessment: An overview of the results. *Hydrobiologia*, 595(1), 627-637.
- Baruah, C., Devi, P., & Sharma, D.K. (2016). Comparative morphometry and biogeography of the freshwater turtles of genus *Pangshura* (Testudines: Geoemydidae: *Pangshura*). *International Journal of Pure and Applied Zoology*, 4(1).
- Bashir, T., Behera, S.K., Khan, A., & Gautam, P. (2012). An inventory of mammals, birds and reptiles along

- a section of the river and banks of Upper Ganges, India. *Journal of Threatened Taxa*, 4(9), 2900-2910.
- Beechie, T., Pess, G., Roni, P., & Giannico, G. (2008). Setting river restoration priorities: A review of approaches and a general protocol for identifying and prioritizing actions. *North American Journal of Fisheries Management*, 28(3), 891-905.
- Behera, S.K. (1995). Studies on population dynamics, habitat utilization and conservation aspects of Gangetic dolphin (*Platanista gangetica*) in a stretch of Ganga River from Rishekesh to Kanpur. Ph.D. thesis, School of Studies in Zoology, Jiwaji University Gwalior, 198 pp.
- Behera, S.K., Singh, H., & Sagar, V. (2013). Status of Ganges river dolphin (*Platanista gangetica gangetica*) in the Ganga River basin, India: A review. *Aquatic Ecosystem Health & Management*, 16(4), 425-432.
- Berger, J. (1997). Population constraints associated with the use of black rhinos as an umbrella species for desert herbivores. *Conservation Biology*, 11(1), 69-78.
- Bergman, Å., Rydén, A., Law, R.J., de Boer, J., Covaci, A., Alaei, M., & Van den Eede, N. (2012). A novel abbreviation standard for organobromine, organochlorine and organophosphorus flame retardants and some characteristics of the chemicals. *Environment International*, 49, 57-82.
- Bhaduria, R.S., Singh, A.N. & Mishra, S.B. (1995). Morphometrics and taxonomy of spotted black terrapin *Geoclemys hamiltonii* (Gray 1831). *Tigerpaper*, 22(3), 27-30.
- Bhuiyan, A.L. (1964). *Fishes of Dacca* (No. 13). Asiatic Society of Pakistan.
- Bilgrami, K.S. (1991). Biological profile of the Ganga: Zooplankton, fish, birds, and other minor fauna. In Krishna Murti, C.R., Bilgrami, K.S., Das, T.M., & Mathur, R.P. (Eds.). *The Ganga: A Scientific Study*, (pp. 83-94). New Delhi: Ganga Project Directorate, Ministry of Environment & Forests, Government of India.
- BirdLife International (2016). *Mycteria leucocephala*. The IUCN Red List of Threatened Species 2016. www.birdlife.org (accessed on 13 June 2017).
- BirdLife International (2017). Important Bird Areas factsheet: Narora. www.birdlife.org (accessed on 13 June 2017).
- Bista, D., & Shah, K. (2010). Diversity and status of the turtles in Ghodaghodi Lake Area, Kailali District, Far West Nepal. *Journal of Natural History Museum*, 25, 366-373.
- Blanford, W.T. (1881). *Fauna of British India, Including Ceylon and Burma: Mammalia*. London: Taylor and Francis.
- Boulenger, G.A. (1890). *The Fauna of British India Including Ceylon and Burma: Published Under the Authority of the Secretary of State for India in Council. Reptilia and Batrachia*. London: Taylor and Francis.
- Bustard, H. R. (1980). The Government of India Crocodile Project. *Cheetal*, 22, 11-16.
- Bustard, H.R. (1984). Breeding the gharial *Gavialis gangeticus*: captive breeding a key conservation strategy for endangered crocodilians. Symposium of the Zoological Society of London 52: 385-406.
- Bustard, H.R., & Choudhury, B.C. (1981). Conservation future of saltwater crocodile (*Crocodylus porosus Schneider*) in India. *Journal of the Bombay Natural History Society*, 77(2), 201-214.
- Carrizo, S.F., Smith, K.G., & Darwall, W.R.T. (2013). Progress towards a global assessment of the status of freshwater fishes (Pisces) for the IUCN Red List: Application to conservation programmes in zoos and aquariums. *International Zoo Yearbook*, 47(1), 46-64.
- Casavas, I., Doulman, D.J., Petr, T.O., Padro, J., Debas, L. (1996). *Cambodia: Rehabilitation and Development Needs of the Fishery Sector*. Rome: FAO.

- Champion, S.H., & Seth, S.K. (1968). *A Revised Survey of the Forest Types of India*. Dehradun: Natraj Publishers.
- Chandler, R.J. (2009). *Shorebirds of the Northern Hemisphere*. Christopher Helm.
- Choudhary, D.N., & Mishra, A. (2006). Sighting of some threatened bird species in Vikramshila Gangetic Dolphin Sanctuary (VGDS), Bhagalpur, Bihar. *Newsletter for Birdwatchers*, 46(5), 68-70.
- Choudhary, S., Dey, S., Dey, S., Sagar, V., Nair, T., & Kelkar, N. (2012). River dolphin distribution in regulated river systems: Implications for dry-season flow regimes in the Gangetic basin. *Aquatic Conservation: Marine and Freshwater Ecosystems*, 22(1), 11-25.
- Choudhary, S.K., Smith, B.D., Dey, S., Dey, S., & Prakash, S. (2006). Conservation and biomonitoring in the Vikramshila Gangetic Dolphin Sanctuary, Bihar, India. *Oryx*, 40(2), 189-197.
- Chaudhuri, B.L. (1911). Freshwater stingrays of the Ganges. *Journal and Proceedings of the Asiatic Society of Bengal*, 3, 625-629.
- Chaudhuri, B.L. (1912). Aquatic tortoises of the middle Ganga and Brahmaputra. *Records of the Indian Museum*, 7, 212-214.
- Choudhury, B.C. (1998). Data on sarus crane. *Newsletter for Birdwatchers*, 38(4), 69.
- Chowfin, S.M., & Leslie, A.J. (2013). A preliminary investigation into nesting and nest predation of the Critically Endangered gharial (*Gavialis gangeticus*) at Boksar in Corbett Tiger Reserve, Uttarakhand, India. *International Journal of Biodiversity and Conservation*, 5(2), 54-57.
- CIFRI (Central Inland Fisheries Research Institute) (1964). Annual Report for 1963-1964. Barrackpore, West Bengal.
- Collen, B., Loh, J., Whitmee, S., McRAE, L., Amin, R., & Baillie, J.E. (2009). Monitoring change in vertebrate abundance: The Living Planet Index. *Conservation Biology*, 23(2), 317-327.
- Collen, B., Whitton, F., Dyer, E.E., Baillie, J.E., Cumberlidge, N., Darwall, W.R., & Böhm, M. (2014). Global patterns of freshwater species diversity, threat and endemism. *Global Ecology and Biogeography*, 23(1), 40-51.
- Collins, R.A., Armstrong, K.F., Meier, R., Yi, Y., Brown, S.D., Cruickshank, R.H., & Johnston, C. (2012). Barcoding and border biosecurity: Identifying cyprinid fishes in the aquarium trade. *PloS One*, 7(1), e28381.
- Compagno, L.J.V. (2007). *Glyphis gangeticus*. The IUCN Red List of Threatened Species 2007. www.iucnredlist.org (accessed on 15 November 2017).
- CPCB (Central Pollution Control Board) (2013). *Pollution Assessment: River Ganga*. New Delhi: Ministry of Environment and Forests, Government of India.
- Daniel, J.C. (1970). A review of the present status and position of endangered species of Indian reptiles. In: *Proceedings of the IUCN 11th Technical Meeting, New Delhi* (pp. 75-76). IUCN Publications Series No. 18.
- Daniel, J.C. (2002). *The Book of Indian Reptiles and Amphibians*. Oxford University Press.
- Daniel, J.C., & Shull, E.M. (1964). A list of the reptiles and amphibians of Surat Dangs, South Gujarat. *Journal of the Bombay Natural History Society*, 60, 737-743.
- Darwall, W.R., Holland, R.A., Smith, K.G., Allen, D., Brooks, E.G., Katarya, V., & Cuttelod, A. (2011). Implications of bias in conservation research and investment for freshwater species. *Conservation Letters*, 4(6), 474-482.
- Das, I. (1987). Status and distribution of estuarine turtles in India. *Marine Fisheries Information Service, Technical and Extension Series*, 72, 21-22.

- Das, I. (1988). New locality record for the Indian Peacock Softshell Turtle *Trionyx hurum*. *Journal of the Bombay Natural History Society*, 84(3), 691-692.
- Das, I. (1991). *Colour Guide to the Turtles and Tortoises of the Indian Subcontinent*. R & A Publishing.
- Das, I. (1995). *Turtles and Tortoises of India*. Bombay: Oxford University Press.
- Das, S. (2014). Ganga - Our endangered heritage. In Sanghi, R. (Ed.), *Our National River Ganga* (pp. 45-74). Cham: Springer.
- Das, I., Basu, D., & Singh, S. (2010). *Nilssonina hurum* (Gray 1830): Indian peacock softshell turtle. In: Rhodin, A.G.J., Pritchard, P.C.H., van Dijk, P.P., Saumure, R.A., Buhlmann, K.A., Iverson, J.B., & Mittermeier, R.A. (Eds.). *Conservation Biology of Freshwater Turtles and Tortoises: A Compilation Project of the IUCN/SSC Tortoise and Freshwater Turtle Specialist Group*. Chelonian Research Monographs No. 5, 048.1-048.6.
- Das, I., & Bhupathy, S. (2009a). *Melanochelys trijuga* (Schweigger 1812): Indian black turtle. In: *Conservation Biology of Freshwater Turtles and Tortoises: A Compilation Project of the IUCN/SSC Tortoise and Freshwater Turtle Specialist Group*. Chelonian Research Monographs No. 5.
- Das, I., & Bhupathy, S. (2009b). *Hardella thurjii* (Gray 1831): Crowned river turtle. In: *Conservation Biology of Freshwater Turtles and Tortoises: A Compilation Project of the IUCN/SSC Tortoise and Freshwater Turtle Specialist Group*. Chelonian Research Monographs No. 5.
- Das, I., & Singh, S. (2009). *Chitra indica* (Gray 1830): Narrow-headed softshell turtle. In: Rhodin, A.G.J., Pritchard, P.C.H., van Dijk, P.P., Saumure, R.A., Buhlmann, K.A., Iverson, J.B., & Mittermeier, R.A. (Eds.). *Conservation Biology of Freshwater Turtles and Tortoises: A Compilation Project of the IUCN/SSC Tortoise and Freshwater Turtle Specialist Group*. Chelonian Research Monographs No. 5, 027.1-027.7.
- Das, M.K., Sharma, A.P., Vass, K.K., Tyagi, R.K., Suresh, V.R., Naskar, M., & Akolkar, A.B. (2013). Fish diversity, community structure and ecological integrity of the tropical river Ganges, India. *Aquatic Ecosystem Health & Management*, 16(4), 395-407.
- Das, S., & Bhattacharyya, B.K. (2013). Municipal solid waste characteristics and management in Kolkata, India. In: *The 19th International Conference on Industrial Engineering and Engineering Management* (pp. 1399-1409). Berlin, Heidelberg: Springer.
- Day, F. (1878). *The Fishes of India: Being a Natural History of the Fishes Known to Inhabit the Seas and Fresh Waters of India, Burma, and Ceylon*, Vol. I. London: Bernard Quaritch, Norman & Son.
- de Vos, A. (1984). Crocodile conservation in India. *Biological Conservation*, 29(2), 183-189.
- del Hoyo, J., Elliott, A., & Sargatal, J. (1996). *Handbook of the Birds of the World, 3, Hoatzin to Auks*. Barcelona: Lynx Edicions.
- Delany, S., & Scott, S., IV (Eds.) (2006). *Waterbird Population Estimates*. Wageningen, The Netherlands: Wetlands International.
- Devaprakash, J., (2015). Project Turtle: Narora. *Science Reporter*, 43-45.
- Dey, S., Dey, S., & Kelkar, S.K.C.N. (2014). An annotated bird checklist of the Vikramshila Gangetic Dolphin Sanctuary, Bhagalpur, Bihar, India, with an assessment of threats to bird conservation. *Forktail*, 30(30), 34-40.
- Dhammika, S. (1993). *The Edicts of King Asoka: An English Rendering*. The Wheel Publication, 17 pp.
- Dhu, S., (1923). *The Angler in India or the Mighty Mahseer*. Dehradun: Natraj Publishers.
- Dudgeon, D., Arthington, A.H., Gessner, M.O., Kawabata, Z.I., Knowler, D.J., Lévêque, C., & Sullivan, C.A. (2006). Freshwater biodiversity: Importance, threats, status and conservation challenges. *Biological Reviews*, 81(2), 163-182.

- Duffy, J.E., Cardinale, B.J., France, K.E., McIntyre, P.B., Thébault, E., & Loreau, M. (2007). The functional role of biodiversity in ecosystems: Incorporating trophic complexity. *Ecology Letters*, 10(6), 522-538.
- Dwivedi, A.C. (2009). Ecological assessment of fishes and population dynamics of *Labeo rohita* (Hamilton), *Tor tor* (Hamilton) and *Labeo calbasu* (Hamilton) in the Paisuni river. *Aquacult*, 10(2), 249-259.
- Dwivedi, A.C. (2013). Population dynamics, age, growth and sex ratio of *Labeo bata* (Hamilton) from the middle stretch of the Ganga river. *India Flora and Fauna*, 19(1), 133-137.
- Dwivedi, A.C., Mishra, A.S., Mayank, P., & Tiwari, A. (2016). Persistence and structure of the fish assemblage from the Ganga river (Kanpur to Varanasi section), India. *Journal of Geography and Natural Disasters*, 6(159), 2167-0587.
- Dwivedi, A.C., & Nautiyal, P. (2010). *Population Dynamics of Important Fishes in the Vindhyan Region, India*. Lap Lambert Academic Publishing GmbH & Co. KG, 220 pp.
- Dwivedi, A.C., Singh, K.R., & Mayank, P. (2007). Age structure and growth of *Labeo calbasu* (Hamilton) from the Ghaghara river in Faizabad region (U.P.). *Journal of Natural Resource and Development*, 2(1), 49-54.
- Dwivedi, A.C., Tewari, N.P., & Singh, K.R. (2004). Present structure of capture and culture fishery of the Faizabad District (U.P.). *Bioved*, 15(1, 2), 95-98.
- Edds, D. (1998). Distribution of *Kachuga smithii smithii*, *Kachuga tecta* and *Kachuga tentoria flaviventer*. *Herpetological Review*, 29(2), 109.
- Elton, C. (1958). *The Ecology of Invasions by Plants and Animals*. London: Chapman & Hall.
- Erlinge, S. (1972). Interspecific relations between otter *Lutra lutra* and mink *Mustela vison* in Sweden. *Oikos*, 327-335.
- Ernst, C.H., Altenburg, R.G.M., & Barbour, R.W. (1997). *Turtles of the World*. The Netherlands: ETI Information Systems Ltd.
- FAO (1974). India: A Preliminary Survey of the Prospects for Crocodile Farming (based on the work of H.R. Bustard, Consultant). Rome: FAO.
- Feld, C.K., Birk, S., Bradley, D.C., Hering, D., Kail, J., Marzin, A., & Pont, D. (2011). From natural to degraded rivers and back again: A test of restoration ecology theory and practice. In: *Advances in Ecological Research* 44 (pp. 119-209). Academic Press.
- Fenberg, P.B., & Roy, K. (2008). Ecological and evolutionary consequences of size-selective harvesting: How much do we know? *Molecular Ecology*, 17(1), 209-220.
- Froese, R., & Pauly, D. (Eds.) (2017). FishBase. World Wide Web electronic publication. www.fishbase.org (accessed on 8 March 2018)
- Gardner, M.R., & Ashby, W.R. (1970). Connectance of large dynamic (cybernetic) systems: Critical values for stability. *Nature*, 228(5273), 784.
- Ghosh, A.K. (1991). The Ganga: A profile and biological resources. *Faunal Resources of Ganga*, Part I (pp. 1-12). Calcutta: Zoological Survey of India.
- Giri, S.S., Sahoo, S.K., Sahu, B.B., Sahu, A.K., Mohanty, S.N., Mukhopadhyay, P.K., & Ayyappan, S. (2002). Larval survival and growth in *Wallago attu* (Bloch and Schneider): Effects of light, photoperiod and feeding regimes. *Aquaculture*, 213(1-4), 151-161.
- GOI (2011). Census of India. Available online at http://www.censusindia.gov.in/2011-prov-results/data_files/india/FinalPPT2011chapter7.pdf%5Cnhttp://www.censusindia.go
- Gopal, B., & Chauhan, M. (2006). Biodiversity and its conservation in the Sundarban mangrove ecosystem. *Aquatic Sciences: Research Across Boundaries*, 68(3), 338-354.

- Gray, J. (2017, April 21). *Batagur kachuga* confiscation. Retrieved from Turtle Survival Alliance, www.turtlesurvival.org (accessed on 2 February 2018).
- Gunther, A.C. (1864). *The Reptiles of British India*. Piccadilly: Robert Hardwicke.
- Hamilton, F. (1822). *An Account of the Fishes Found in the River Ganges and Its Branches*. London: Archibald Constable and Company.
- Hanfee, F., (1999). *A WWF India Field Guide to Fresh Water Turtles and Tortoises of India*. TRAFFIC-India/WWF-India.
- Hinton, A.C.M., & Fry, T.B. (1923). BNHS's mammal survey of India, Burma and Ceylon. *Journal of the Bombay Natural History Society*, 29, 415-428.
- Hudson, R. (2016). River terrapins hatch throughout Asia. Retrieved from Turtle Survival Alliance, www.turtlesurvival.org (accessed on 2 February 2018)
- Hussain, S.A. (1999). Reproductive success, hatchling survival and growth of a managed population of Ganges gharial. *Biological Conservation*, 87, 261-268.
- Hussain, S.A. (2002). Conservation status of otters in the Tarai and Lower Himalayas of Uttar Pradesh, India. *IUCN Otter Specialists Group Bulletin A*, 19, 131-142.
- Hussain, S. A. (2009). Basking site and water depth selection by gharial *Gavialis gangeticus* Gmelin 1789 (Crocodylia, Reptilia) in National Chambal Sanctuary, India and its implication for river conservation. *Aquatic Conservation: Marine and Freshwater Ecosystems*, 19(2), 127-133.
- Hussain, S.A., Gupta, S.K., & de Silva, P.K. (2011). Biology and ecology of Asian small-clawed otter *Aonyx cinereus* (Illiger, 1815): a review. *IUCN Otter Specialist Group Bulletin*, 28(2), 63-75.
- ICAR-CIFRI (2017). Annual Report 2016-17. ICAR-Central Inland Fisheries Research Institute, Barrackpore, 232 pp.
- IIT (2012). Status of Higher Aquatic Vertebrates in the Ganga River. Ganga River Basin. Environment Management Plan by Indian Institutes of Technology.
- Inglis, C.M., Travers, W.L., O'Donel, H.V., & Shebbeare, E.O. (1919). A tentative list of the vertebrates of the Jalpaiguri District, Bengal. *Journal of the Bombay Natural History Society*, 26(3), 819-825.
- Islam, M.Z.U. & Rahmani, A.R. (2002). The threatened birds of India. *Buceros*, 7(1&2), 1-102.
- Islam, M.Z.U., & Rahmani, A.R. (2008). *Potential and Existing Ramsar Sites in India*. Oxford University Press.
- IUCN (International Union for Conservation of Nature) (2009). IUCN Red List of Threatened Species. Ver. 2009.1
- IUCN (International Union for Conservation of Nature) (2016). The IUCN Red List of Threatened Species. Version 2016-3.
- Iverson, J.B. (1992). A revised checklist with distribution maps of the turtles of the world (No. C/598.13 I9).
- IWPA (Indian Wildlife Protection Act) (1972). *The Indian Wildlife (Protection) Act, 1972*. Ministry of Environment and Forests, Government of India, New Delhi.
- Jain, S.K., Jeuland, M.A., Bharati, L., & Khan, Z.H. (2016). Surface water resources. In: Bharati, L., Sharma, B.R., & Smakhtin, V. (Eds.), *The Ganges River Basin: Status and Challenges in Water, Environment and Livelihoods* (pp. 8-23). London & New York: Routledge, Taylor and Francis Group.
- Jayaram, K.C., & Majumdar, N. (1976). On a collection of fish from the Mahanadi. *Records the Zoological Survey of India*, 69, 305-323.
- Jerdon, T.C. (1874). *The Mammals of India: A Natural History of All the Animals Known to Inhabit Continental India*. London: John Wheldon.

- Jha, B.R., & Rayamajhi, A. (2010). *Tor putitora*. The IUCN Red List of Threatened Species 2010.
- Jha, K.K., & McKinley, C.R. (2014). Demography and ecology of Indian sarus crane *Grus antigone antigone* in Uttar Pradesh, northern India. *Asian Journal of Conservation Biology*, 3(1), 8-18.
- Jha, S. (2006). Records of some rare birds from Farakka Barrage (West Bengal, India). *Indian Birds*, 2(4), 106.
- Jhingran, V.G. (1991). *Fish and Fisheries of India*, 3rd edn. New Delhi: Hindustan Publishing Corp.
- Jones, S. (1982). The present status of the Gangetic susu, *Platanista gangetica* (Roxburgh), with comments on the Indus susu, *Platanista minor* Owen. FAO Advisory Committee on Marine Resources Research Working Party on Marine Mammals. *FAO Fish. Ser. (5)* 4, 97-115.
- Joshi, R. (2013). Range extension of mugger crocodile *Crocodylus palustris* (Lesson, 1831) in Upper Ganges and tributaries, lesser Himalayan zone, north India. *Journal of Biological Earth Science*, 3(1), 100-109.
- Kar, S.K., & Bustard, H.R. (1986). Status of the saltwater crocodile (*Crocodylus porosus* Schneider) in the Bhitarkanika Wildlife Sanctuary, Orissa, India. *Journal of the Bombay Natural History Society*, 86(2), 141-150.
- Katdare, S., Srivathsa, A., Joshi, A., Panke, P., Pande, R., Khandal, D., & Everard, M. (2011). Gharial (*Gavialis gangeticus*) populations and human influences on habitat on the River Chambal, India. *Aquatic Conservation: Marine and Freshwater Ecosystems*, 21(4), 364-371.
- Kaur, J., Sundar, K.S.G. & Choudhury, B.C. (2002). *Sarus Crane Count-2001*. Dehradun: Wildlife Institute of India.
- Kelkar, N., Krishnaswamy, J., Choudhary, S., & Sutaria, D. (2010). Coexistence of fisheries with river dolphin conservation. *Conservation Biology*, 24(4), 1130-1140.
- Khan, M.S., Dimri, N.K., Nawab, A., Ilyas, O., & Gautam, P. (2014). Habitat use pattern and conservation status of smooth-coated otters *Lutrogale perspicillata* in the Upper Ganges Basin, India. *Animal Biodiversity and Conservation*, 37(1), 69-76.
- Kingsford, R.T., Dunn, H., Love, D., Nevill, J., Stein, J., & Tait, J. (2005). Protecting Australia's rivers, wetlands and estuaries of high conservation value. Canberra, Australia: Department of Environment and Heritage.
- Kruuk, H., Carss, D.N., Conroy, J.W.H., & Durbin, L. (1993). Otter (*Lutra lutra* L.) numbers and fish productivity in rivers in north-east Scotland. In: *Symp. Zool. Soc. Lond*, 65, 171-191.
- Kukkala, A.S., & Moilanen, A. (2013). Core concepts of spatial prioritisation in systematic conservation planning. *Biological Reviews*, 88(2), 443-464.
- Kumar, D. (2017a). River Ganges: Historical, cultural and socioeconomic attributes. *Aquatic Ecosystem Health & Management*, 20(1-2), 8-20.
- Kumar, A. (2017b). Effect of habitat characteristics on waterbird diversity along river Ganga in Allahabad, Uttar Pradesh. M.Sc. dissertation, Wildlife Institute of India, Dehradun.
- Lake, P.S., Bond, N., & Reich, P. (2007). Linking ecological theory with stream restoration. *Freshwater Biology*, 52(4), 597-615.
- Lakra, W.S., Sarkar, U.K., Gopalakrishnan, A., & Kathirvelpandian, A. (2010a). *Threatened Freshwater Fishes of India*. Lucknow: National Bureau of Fish Genetic Resources.
- Lakra, W.S., Sarkar, U.K., Kumar, R.S., Pandey, A., Dubey, V.K., & Gusain, O.P. (2010b). Fish diversity, habitat ecology and their conservation and management issues of a tropical river in Ganga basin, India. *The Environmentalist*, 30(4), 306-319.
- Latif, M., Ullah, M.Z., Minhas, I.B., & Latif, S. (2017). Morphometric study of *Puntius sophore* (Hamilton, 1822) with special reference to body length-weight from Chenab River, Punjab, Pakistan. *Journal of Entomology and Zoology Studies*, 5(6), 2032-2035.

- Lau, M., & Shi, H.T. (2000). Conservation and trade of terrestrial and freshwater turtles and tortoises in the People's Republic of China (pp. 30-38). Chelonian Research Monographs No. 2.
- Launer, A.E., & Murphy, D.D. (1994). Umbrella species and the conservation of habitat fragments: A case of a threatened butterfly and a vanishing grassland ecosystem. *Biological Conservation*, 69(2), 145-153.
- Lebreton, L.C., Van der Zwet, J., Damsteeg, J.W., Slat, B., Andrady, A., & Reisser, J. (2017). River plastic emissions to the world's oceans. *Nature Communications*, 8, 15611.
- MacArthur, R. (1955). Fluctuations of animal populations and a measure of community stability. *Ecology*, 36(3), 533-536.
- Macdonald, A. St. J. (1935). A fishing trip in Kumaon. *Journal of the Bombay Natural History Society*, 38, 598-600.
- Maheswaran, G., Deuti, K., & Khan, R.A. (2010). Rapid survey of Indian sarus crane (*Grus antigone*) in Uttar Pradesh. *Records of Zoological Survey of India*, 110 (Part 1), 71-81.
- Mall, R.K., Gupta, A., Singh, R., Singh, R.S., & Rathore, L.S. (2006). Water resources and climate change: An Indian perspective. *Current Science*, 1610-1626.
- Manakadan, R., & Pittie, A. (2001). Standardised common and scientific names of the birds of the Indian subcontinent. *Buceros*, 6(1), 1-37.
- Manna, R.K., Satpathy, B.B., Roshith, C.M., Naskar, M., Bhaumik, U., & Sharma, A.P. (2013). Spatio-temporal changes of hydro-chemical parameters in the estuarine part of the river Ganges under altered hydrological regime and its impact on biotic communities. *Aquatic Ecosystem Health & Management*, 16(4), 433-444.
- Martikainen, P., Kaila, L., & Haila, Y. (1998). Threatened beetles in white-backed woodpecker habitats. *Conservation Biology*, 12(2), 293-301.
- Martin, S. (2008). Global diversity of crocodiles (Crocodylia, Reptilia) in freshwater. *Hydrobiologia*, 595(1), 587-591.
- Mason, C.F., & Macdonald, S.M. (1987). The use of spraints for surveying otter *Lutra lutra* populations: An evaluation. *Biological Conservation*, 41(3), 167-177.
- May, R.M. (1973). Qualitative stability in model ecosystems. *Ecology*, 54(3), 638-641.
- McCann, C. (1940). A reptile and amphibian miscellany. *Journal of the Bombay Natural History Society*, 41, 742-764.
- McDonald, T., Gann, G.D., Jonson, J., & Dixon, K.W. (2016). *International Standards for the Practice of Ecological Restoration Including Principles and Key Concepts*. Washington, D.C.: Society for Ecological Restoration.
- McIntyre, P.B., Liermann, C.A.R., & Revenga, C. (2016). Linking freshwater fishery management to global food security and biodiversity conservation. *Proceedings of the National Academy of Sciences*, 113(45), 12880-12885.
- Mellin, C., Bradshaw, C.J.A., Fordham, D.A., & Caley, M.J. (2014). Strong but opposing α -diversity-stability relationships in coral reef fish communities. *Proceedings of the Royal Society of London B: Biological Sciences*, 281(1777), 20131993.
- Mishra, H., Kumar, V., & Kumar, A. (2016). Foraging guild status, diversity and population structure of waders of the River Ganges in District Rae Bareilly, Uttar Pradesh, India. *Journal of Entomology and Zoology Studies*, 4(6), 415-419.
- Misra, A.K. (2011). Impact of urbanization on the hydrology of Ganga Basin (India). *Water Resources Management*, 25(2), 705-719.

- Mohapatra, P. P., Palei, H. S., & Hussain, S. A. (2014). Occurrence of Asian small clawed otter *Aonyx cinereus* (Illiger, 1815) in Eastern India. *Current Science*, 107(3), 367-370.
- Moll, E.O. (1986). Survey of the freshwater turtles of India, Part I: The genus *Kachuga*. *Journal of the Bombay Natural History Society*, 83, 538-552.
- Moll, E.O. (1987). Survey of the freshwater turtles of India. Part II: The genus *Kachuga*. *Journal of the Bombay Natural History Society*, 84(1), 7-25.
- Moll, E.O., & Vijaya, J. (1986). Distributional records for some Indian turtles. *Journal of the Bombay Natural History Society*, 83(1), 57-62.
- Moll, E.O., Platt, K., Platt, S.G., Praschag, P., & Van Dijk, P.P. (2009). *Batagur baska* (Gray 1830): Northern river terrapin. *Chelonian Research Monographs* No. 5, 037-1.
- Montana, C.G., Choudhary, S.K., Dey, S., & Winemiller, K.O. (2011). Compositional trends of fisheries in the River Ganges, India. *Fisheries Management and Ecology*, 18(4), 282-296.
- Muller, J., & Henle, F.G.J. (1839). *Systematische Beschreibung der Plagiostome*. Berlin: Veit & Co.
- Nair, A.K. (2009). The status and distribution of major aquatic fauna in the National Chambal Gharial Sanctuary in Rajasthan with special reference to the Gangetic Dolphin *Platanista gangetica gangetica* (Cetartiodactyla: Platanistidae). *Journal of Threatened Taxa*, 1(3), 141-146.
- Nautiyal, P. (2010). Food chains of Ganga River ecosystems in the Himalayas. *Aquatic Ecosystem Health & Management*, 13(4), 362-373.
- Nawab, A. (2007). Ecology of otters in Corbett Tiger Reserve, Uttarakhand, India. Ph.D. thesis. Dehradun, Uttarakhand, India: Forest Research Institute (University). 174 pp.
- Nawab, A., & Hussain, S.A. (2012). Factors affecting the occurrence of smooth-coated otter in aquatic systems of the Upper Gangetic Plains, India. *Aquatic Conservation: Marine and Freshwater Ecosystems*, 22, 616-625.
- Nawab, A., Sinha, R.K., Thompson, P.M., & Sharma, S. (2016). Ecosystem services and conservation assessment of freshwater biodiversity. In: Bharati, L., Sharma, B.R., & Smakhtin, V. (Eds.). *The Ganges River Basin: Status and Challenges in Water, Environment and Livelihoods* (pp. 188-204). London & New York: Routledge, Taylor and Francis Group.
- Nel, J.L., Roux, D.J., Abell, R., Ashton, P.J., Cowling, R.M., Higgins, J.V., & Viers, J.H. (2009). Progress and challenges in freshwater conservation planning. *Aquatic Conservation: Marine and Freshwater Ecosystems*, 19(4), 474-485.
- Ng, H.H. (2010). *Notopterus Notopterus*. The IUCN Red List of Threatened Species. www.iucnredlist.org (accessed on 4 December 2017)
- NGRBA (National Ganga River Basin Authority) (2011). Environmental and Social Management Framework (ESMF). *Environmental and Social Analysis*, Vol. 1. New Delhi: Ministry of Environment, Forest & Climate Change, Government of India.
- Nilsson, C., Reidy, C.A., Dynesius, M., & Revenga, C. (2005). Fragmentation and flow regulation of the world's large river systems. *Science*, 308(5720), 405-408.
- Noss, R.F., O'Connell, M., & Murphy, D.D. (1997). *The Science of Conservation Planning: Habitat Conservation Under the Endangered Species Act*. Island Press.
- NRCD (National River Conservation Directorate) (2009). Status Paper on River Ganga. Ministry of Environment, Forest & Climate Change, Government of India.
- Odum, E.P. (1953). *Fundamentals of Ecology*. Philadelphia: Saunders.
- Pallas, P.S. (1769). *Spicilegia zoologica: quibus novae imprimis et obscurae animalium species iconibus, descriptionibus atque commentariis illustrantur*. Prostant Apud Joachimum Pauli.

- Palmer, M.A., Hart, D.D., Allan, J.D., & Bernhardt, E. (2004). Bridging engineering, ecological, and geomorphic science to enhance riverine restoration: Local and national efforts. In: Bizier, P., & DeBarry, P. (Eds.), *Proceedings of a National Symposium on Urban and Rural Stream Protection and Restoration* (pp. 29-37). EWRI World Water and Environmental Congress, Philadelphia, PA (June 2003). Reston, VA: American Society of Civil Engineers.
- Palmer, M.A., Menninger, H.L., & Bernhardt, E.S. (2010). River restoration, habitat heterogeneity and biodiversity: A failure of theory or practice? *Freshwater Biology*, 55(s1), 205-222.
- Palmer, M.A., Allan, J.D., Meyer, J.L., & Bernhardt, E.S. (2007). River restoration in the twenty-first century: Data and experiential knowledge to inform future efforts. *Restoration Ecology*, 15(3), 472-481.
- Pawar, S.S., & Choudhury, B.C. (2000). An inventory of chelonians from Mizoram, north-east India: New records and some observations on threats. *Hamadryad*, 25(2), 144-158.
- Payne, A.I., Sinha, R., Singh, H.R., & Huq, S. (2004). A review of the Ganges Basin, its fish and fisheries. In: Welcomme, R.L., & Petr, R.L. (Eds.) *Proceedings of the Second International Symposium on the Management of Large Rivers for Fisheries* Volume 1. Food and Agriculture Organization of the United Nations & Mekong River Commission. FAO Regional Office for Asia and the Pacific, Bangkok. RAP Publication 2004/16, 229-252.
- Perennou, C., Mundkur, T., Scott, D.A., Follestad, A., & Kvenild, L. (1994). *The Asian Waterfowl Census 1987-91: Distribution and Status of Asian Waterfowl*. Asian Wetland Bureau.
- Petr, T. (Ed.) (1999). *Fish and Fisheries at Higher Altitudes: Asia* (No. 385). FAO.
- Pocock, R.I. (1941). *The Fauna of British India, Including Ceylon and Burma*, Volume 2, Mammals. London: Taylor and Francis.
- Praschag, P., Sommer, R.S., McCarthy, C., & Gemel, R. (2008). Naming one of the world's rarest chelonians, the southern batagur. *Zootaxa*, 1758, 61-68.
- Pressey, R.L., Cabeza, M., Watts, M.E., Cowling, R.M., & Wilson, K.A. (2007). Conservation planning in a changing world. *Trends in Ecology & Evolution*, 22(11), 583-592.
- Raghavan, R., Das, S., Nameer, P.O., Bijukumar, A., & Dahanukar, N. (2016). Protected areas and imperilled endemic freshwater biodiversity in the Western Ghats hotspot. *Aquatic Conservation: Marine and Freshwater Ecosystems*, 26(S1), 78-90.
- Rahmani, A.R. (1981). Narora reservoir, UP: A potential bird sanctuary. *Journal of the Bombay Natural History Society*, 78(1), 88-92.
- Rao, R.J. (1993). Status and conservation of aquatic species diversity in certain water bodies of Madhya Pradesh. In: *Proceedings of Taal, the 12th World Lake Conference* (Vol. 2000, p. 2008).
- Rao, R.J. (2001). Biological resources of the Ganga river, India. *Hydrobiologia*, 458(1), 159-168.
- Rashid, S.M.A., & Khan, S.M.H. (2000). Trade and conservation status of freshwater turtles and tortoises in Bangladesh. In: *Asian Turtle Trade: Proceedings of a Workshop on Conservation and Trade of Freshwater Turtles and Tortoises in Asia*. Chelonian Research Monographs No. 2, pp. 77-85.
- Ray, P. (1998). *Ecological Imbalance of the Ganga River System: Its Impact on Aquaculture*. New Delhi: Daya Publishing House.
- Revenge, C., Campbell, I., Abell, R., De Villiers, P., & Bryer, M. (2005). Prospects for monitoring freshwater ecosystems towards the 2010 targets. *Philosophical Transactions of the Royal Society B: Biological Sciences*, 360(1454), 397-413.
- Rhodin, A.G.J. (2006). Turtles and humans in Florida and the world: A global perspective on diversity, threats, and economic development. In: Meylan, P.A. (Ed.), *Biology and Conservation of Florida Turtles* (pp. 19-27). Chelonian Research Monographs, Chelonian Research Foundation.

- Rodgers, W.A., & Panwar, H.S. (1988). Planning a Wildlife Protected Area Network for India: A Report Prepared for the Department of Environment, Forests & Wildlife, Government of India. Dehradun: Wildlife Institute of India.
- Rodgers, W.A., Panwar, H.S., & Mathur, V.B. (2000). *Wildlife Protected Area Network in India: A review* (Executive Summary). Dehradun: Wildlife Institute of India.
- Saha, M., Sarkar, A., & Bandyopadhyay, B. (2011). Study of saltwater crocodile (*Crocodylus porosus*) in mangrove forest at Sundarban in West Bengal for conservation and management. In: Gupta, V.K., & Verma, A.K. (Eds.), *Animal Diversity, Natural History and Conservation*. New Delhi: Daya Publishing House.
- Sanghi, R., & Kaushal, N. (2014). Introduction to our national river Ganga via cmaps. In: Sanghi, R. (Ed.), *Our National River Ganga* (pp. 3-44). Cham: Springer.
- Santiapillai, C., & de Silva, M. (2001). Status, distribution and conservation of crocodiles in Sri Lanka. *Biological Conservation*, 97(3), 305-318.
- Sarkar, U.K., Pathak, A.K., Sinha, R.K., Sivakumar, K., Pandian, A.K., Pandey, A., Dubey, V.K., & Lakra, W.S. (2012). Freshwater fish biodiversity in the River Ganga (India): Changing pattern, threats and conservation perspectives. *Reviews in Fish Biology and Fisheries*, 22(1), 251-272.
- Saunders, D.L., Meeuwig, J.J., & Vincent, A.C.J. (2002). Freshwater protected areas: Strategies for conservation. *Conservation Biology*, 16(1), 30-41.
- Sclater, W.L. (1891). *Catalogue of Mammalia in the Indian Museum, Kolkata*, Part 11. Printed by the Superintendent of Government Printing, Kolkata.
- Scott, J.M., Davis, F., Csuti, B., Noss, R., Butterfield, B., Groves, C., & Ulliman, J. (1993). Gap analysis: A geographic approach to protection of biological diversity. *Wildlife Monographs*, 3-41.
- Senthilkumar, K., Kannan, K., Sinha, R.K., Tanabe, S., & Giesy, J.P. (1999). Bioaccumulation profiles of polychlorinated biphenyl congeners and organochlorine pesticides in Ganges River dolphins. *Environmental Toxicology and Chemistry*, 18(7), 1511-1520.
- Sharma, A. (2008). Sighting of Indian skimmer *Rynchops albicollis* (Swinson) in the Purbasthali-Ganges Islets, Burdwan District, West Bengal. *Journal of the Bombay Natural History Society*, 105(1), 92-93.
- Sharma, P.D. (2005). *Ecology and Environment*. Meerut: Rastogi Publications.
- Sharma, R., & Basu, D. (2004). Recent reversals in the population trends in the population of gharial in the National Chambal Sanctuary in north India: Implications and a suggested strategy for the conservation of one of the world's most endangered crocodilians. In: *Crocodiles. Proceedings of the 17th Working Meeting of the Crocodile Specialist Group, IUCN-The World Conservation Union* (pp. 180-186).
- Sharma, R.C. (1998). *Fauna of India. Reptilia: Testudines and Crocodilians*. Vol 1. Zoological Survey of India, Kolkata.
- Sharma, R.C. (2003). Protection of an endangered fish *Tor tor* and *Tor putitora* population impacted by transportation network in the area of Tehri Dam Project, Garhwal Himalaya, India. In: *Proceedings of International Conference on Ecology & Transportation, Lake Placid, New York* (pp. 83-90).
- Sharma, R.K. (2004). Survey of Aquatic Animals in National Chambal Sanctuary, Morena, Madhya Pradesh. Report submitted to Superintendent of National Chambal Sanctuary, Madhya Pradesh, 30 pp.
- Sharma, R.K. (2005). Survey of Aquatic Animals in National Chambal Sanctuary, Morena, Madhya Pradesh. Report submitted to Superintendent of National Chambal Sanctuary, Madhya Pradesh, 24 pp.
- Sharma, R.K. (2006). Survey of Aquatic Animals in National Chambal Sanctuary, Morena, Madhya Pradesh. Report submitted to Superintendent of National Chambal Sanctuary, Madhya Pradesh, 26 pp.

- Sharma, R.K. (2007). Annual Faunal Survey in National Chambal Sanctuary, Morena, Madhya Pradesh. Report submitted to Superintendent of National Chambal Sanctuary, Madhya Pradesh, 15 pp.
- Sharma, R.K., & Singh, L.A.K. (2015). Status of mugger crocodile (*Crocodylus palustris*) in National Chambal Sanctuary after thirty years and its implications on conservation of gharial (*Gavialis gangeticus*). *Zoo's Print*, 30(5), 9-16.
- Sial, N., Haider, M.K., Shahzad, M.I., Abdullah, M., Malik, S., Tabasum, S., Mustafa, G., Azam, S.M. & Iqbal, H. (2016). Feeding behavior of freshwater turtle (*Pangshura smithii*) with respect to seasonal temperature variations. *Ciência e Técnica Vitivinícola*, 31, 254-223.
- Singh, D.S., Tangri, A.K., Kumar, D., Dubey, C.A., & Bali, R. (2017). Pattern of retreat and related morphological zones of Gangotri Glacier, Garhwal Himalaya, India. *Quaternary International*, 444, 172-181.
- Singh, I.B., Bajpai, V.N., Kumar, A., & Singh, M. (1990). Changes in the channel characteristics of Ganga River during late Pleistocene-Holocene. *Journal of the Geological Society of India*, 36(1), 67-73.
- Singh, L.A.K., & Sharma, R.K. (1985). Gangetic dolphin, *Platanista gangetica*: Observations on habits and distribution pattern in National Chambal Sanctuary. *Journal of the Bombay Natural History Society*, 82(3), 648-653.
- Singh, R.K., & Francis, R.C. (2017). Case study and comparison of data for solid waste management in Allahabad, Varanasi and Kanpur districts in U.P. (India). *International Journal of Mechanical Engineering and Technology*, 8(7), 1060-1067.
- Sinha, R., Mohanta, H., Jain, V., & Tandon, S.K. (2017). Geomorphic diversity as a river management tool and its application to the Ganga River, India. *River Research and Applications*, 33(7), 1156-1176.
- Sinha, R., Tandon, S.K., Gibling, M.R., Bhattacharjee, P.S., & Dasgupta, A.S. (2005). Late Quaternary geology and alluvial stratigraphy of the Ganga basin. *Himalayan Geology*, 26(1), 223-240.
- Sinha, R.K. (1999). The Ganges River Dolphin: A Tool for Baseline Assessment of Biological Diversity in River Ganges (No. 1/99). India. Final Technical Report No. 1/99. Patna University, Patna, India.
- Sinha, R.K., Behera, S.K., & Choudhary, B.C. (2010). *The Conservation Action Plan for the Ganges River Dolphin 2010-2020*. New Delhi: National Ganga River Basin Authority (NGBRA), Ministry of Environment, Forest and Climate Change, Government of India.
- Sinha, R.K., & Kannan, K. (2014). Ganges river dolphin: An overview of biology, ecology, and conservation status in India. *Ambio*, 43(8), 1029-1046.
- Sinha, R.K., Kedia, D.K., & Kumari, A. (2014). Overview of higher vertebrates in the Ganges-Brahmaputra-Meghna river basin: Their status, threats and conservation. In: Sinha, R.K., & Ahmed, B. (Eds.), *Rivers for Life: Proceedings of the International Symposium on River Biodiversity: Ganges-Brahmaputra-Meghna River System, Ecosystems for Life, Bangladesh-India Initiative*. IUCN, 105 pp.
- Sinha, R.K., & Sharma, G. (2003). Current status of the Ganges river dolphin, *Platanista gangetica* in the rivers Kosi and Son, Bihar, India. *Journal of the Bombay Natural History Society*, 100(1), 27-37.
- Sinha, R.K., Smith, B.D., Sharma, G., Prasad, G., Choudhury, B.C., Sapkota, K., & Behera, S.K. (2000). Status and distribution of the Ganges susu (*Platanista gangetica*) in the Ganges River system of India and Nepal. *Occasional Paper of the IUCN Species Survival Commission*, 23, 54-61.
- Smith, A.M. (1931). *The Fauna of British India, Including Ceylon and Burma: Reptilia and Amphibia*, Vol. I. London: Taylor and Francis.
- Srivastava, S.M., Singh, S.P., Kumari, A., & Pandey, A.K. (2012). Fecundity of threatened bronze featherback, *Notopterus notopterus* (Pallas) from river Gomti, Lucknow (India). *National Journal of Life Science*, 9(2), 193-200.
- Stevens, E.R. (1905). A morning's mahseer fishing in the Ganges. *Indian Forester*, 31, 406.
- Strayer, D.L. (2010). Alien species in fresh waters: Ecological effects, interactions with other stressors, and prospects for the future. *Freshwater Biology*, 55, 152-174.

- Strayer, D.L., & Dudgeon, D. (2010). Freshwater biodiversity conservation: Recent progress and future challenges. *Journal of the North American Benthological Society*, 29(1), 344-358.
- Stuart, B.L., & Thorbiarnarson, J. (2003). Biological prioritization of Asian countries for turtle conservation. *Chelonian Conservation and Biology*, 4(3), 642-647.
- Sundar, K.S.G. (2004). Observations on breeding Indian skimmers *Rynchops albicollis* in the National Chambal Sanctuary, Uttar Pradesh, India. *Forktail*, 20, 89-90.
- Sundar, K.S.G. (2008). Uttar Pradesh: An unlikely Shangri-La. *ICF Bugle*, 34(2), 6.
- Sundar, K.S.G., Kaur, J., & Choudhury, B.C. (2000). Distribution, demography and conservation status of the Indian sarus crane (*Grus antigone antigone*) in India. *Journal of the Bombay Natural History Society*, 97(3), 319-339.
- Talwar, K.P., & Jhingran, A.G. (1991). *Inland Fishes of India and Adjacent Countries*, Vol. 2. New Delhi: Oxford & IBH.
- Talwar, P.K. (1991). Pisces. In: Director, Zoological Survey of India (Ed.), *Faunal Resources of Ganga*, Part 1, (pp. 59-145). Calcutta: Zoological Survey of India.
- Tandon, S.K., & Sinha, R. (2018). The Ganga River: A summary view of a large river system of the Indian sub-continent. In: Singh, D.S. (Ed.), *The Indian Rivers: Scientific and Socio-economics Aspects* (pp. 61-73). Singapore: Springer.
- Theobald, W. (1876). *Descriptive Catalogue of the Reptiles of British India*. Calcutta: Thacker, Spink and Co.
- Tripathi, S., Gopesh, A., Joshi, K.D., Dwivedi, A.C., & Mayank, P. (2013). Studies on feeding behaviour of *Labeo bata* (Hamilton, 1822) from the lower stretch of the Yamuna river, Uttar Pradesh. *Journal of the Kalash Science*, 2, 49-52.
- Trivedi, R.C. (2010). Water quality of the Ganga River: An overview. *Aquatic Ecosystem Health & Management*, 13(4), 347-351.
- Valdiya, K.S. (2016). Indo-Gangetic plains: Evolution and later developments. In: *The Making of India* (pp. 723-745). Cham: Springer.
- van der Ploeg, J., Cauillan-Cureg, M., van Weerd, M., & Persoon, G. (2011). 'Why must we protect crocodiles?'- Explaining the value of the Phillipine crocodile to rural communities. *Journal of Integrative Environmental Sciences*, 8(4), 287-298.
- Vass, K.K., Mondal, S.K., Samanta, S., Suresh, V.R., & Katiha, P.K. (2010). The environment and fishery status of the river Ganges. *Aquatic Ecosystem Health and Management Society*, 13(4), 385-394.
- Vasudeva, V. (2017). Waterbird abundance and diversity along the stretch of Ganga River impacted by the Narora barrage. M.Sc. dissertation, GGS Indraprastha University, New Delhi.
- Vijaya, J. (1983). Range extension for the spotted pond turtle *Geoclemys hamiltonii*. *Hamadryad*, 8(2).
- Vorosmarty, C.J., McIntyre, P.B., Gessner, M.O., Dudgeon, D., Prusevich, A., Green, P., & Davies, P.M. (2010). Global threats to human water security and river biodiversity. *Nature*, 467(7315), 555.
- Wadia, D.N. (1961). *Geology of India*. London: MC Muller & Company.
- Wetlands International (Eds.) (2012). Waterbird Population Estimates. Summary report. Wageningen, The Netherlands: Wetlands International.
- Whitaker, R. (1979). The crocodilians of Corbett National Park. *Indian Journal of Forestry*, 2(1), 38-40.
- Whitaker, R. (1997). Turtle rearing in village ponds. In: *Proceedings: Conservation, Restoration, and Management of Tortoises and Turtles: An International Conference* (pp. 106-108). NY Turtle and Tortoise Society.
- Whitaker, R. (2007). The gharial: Going extinct again. *Iguana*, 14(1), 25-32.

- Whitaker, R., & Daniel, J.C. (1980). The status of Indian crocodilians. *Journal of the Bombay Natural History Society*, 75, 1238-1245.
- Wink, A. (2002). From the Mediterranean to the Indian Ocean: Medieval history in geographic perspective. *Comparative Studies in Society and History*, 44(3), 416-445.
- Wright, L. (1982). *The IUCN Amphibia-Reptilia Red Data Book*, Vol. 1. IUCN.
- WWF (2016). Living Planet Report 2016 Risk and Resilience in a New Era. Gland, Switzerland: WWF International.
- Yeung, L.W., Yamashita, N., Taniyasu, S., Lam, P.K., Sinha, R.K., Borole, D.V., & Kannan, K. (2009). A survey of perfluorinated compounds in surface water and biota including dolphins from the Ganges River and in other waterbodies in India. *Chemosphere*, 76(1), 55-62.
- Zöckler, C., Balachandran, S., Bunting, G.C., Fanck, M., Kashiwagi, M., Lappo, E.G., Maheswaran, G., Sharma, A., Syroechkovski, E.E., & Webb, K. (2005). The Indian Sunderbans: An important wintering site for Siberian waders. *Wader Study Group Bulletin*, 108, 42-46.
- ZSI (1991). *Faunal Resources of Ganga*, Part I. Calcutta: Zoological Survey of India.
- ZSI (1992). *Fauna of West Bengal*, Part 2. State Faunal Series 3. Calcutta: Zoological Survey of India.

ANNEXURE I

Mammalian species of the Ganga River

Family	Common Name/s	Scientific Name	IUCN status	IWPA status	Previous studies	Present study	Ganga specific biogeographic province/s (2B, 7A, 7B, 8B)
Catecea	Gangetic river dolphin	<i>Platanista gangetica</i> (Roxburgh, 1801)	EN	Schedule I	b, i, j, k, n, o	+	7A, 7B
	Irrawaddy dolphin	<i>Orcaella brevirostris</i> (Owen in Gray, 1866)	EN	Schedule I	I		8B
Carnivora	Smooth-coated otter	<i>Lutrogale perspicillata</i> (L. Geoffroy Saint-Hilaire, 1826)	VU	Schedule II	c, d, f, g, h, m, n, o	+	2B, 7A, 7B
	Eurasian otter	<i>Lutra lutra</i> (Linnaeus, 1758)	NT	Schedule II	d, e, m		2B
	Asian small clawed otter	<i>Aonyx cinereus</i> (Illiger, 1815)	VU	Schedule I	a, c, m		2B, 8B

^aJerdon (1874); ^bAnderson (1878); ^cBlanford (1881); ^dAtkinson (1882); ^eSclater (1891); ^fHinton and Fry (1923); ^gPocock (1941), ^hAtkinson (1974); ⁱJones (1982); ^jBehera (1995); ^kSinha (1999); ^lSmith et al. (2006); ^mHussain (2002); ⁿChoudhary et al. (2006); ^oSinha et al. (2010); ^pNawab & Hussain (2012)

REFERENCES

- Anderson, J. (1878). *Anatomical and Zoological Researches:Comprising an Account of Zoological Results of the Two Expeditions to Western Yunnan in 1868 and 1875; and a Monograph of the Two Cetacean Genera, Platanista and Orcella*. London: Bernard Quaritch.
- Atkinson, E.T. (1882). *The Himalayan Districts of the North Western Province of India*. Allahabad. Reprinted in 1996 as the Himalayan Gazetteer, Vol. 2, Part I. Dehradun:Natraj Publishers.
- Atkinson, E.T. (1974). *Fauna of the Himalayas, Containing Species of Kumaon, Garhwal, Nepal and Tibet*. New Delhi: Cosmo Publications.
- Behera, S.K. (1995). Studies on population dynamics, habitat utilization and conservation aspects of Gangetic dolphin (*Platanista gangetica*) in a stretch of Ganga River from Rishekech to Kanpur. Ph.D. thesis, School of Studies in Zoology, Jiwaji University Gwalior, 198 pp.
- Blanford, W.T. (1881). *Fauna of British India, Including Ceylon and Burma: Mammalia*. London: Taylor and Francis.
- Choudhary, S.K., Smith, B.D., Dey, S., Dey, S., & Prakash, S. (2006). Conservation and biomonitoring in the Vikramshila Gangetic Dolphin Sanctuary, Bihar, India. *Oryx*, 40(2), 189-197.
- Hinton, A.C.M., & Fry, T.B. (1923). BNHS's mammal survey of India, Burma and Ceylon. *Journal of the Bombay Natural History Society*, 29, 415-428.

- Hussain, S.A. (2002). Conservation status of otters in the Tarai and Lower Himalayas of Uttar Pradesh, India. *IUCN Otter Specialists Group Bulletin A*, 19, 131-142.
- Jerdon, T.C. (1874). *The Mammals of India: A Natural History of All the Animals Known to Inhabit Continental India*. London: John Wheldon.
- Jones, S. (1982). The present status of the Gangetic susu, *Platanista gangetica* (Roxburgh), with comments on the Indus susu, *Platanista minor* (Owen). FAO Advisory Committee on Marine Resources Research Working Party on Marine Mammals. FAO Fish. Ser. (5) 4, 97-115.
- Nawab, A., & Hussain, S.A. (2012). Factors affecting the occurrence of smooth-coated otter in aquatic systems of the Upper Gangetic Plains, India. *Aquatic Conservation: Marine and Freshwater Ecosystems*, 22, 616-625.
- Pocock, R.I. (1941). *The Fauna of British India, Including Ceylon and Burma, Volume 2, Mammals*. London: Taylor and Francis.
- Sclater, W.L. (1891). *Catalogue of Mammalia in the Indian Museum, Kolkata*, Part 11. Printed by the Superintendent of Government Printing, Kolkata.
- Sinha, R.K. (1999). The Ganges River Dolphin: A Tool for Baseline Assessment of Biological Diversity in River Ganges (No. 1/99). India. Final Technical Report No. 1/99. Patna University, Patna, India.
- Sinha, R.K., Behera, S.K., & Choudhary, B.C. (2010). *The Conservation Action Plan for the Ganges River Dolphin 2010-2020*. New Delhi: National Ganga River Basin Authority (NGBRA), Ministry of Environment, Forest and Climate Change, Government of India.
- Smith, B. D., Braulik, G., Strindberg, S., Ahmed, B., & Mansur, R. (2006). Abundance of Irrawaddy dolphins (*Orcaella brevirostris*) and Ganges river dolphins (*Platanista gangetica gangetica*) estimated using concurrent counts made by independent teams in waterways of the Sundarbans mangrove forest in Bangladesh. *Marine Mammal Science*, 22(3), 527-547.

ANNEXURE II



Waterbird and water associated bird species of the Ganga River

Family	Common Name/s	Scientific Name (Synonyms)	IUCN Status
Podicipedidae	Little grebe	<i>Tachybaptus ruficollis</i> (Pallas, 1764)	LC
	Great crested grebe	<i>Podiceps cristatus</i> (Linnaeus, 1758)	LC
Phalacrocoracidae	Little cormorant	<i>Microcarbo niger</i> (Vieillot, 1817)	LC
	Indian cormorant	<i>Phalacrocorax fuscicollis</i> (Stephens, 1826)	LC
	Great cormorant	<i>Phalacrocorax carbo</i> (Linnaeus, 1758)	LC
Anhingidae	Oriental darter	<i>Anhinga melanogaster</i> (Pennant, 1769)	NT
Ardeidae	Little egret	<i>Egretta garzetta</i> (Linnaeus, 1766)	LC
	Grey heron	<i>Ardea cinerea</i> (Linnaeus, 1758)	LC
	Purple heron	<i>Ardea purpurea</i> (Linnaeus, 1766)	LC
	Great white egret	<i>Ardea alba</i> (Linnaeus, 1758)	LC
	Intermediate egret	<i>Ardea intermedia</i> (Wagler, 1829)	LC
	Cattle egret	<i>Bubulcus ibis</i> (Linnaeus, 1758)	LC
	Indian pond-heron	<i>Ardeola grayii</i> (Sykes, 1832)	LC
	Green-backed heron	<i>Butorides striata</i> (Linnaeus, 1758)	LC
	Black-crowned Night-heron	<i>Nycticorax nycticorax</i> (Linnaeus, 1758)	LC
	Yellow bittern	<i>Ixobrychus sinensis</i> (Gmelin, 1789)	LC
	Black bittern	<i>Ixobrychus flavicollis</i> (Latham, 1790)	LC
Ciconiidae	Painted stork	<i>Mycteria leucocephala</i> (Pennant, 1769)	NT
	Asian openbill-stork	<i>Anastomus oscitans</i> (Boddaert, 1783)	LC

IWPA status	Previous Study	Present Study	Ganga specific biogeographic province/s (2B, 7A, 7B, 8B)
Schedule IV	d, f, h, j	+	2B, 7A, 7B
Schedule IV	f, j	+	2B, 7A, 7B
Schedule IV	d, h, j	+	2B, 7A, 7B
Schedule IV	d, h, j	+	2B, 7A, 7B
Schedule IV	a	+	2B, 7A, 7B
Schedule IV	d, g	+	2B, 7A, 7B
Schedule IV	d, h, j	+	2B, 7A, 7B
Schedule IV	a	+	2B, 7A, 7B
Schedule IV	a, d, h, j	+	2B, 7A, 7B
Schedule IV	d, h, j	+	2B, 7A, 7B
Schedule IV	d, h	+	2B, 7A, 7B
Schedule IV	d, h, j	+	2B, 7A, 7B
Schedule IV	a, h, j	+	2B, 7A, 7B
Schedule IV	e	+	2B, 7A, 7B
Schedule IV	d	+	2B, 7A, 7B
Schedule IV	a, j		2B, 7A, 7B
Schedule IV		+	2B, 7A, 7B
Schedule IV	d, g	+	2B, 7A, 7B
Schedule IV	d, h	+	2B, 7A, 7B

Family	Common Name/s	Scientific Name (Synonyms)	IUCN Status
	Asian wolly-necked stork	<i>Ciconia episcopus</i> (Boddaert, 1783)	VU
	White stork	<i>Ciconia ciconia</i> (Linnaeus, 1758)	LC
	Black-necked stork	<i>Ephippiorhynchus asiaticus</i> (Latham, 1790)	NT
	Lesser Adjutant	<i>Leptoptilos javanicus</i> (Horsfield, 1821)	VU
	Greater adjutant	<i>Leptoptilos dubius</i> (Gmelin, 1789)	EN
Threskiornithidae	Glossy ibis	<i>Plegadis falcinellus</i> (Linnaeus, 1766)	LC
	Black-headed ibis	<i>Threskiornis melanocephalus</i> (Latham, 1790)	NT
	Red-naped ibis	<i>Pseudibis papillosa</i> (Temminck, 1824)	LC
	Eurasian spoonbill	<i>Platalea leucorodia</i> (Linnaeus, 1758)	LC
Anatidae	Fulvous whistling-duck	<i>Dendrocygna bicolor</i> (Vieillot, 1816)	LC
	Lesser whistling-duck	<i>Dendrocygna javanica</i> (Horsfield, 1821)	LC
	Lesser white-fronted goose	<i>Anser erythropus</i> (Linnaeus, 1758)	VU
	Greylag goose	<i>Anser anser</i> (Linnaeus, 1758)	LC
	Bar-headed goose	<i>Anser indicus</i> (Latham, 1790)	LC
	Ruddy shelduck	<i>Tadorna ferruginea</i> (Pallas, 1764)	LC
	Common shelduck	<i>Tadorna tadorna</i> (Linnaeus, 1758)	LC
	African Comb Duck	<i>Sarkidiornis melanotos</i> (Pennant, 1769)	LC
	Cotton teal	<i>Nettapus coromandelianus</i> (Gmelin, 1789)	LC
	Gadwall	<i>Mareca strepera</i> (Linnaeus, 1758)	LC

IWPA status	Previous Study	Present Study	Ganga specific biogeographic province/s (2B, 7A, 7B, 8B)
Schedule IV	d	+	2B, 7A, 7B
Schedule I	a		2B, 7A
Schedule IV	a, g	+	2B, 7A, 7B
Schedule IV	g, h	+	2B, 7A, 7B
Schedule IV	a, g	+	7A, 7B
Schedule IV	j	+	2B, 7A, 7B
Schedule IV	a, g, h, j	+	2B, 7A, 7B
Schedule IV	a, d, g, h	+	2B, 7A, 7B
Schedule IV	a, d	+	2B, 7A, 7B
Schedule I	f, h		7B
Schedule IV	d, f, h, j	+	2B, 7A, 7B
Schedule IV	g		7A
Schedule IV	a, f, g	+	2B, 7A, 7B
Schedule IV	a, d, f, g	+	2B, 7A, 7B
Schedule IV	a, d, f, g, j	+	2B, 7A, 7B
Schedule IV	a	+	2B, 7A, 7B
Schedule IV	d, g, j	+	2B, 7A, 7B
Schedule IV	d, f, g, j	+	2B, 7A, 7B
Schedule IV	a, d, f, g	+	2B, 7A, 7B

Family	Common Name/s	Scientific Name (Synonyms)	IUCN Status
	Falcated duck	<i>Mareca falcata</i> (Georgi, 1775)	NT
	Eurasian wigeon	<i>Mareca penelope</i> (Linnaeus, 1758)	LC
	Mallard	<i>Anas platyrhynchos</i> (Linnaeus, 1758)	LC
	Indian spot-billed duck	<i>Anas poecilorhyncha</i> (J.R.Forester, 1781)	LC
	Northern shoveler	<i>Spatula clypeata</i> (Linnaeus, 1758)	LC
	Northern pintail	<i>Anas acuta</i> (Linnaeus, 1758)	LC
	Garganey	<i>Spatula querquedula</i> (Linnaeus, 1758)	LC
	Eurasian teal	<i>Anas crecca</i> (Linnaeus, 1758)	LC
	Marbled teal	<i>Marmaronetta angustirostris</i> (Menetries, 1832)	VU
	Red-crested pochard	<i>Netta rufina</i> (Pallas, 1773)	LC
	Common pochard	<i>Aythya ferina</i> (Linnaeus, 1758)	VU
	Ferruginous duck	<i>Aythya nyroca</i> (Guldenstadt, 1770)	NT
	Baer's pochard	<i>Aythya baeri</i> (Radde, 1863)	CR
	Tufted duck	<i>Aythya fuligula</i> (Linnaeus, 1758)	LC
	Goosander	<i>Mergus merganser</i> (Linnaeus, 1758)	LC
	Tufted duck	<i>Aythya fuligula</i>	LC
Accipitridae	Black-shouldered kite	<i>Elanus caeruleus</i> (Desfontaines, 1789)	LC
	Brahminy kite	<i>Haliastur indus</i> (Boddaert, 1783)	LC
	Pallas's fish-eagle	<i>Haliaeetus leucoryphus</i> (Pallas, 1771)	EN
	Lesser fish-eagle	<i>Ichthyophaga humilis</i> (S.Muller & Schlegel, 1841)	NT
	Grey-headed fish-eagle	<i>Ichthyophaga ichthyaetus</i> (Horsfield, 1821)	NT
	Egyptian vulture	<i>Neophron percnopterus</i> (Linnaeus, 1758)	EN

IWPA status	Previous Study	Present Study	Ganga specific biogeographic province/s (2B, 7A, 7B, 8B)
Schedule IV	g		2B, 7B
Schedule IV	f, j	+	2B, 7A, 7B
Schedule IV	d	+	2B, 7A, 7B
Schedule IV	d, g, j	+	2B, 7A, 7B
Schedule IV	a, d, f, j	+	2B, 7A, 7B
Schedule IV	d, f, h, j	+	2B, 7A, 7B
Schedule IV	d, j	+	2B, 7A, 7B
Schedule IV	f, j	+	2B, 7A, 7B
Schedule IV	g		2B
Schedule IV	a, f, g, h, j	+	2B, 7A, 7B
Schedule IV	d, f, g, j	+	2B, 7A, 7B
Schedule IV	f, g, h, j	+	2B, 7A, 7B
Schedule IV	g		7B
Schedule IV	f, j	+	2B, 7A, 7B
Schedule IV	b	+	2B, 7A
	e		2B, 7A, 7B
Schedule IV	d, j		2B, 7A, 7B
Schedule IV	e	+	2B, 7A, 7B
Schedule IV	g	+	2B, 7A, 7B
Schedule IV	g		2B
Schedule IV	g	+	7A, 7B
Schedule IV	g		2B, 7A, 7B

Family	Common Name/s	Scientific Name (Synonyms)	IUCN Status
	Western marsh-harrier	<i>Circus aeruginosus</i> (Linnaeus, 1758)	LC
	Greater spotted eagle	<i>Clanga clanga</i> (Pallas, 1811)	VU
Pandionidae	Osprey	<i>Pandion haliaetus</i> (Linnaeus, 1758)	LC
Falconidae	Red-headed falcon	<i>Falco chicquera</i> (Daudin, 1800)	NT
	Peregrine falcon	<i>Falco peregrinus</i> (Tunstall, 1771)	LC
Gruidae	Sarus crane	<i>Antigone antigone</i> (Linnaeus, 1758),	VU
	Demoiselle crane	<i>Anthropoides virgo</i> (Linnaeus, 1758),	LC
	Common crane	<i>Grus grus</i> (Linnaeus, 1758)	LC
Rallidae	White-breasted waterhen	<i>Amauornis phoenicurus</i> (Pennant, 1769)	LC
	Baillon's crake	<i>Zapornia pusilla</i> (Pallas, 1776)	LC
	Purple moorhen	<i>Porphyrio porphyrio</i> (Linnaeus, 1758)	LC
	Common moorhen	<i>Gallinula chloropus</i> (Linnaeus, 1758)	LC
	Common coot	<i>Fulica atra</i> (Linnaeus, 1758)	LC
Jacanidae	Pheasant-tailed jacana	<i>Hydrophasianus chirurgus</i> (Scopoli, 1786)	LC
	Bronze-winged jacana	<i>Metopidius indicus</i> (Latham, 1790)	LC
Charadriidae	Little ringed plover	<i>Charadrius dubius</i> (Scopoli, 1786)	LC
	Kentish plover	<i>Charadrius alexandrinus</i> (Linnaeus, 1758)	LC
	River lapwing	<i>Vanellus duvaucelii</i> (Lesson, 1826)	NT
	Grey-headed lapwing	<i>Vanellus cinereus</i> (Blyth, 1842)	LC
	Red-wattled lapwing	<i>Vanellus indicus</i> (Boddaert, 1783)	LC
	White-tailed lapwing	<i>Vanellus leucurus</i> (Lichtenstein, 1823)	LC

IWPA status	Previous Study	Present Study	Ganga specific biogeographic province/s (2B, 7A, 7B, 8B)
Schedule IV	d, f, j	+	2B, 7A, 7B
Schedule IV	g	+	2B, 7A, 7B
Schedule I	j	+	2B, 7A, 7B
Schedule I	e		2B, 7A, 7B
Schedule I	e	+	2B, 7A, 7B
Schedule IV	d, f, g	+	2B, 7A
Schedule IV	a, d	+	2B, 7A
Schedule IV	a	+	2B, 7A, 7B
Schedule IV	d, h, j	+	2B, 7A, 7B
Schedule IV	j		2B, 7A
Schedule IV	d, g, h, j	+	Not in Ganga state
Schedule IV	d, j	+	2B, 7A, 7B
Schedule IV	d, f, h, j	+	2B, 7A, 7B
Schedule IV	d, h, j	+	2B, 7A, 7B
Schedule IV	d, h, j	+	2B, 7A, 7B
Schedule IV	d, j	+	2B, 7A, 7B
Schedule IV	e	+	2B, 7A, 7B
Schedule IV	d, g	+	2B, 7A, 7B
Schedule IV	f, h, j	+	2B, 7A, 7B
Schedule IV	d, h, j	+	2B, 7A, 7B
Schedule IV	e		2B, 7A, 7B

Family	Common Name/s	Scientific Name (Synonyms)	IUCN Status
Scolopacidae	Common snipe	<i>Gallinago gallinago</i> (Linnaeus, 1758)	LC
	Black-tailed godwit	<i>Limosa limosa</i> (Linnaeus, 1758)	NT
	Eurasian curlew	<i>Numenius arquata</i> (Linnaeus, 1758)	NT
	Spotted redshank	<i>Tringa erythropus</i> (Pallas, 1764)	LC
	Common redshank	<i>Tringa totanus</i> (Linnaeus, 1758)	LC
	Marsh sandpiper	<i>Tringa stagnatilis</i> (Bechstein, 1803)	LC
	Common greenshank	<i>Tringa nebularia</i> (Gunnerus, 1767)	LC
	Green sandpiper	<i>Tringa ochropus</i> (Linnaeus, 1758)	LC
	Wood sandpiper	<i>Tringa glareola</i> (Linnaeus, 1758)	LC
	Common sandpiper	<i>Actitis hypoleucos</i> (Linnaeus, 1758)	LC
	Little stint	<i>Calidris minuta</i> (Leisler, 1812)	LC
	Curlew sandpiper	<i>Calidris ferruginea</i> (Pontoppidan, 1763)	NT
	Ruff	<i>Calidris pugnax</i> (Linnaeus, 1758)	LC
Recurvirostridae	Black-winged stilt	<i>Himantopus himantopus</i> (Linnaeus, 1758)	LC
	Pied avocet	<i>Recurvirostra avosetta</i> (Linnaeus, 1758)	LC
Burhinidae	Great thick-knee	<i>Esacus recurvirostris</i> (Cuvier, 1829)	NT
	Indian thick knee	<i>Burhinus indicus</i> (Salvadori, 1865)	LC
Glareolidae	Little pratincole	<i>Glareola lactea</i> (Temminck, 1820)	LC
Laridae	Pallas's gull	<i>Larus ichthyæus</i> (Pallas, 1773)	LC
	Brown-headed gull	<i>Larus brunnicephalus</i> (Jerdon, 1840)	LC
	Black-headed gull	<i>Larus ridibundus</i> (Linnaeus, 1766)	LC
	Gull-billed tern	<i>Gelochelidon nilotica</i> (Gmelin, 1789)	LC
	River tern	<i>Sterna aurantia</i> (J.E.Gray, 1831)	NT
	Common tern	<i>Sterna hirundo</i> (Linnaeus, 1758)	LC
	Little tern	<i>Sternula albifrons</i> (Pallas, 1764)	LC
	Black-bellied tern	<i>Sterna acuticauda</i> (J.E.Gray, 1831)	EN

IWPA status	Previous Study	Present Study	Ganga specific biogeographic province/s (2B, 7A, 7B, 8B)
Schedule IV	h, j		2B, 7A, 7B
Schedule IV	g, i	+	7B
Schedule IV	g	+	2B, 7A, 7B
Schedule IV	e	+	2B, 7A, 7B
Schedule IV	f, j	+	2B, 7A, 7B
Schedule IV	j	+	2B, 7A, 7B
Schedule IV	d, j	+	2B, 7A, 7B
Schedule IV	d, f, j	+	2B, 7A, 7B
Schedule IV	f, h, j	+	2B, 7A, 7B
Schedule IV	d, f, j	+	2B, 7A, 7B
Schedule IV	h, j	+	2B, 7A, 7B
Schedule IV	d		2B, 7A, 7B
Schedule IV	c	+	2B, 7A, 7B
Schedule IV	d, j	+	2B, 7A, 7B
Schedule IV	d	+	2B, 7A, 7B
Schedule IV	d, g	+	2B, 7A
NL	e		2B, 7A, 7B
NL	d, g, j	+	2B, 7A, 7B
Schedule IV	d	+	2B, 7A, 7B
Schedule IV	a, h	+	2B, 7A, 7B
Schedule IV	h	+	2B, 7A, 7B
NL	d		2B, 7A, 7B
NL	a, d, g	+	2B, 7A, 7B
NL	e	+	7B
NL	g	+	2B, 7A, 7B
NL	d, g	+	2B, 7A, 7B

Family	Common Name/s	Scientific Name (Synonyms)	IUCN Status
Laridae	Indian skimmer	<i>Rynchops albicollis</i> (Swainson, 1838)	VU
Alcedinidae	Small blue kingfisher	<i>Alcedo atthis</i> (Linnaeus, 1758)	LC
	Stork-billed kingfisher	<i>Pelargopsis capensis</i> (Linnaeus, 1766)	LC
	White-breasted kingfisher	<i>Halcyon smymensis</i> (Linnaeus, 1758)	LC
	Crested kingfisher	<i>Megaceryle lugubris</i> (Temminck, 1834)	LC
	Pied kingfisher	<i>Ceryle rudis</i> (Linnaeus, 1758)	LC
Hirundinidae	Plain martin	<i>Riparia paludicola</i> (Vieillot, 1871)	LC
	Barn swallow	<i>Hirundo rustica</i> (Linnaeus, 1758)	LC
	Wire-tailed swallow	<i>Hirundo smithii</i> (Leach, 1818)	LC
	Red rumped swallow	<i>Cecropis daurica</i> (Linnaeus, 1771)	LC
	Streak-throated swallow	<i>Petrochelidon fluvicola</i> (Blyth, 1855)	LC
Motacillidae	White wagtail	<i>Motacilla alba</i> (Linnaeus, 1758)	LC
	Large pied wagtail	<i>Motacilla maderaspatensis</i> (Gmelin, 1789)	LC
	Citrine wagtail	<i>Motacilla citreola</i> (Pallas, 1776)	LC
	Western yellow wagtail	<i>Motacilla flava</i> (Linnaeus, 1758)	LC
	Grey wagtail	<i>Motacilla cinerea</i> (Tunstall, 1771)	LC
Muscicapidae	Black redstart	<i>Phoenicurus ochruros</i> (Gmelin, 1774)	LC
	Plumbeous water redstart	<i>Phoenicurus fuliginosus</i> (Vigors, 1831)	LC
	White-capped water redstart	<i>Phoenicurus leucocephalus</i> (Vigors, 1831)	LC
Pelecanidae	Great white pelican	<i>Pelecanus onocrotalus</i> (Linnaeus, 1758)	LC
Alaudidae	Oriental skylark	<i>Alauda gulgula</i> (Franklin, 1831)	LC
Sturnidae	Asian pied starling	<i>Gracupica contra</i> (Linnaeus, 1758)	LC

^aBilgrami (1991); ^bdel Hoyo et al. (1996); ^cBarter, M.A. (2002); ^dKhan et al. (2013); ^eDey et al. (2014); ^fKhan et al. (2016); ^gRahmani et al. (2016); ^hSatya Prakash et al. (2016); ⁱDixit et al. (2016); ^jChowdhury (2017)

IWPA status	Previous Study	Present Study	Ganga specific biogeographic province/s (2B, 7A, 7B, 8B)
NL	d, g	+	2B, 7A, 7B
Schedule IV	h, j	+	2B, 7A, 7B
Schedule IV	j		2B, 7A, 7B
Schedule IV	d, h, j	+	2B, 7A, 7B
Schedule IV	e	+	2B
Schedule IV	d, h, j	+	2B, 7A, 7B
NL	e	+	7A, 7B
NL	h	+	2B, 7A, 7B
NL	d		2B, 7A, 7B
NL	e		2B, 7A, 7B
NL	e		2B, 7A, 7B
NL	e	+	2B, 7A, 7B
NL	h, j	+	2B, 7A, 7B
NL	h, j	+	2B, 7A, 7B
NL	h, j	+	2B, 7A, 7B
NL	e	+	2B, 7A, 7B
Schedule IV	e	+	2B, 7A, 7B
Schedule IV	e, i	+	7B, 7A, 2B
Schedule IV	e	+	7B, 7A, 2B
NL	e		2B, 7A
NL	e		2B, 7A, 7B
Schedule-IV	e		2B, 7A, 7B

REFERENCES



- Barter, M.A. (2002). Shorebirds of the Yellow Sea: Importance, threats and conservation status. *Wetlands International Global Series 9, International Wader Studies 12*, Canberra, Australia. pp 118.
- Bilgrami, K.S. (1991). Biological profile of the Ganga: Zooplankton, fish, birds, and other minor fauna. In Krishna Murti, C.R., Bilgrami, K.S., Das, T.M., & Mathur, R.P. (Eds.). *The Ganga: A Scientific Study*, (pp. 83-94). New Delhi: Ganga Project Directorate, Ministry of Environment & Forests, Government of India.
- Chowdhury, G & Patra, A (2017). Behavioral ecology of Blue tailed Bee eaters (*Merops philippinus*) in Hooghly and Burdwan District of West Bengal, India. *International Research Journal of Biological Science.*, 6 (11), 31-35.
- del Hoyo, J., Elliott, A., & Sargatal, J. (1996). *Handbook of the Birds of the World, 3, Hoatzin to Auks*. Barcelona: Lynx Edicions.
- Dey, S., Dey, S., & Kelkar, S.K.C.N. (2014). An annotated bird checklist of the Vikramshila Gangetic Dolphin Sanctuary, Bhagalpur, Bihar, India, with an assessment of threats to bird conservation. *Forktail*, 30(30), 34-40.
- Dixit, S., Joshi, V., & Barve, S. (2016). Bird diversity of the Amrutganga Valley, Kedarnath, Uttarakhand, India with an emphasis on the elevational distribution of species. *Check List*, 12(2), 1874.
- Khan, M. S., Aftab, A., Syed, Z., Nawab, A., Ilyas, O., & Khan, A. (2013). Composition and conservation status of avian species at Hastinapur Wildlife Sanctuary, Uttar Pradesh, India. *Journal of Threatened Taxa*, 5(12), 4714-4721.
- Khan, T.N., A. Sinha & P. Hazra (2016). Population trends and community composition of migratory waterbirds in three emerging wetlands of global significance in southwestern Bengal, India. *Journal of Threatened Taxa* 8(3) 8541-8555.
- Prakash, S., Dwivedi, M. D., Raziuddin, M., Paul, K., Marthak, K., Singh, M., & Samanta, I. (2016). Report on Asian Waterbird Census (AWC), Department of Forest, Environment & Climate Change, Government of Jharkhand, 14. pp 11.
- Rahmani, A.R., Islam, M.Z., & Kasambe, R.M., (2016). *Important Bird and Biodiversity Areas in India: Priority Sites for Conservation (Revised and updated)*. Bombay Natural History Society, Indian Bird Conservation Network, Royal Society for the Protection of Birds and BirdLife International (U.K.). (Pp 1992 + xii).

ANNEXURE III

Crocodilian species of the Ganga River

Family	Common Name/s	Scientific Name	IUCN status	IWPA status	Previous studies	Present study	Ganga specific biogeographic province/s (2B, 7A, 7B, 8B)
Crocodylidae	Mugger	<i>Crocodylus palustris</i> (Lesson, 1831)	VU	Schedule I	a, h, g	+	7A, 7B
	Estuarine crocodile	<i>Crocodylus porosus</i> (Schneider, 1801)	LC	Schedule I	c, d, f		8B
Gavialidae	Gharial	<i>Gavialis gangeticus</i> (Gmelin in Linnaeus, 1789)	CR	Schedule I	a, b, e, i	+	7A, 7B

^aBoulenger (1890); ^bSmith (1931); ^cDaniel (1970); ^dBustard & Choudhury (1981); ^eWright (1982); ^fSaha et al (2011); ^gJoshi (2013);

^hSharma & Singh (2015); ⁱNawab et al. (2016)

REFERENCES

- Boulenger, G.A. (1890). *The Fauna of British India Including Ceylon and Burma: Published Under the Authority of the Secretary of State for India in Council. Reptilia and Batrachia*. London: Taylor and Francis.
- Bustard, H.R., & Choudhury, B.C. (1981). Conservation future of saltwater crocodile (*Crocodylus porosus* Schneider) in India. *Journal of the Bombay Natural History Society*, 77(2), 201-214.
- Daniel, J.C. (1970). A review of the present status and position of endangered species of Indian reptiles. In: *Proceedings of the IUCN 11th Technical Meeting, New Delhi* (pp.75-76). IUCN Publications Series No. 18.
- Joshi, R. (2013). Range extension of mugger crocodile *Crocodylus palustris* (Lesson, 1831) in Upper Ganges and tributaries, lesser Himalayan zone, north India. *Journal of Biological Earth Science*, 3(1), 100-109.
- Nawab, A., Sinha, R.K., Thompson, P.M., & Sharma, S. (2016). Ecosystem services and conservation assessment of freshwater biodiversity. In: Bharati, L., Sharma, B.R., & Smakhtin, V. (Eds.), *The Ganges River Basin: Status and Challenges in Water, Environment and Livelihoods* (pp. 188-204). London & New York: Routledge, Taylor and Francis Group.
- Saha, M., Sarkar, A., & Bandyopadhyay, B. (2011). Study of saltwater crocodile (*Crocodylus porosus*) in mangrove forest at Sundarban in West Bengal for conservation and management. In: Gupta, V.K., & Verma, A.K. (Eds.), *Animal Diversity, Natural History and Conservation*. New Delhi: Daya Publishing House.
- Sharma, R.K., & Singh, L.A.K. (2015). Status of mugger crocodile (*Crocodylus palustris*) in National Chambal Sanctuary after thirty years and its implications on conservation of gharial (*Gavialis gangeticus*). *Zoo's Print*, 30(5), 9-16.
- Smith, A.M. (1931). *The Fauna of British India, Including Ceylon and Burma: Reptilia and Amphibia*, Vol. I. London: Taylor and Francis.
- Wright, L. (1982). *The IUCN Amphibia-Reptilia Red Data Book*, Vol. 1. IUCN.

ANNEXURE IV

Turtle species of the Ganga River

Family	Common Name/s	Scientific Name	IUCN status	IWPA status	Previous studies	Present study	Ganga specific biogeographic province/s (2B, 7A, 7B, 8B)
Geoemydidae	Northern river terrapin	<i>Batagur baska</i> (Gray, 1830)	CR	Schedule I	m, o, s		8B
	Red-crowned roofed turtle	<i>Batagur kachuga</i> (Gray, 1831)	CR	Schedule I	b, s, t		7A, 7B
	Three-striped roofed turtle	<i>Batagur dhongoka</i> (Gray, 1832)	EN	NL	a	+	7A, 7B
	Spotted pond turtle	<i>Geoclemys hamiltonii</i> (Gray, 1830)	VU	Schedule I	b, e	+	7A, 7B
	Crowned river turtle	<i>Hardella thurjii</i> (Gray, 1831)	VU	NL	d,	+	7A, 7B
	Indian black turtle	<i>Melanochelys trijuga</i> (Schweigger, 1812)	NT	NL	g, h,	+	7A, 7B
	Brown roofed turtle	<i>Pangshura smithii</i> (Gray, 1863)	NT	NL	d, h, i	+	2B, 7A, 7B
	Indian roofed turtle	<i>Pangshura tecta</i> (Gray, 1830)	LC	Schedule I	k, l	+	7A, 7B
	Indian tent turtle	<i>Pangshura tentoria</i> (Gray, 1834)	LC	NL	i	+	7A, 7B
Trionychidae	Indian narrow-headed softshell turtle	<i>Chitra indica</i> (Gray, 1830)	EN	Schedule IV	a, c, e, h, n	+	7A, 7B
	Indian softshell turtle	<i>Nilssonina gangetica</i> (Cuvier, 1825)	VU	Schedule I	a, c, p	+	7A, 7B
	Indian peacock softshell turtle	<i>Nilssonina hurum</i> (Gray, 1830)	VU	Schedule I	c, e, g, p, r	+	7A, 7B
	Indian flapshell turtle	<i>Lissemys punctata</i> (Lacépède, 1788)	LC	Schedule I	q, j	+	7A, 7B

^aGunther (1864); ^bAnderson (1871); ^cAnnandale (1912); ^dChaudhuri (1912); ^eSmith (1931); ^fDaniel & Shull (1964); ^gMoll & Vijaya (1986); ^hDas (1991); ⁱRao (1993); ^jWhitaker (1997); ^kChoudhury et al. (2000); ^lPraschag et al. (2007); ^mDas & Bhupathy (2009); ⁿDas & Singh (2009); ^oMoll et al. (2009); ^pDas et al. (2010); ^qBhupathy et al. (2014); ^rSinha, et al. (2014); ^sIUCN (2016); ^tGray (2017)

REFERENCES

- Anderson, J. (1871). On some Indian reptiles. In *Proceedings of the Zoological Society of London*, 11, 149-211.
- Annandale, N. (1912). The Indian mud turtles (Trionychidae). *Records of the Indian Museum*, 7(2), 151-180.

- Bhupathy, S., Webb, R.G., & Praschag, P. (2014). *Lissemys punctata* (Bonnaterre 1789) - Indian Flapshell Turtle. In: Rhodin, A.G.J., Pritchard, P.C.H., van Dijk, P.P., Saumure, R.A., Buhlmann, K.A., Iverson, J.B., & Mittermeier, R.A. (Eds.). *Conservation Biology of Freshwater Turtles and Tortoises: A Compilation Project of the IUCN/SSC Tortoise and Freshwater Turtle Specialist Group*. Chelonian Research Monographs No. 5, 076.1-12.
- Chaudhuri, B.L. (1912). Aquatic tortoises of the middle Ganga and Brahmaputra. *Records of the Indian Museum*, 7, 212-214.
- Choudhury, B.C., Bhupathy, S., & Hanfee, F. (2000). Status information on the tortoises and freshwater turtles of India. In: van Dijk, P.P., Stuart, B.L., and Rhodin, A.G.J. (Eds.). *Asian Turtle Trade: Proceedings of a Workshop on Conservation and Trade of Freshwater Turtles and Tortoises in Asia*. Chelonian Research Monographs No. 2, 86-94.
- Daniel, J.C., & Shull, E.M. (1964). A list of the reptiles and amphibians of Surat Dangs, South Gujarat. *Journal of the Bombay Natural History Society*, 60, 737-743.
- Das, I. (1991). *Colour Guide to the Turtles and Tortoises of the Indian Subcontinent*. R & A Publishing.
- Das, I., & Bhupathy, S. (2009). *Batagur baska* (Gray 1830) Northern River Terrapin. In: Rhodin, A.G.J., Pritchard, P.C.H., van Dijk, P.P., Saumure, R.A., Buhlmann, K.A., Iverson, J.B., & Mittermeier, R.A. (Eds.). *Conservation Biology of Freshwater Turtles and Tortoises: A Compilation Project of the IUCN/SSC Tortoise and Freshwater Turtle Specialist Group*. Chelonian Research Monographs No. 5, 043.1-043.6.
- Das, I., & Singh, S. (2009). *Chitra indica* (Gray 1830): Narrow-headed softshell turtle. In: Rhodin, A.G.J., Pritchard, P.C.H., van Dijk, P.P., Saumure, R.A., Buhlmann, K.A., Iverson, J.B., & Mittermeier, R.A. (Eds.). *Conservation Biology of Freshwater Turtles and Tortoises: A Compilation Project of the IUCN/SSC Tortoise and Freshwater Turtle Specialist Group*. Chelonian Research Monographs No. 5, 027.1-027.7.
- Das, I., Basu, D., & Singh, S. (2010). *Nilssonina hurum* (Gray 1830): Indian peacock softshell turtle. In: Rhodin, A.G.J., Pritchard, P.C.H., van Dijk, P.P., Saumure, R.A., Buhlmann, K.A., Iverson, J.B., & Mittermeier, R.A. (Eds.). *Conservation Biology of Freshwater Turtles and Tortoises: A Compilation Project of the IUCN/SSC Tortoise and Freshwater Turtle Specialist Group*. Chelonian Research Monographs No. 5, 048.1-048.6.
- Gray, J. (2017, April 21). *Batagur kachuga* confiscation. Retrieved from Turtle Survival Alliance, www.turtlesurvival.org (accessed on 2 February 2018).
- Gunther, A.C. (1864). *The Reptiles of British India*. Piccadilly: Robert Hardwicke.
- IUCN (International Union for Conservation of Nature) (2016). The IUCN Red List of Threatened Species. Version 2016-3.
- Moll, E.O., & Vijaya, J. (1986). Distributional records for some Indian turtles. *Journal of the Bombay Natural History Society*, 83(1), 57-62.
- Moll, E.O., Platt, K., Platt, S.G., Praschag, P., & Van Dijk, P.P. (2009). *Batagur baska* (Gray 1830): Northern river terrapin. *Chelonian Research Monographs* No. 5, 037-1.
- Praschag, P., Hundsdoerfer, A.K., & Fritz, U. (2007). Phylogeny and taxonomy of endangered South and South-east Asian freshwater turtles elucidated by mtDNA sequence variation (Testudines: Geoemydidae: Batagur, Callagur, Hardella, Kachuga, Pangshura). *Zoologica Scripta* 36(5):429-442.
- Rao, R.J. (1993). Status and conservation of aquatic species diversity in certain water bodies of Madhya Pradesh. In: *Proceedings of Taal, the 12th World Lake Conference* (Vol. 2000, p. 2008).
- Sinha, R.K., Kedia, D.K., & Kumari, A. (2014). Overview of higher vertebrates in the Ganges-Brahmaputra-Meghna river basin: Their status, threats and conservation. In: Sinha, R.K., & Ahmed, B. (Eds.), *Rivers for Life: Proceedings of the International Symposium on River Biodiversity: Ganges-Brahmaputra-Meghna River System, Ecosystems for Life, a Bangladesh-India Initiative*. IUCN, 105 pp.
- Smith, A.M. (1931). *The Fauna of British India, Including Ceylon and Burma: Reptilia and Amphibia*, Vol. I. London: Taylor and Francis.
- Whitaker, R. (1997). Turtle rearing in village ponds. In: *Proceedings: Conservation, Restoration, and Management of Tortoises and Turtles: An International Conference* (pp. 106-108). NY Turtle and Tortoise Society.

ANNEXURE V

Water Snakes species of the Ganga River

Family	Common Name	Scientific Name
Natricidea	Checkered keelback	<i>Xenochrophis piscator</i> (Schneider, 1799)
	Dark bellied keelback water snake	<i>Xenochrophis cerasogaster</i> (Cantor, 1839)
Homalopsidae	Common smooth water snake	<i>Enhydris enhydris</i> (Schneider, 1799)
	Seibold's water snake	<i>Ferania seiboldii</i> (Schlegel, 1837)
	Crab eating water snake	<i>Fordonia leucobalia</i> (Schlegel, 1837)
	Glossy marsh snake	<i>Gerarda prevostiana</i> (Eydoux & Gervais, 1837)
	Dog faced water snake	<i>Cerberus rynchops</i> (Schneider, 1799)
Acrochordidae	Wart snake	<i>Acrochordus granulatus</i> (Schneider, 1799)

^aSmith (1943); ^bWhitaker et al. (2004); ^cDas (2017)

REFERENCES

Das, I., & Das, A. (2017). *A naturalists guide to the reptiles of India Bangladesh, Bhutan, Nepal, Pakistan and Sri Lanka*. John Beaufoy publishing. 176 pp.

Smith, M.A. (1943). *The Fauna of British India, Ceylon and Burma, Including the Whole of the Indo-Chinese Sub-Region. Reptilia and Amphibia*. Vol. III (Serpentes). London: Taylor and Francis.

Whitaker, R., & Captain, A., (2004). *Snakes of India*. Draco books. pp 500.

IUCN Status	IPWA Status	Previous Studies	Present Studies	Ganga specific biogeographic province/s (2B, 7A, 7B, 8B)
NE	Schedule II	a, b, c	+	7A, 7B, 8B
NE	Schedule IV	a, b, c	+	7A, 7B
LC	Schedule IV	a, b		7B
NE	Schedule IV	a, b, c	+	7A, 7B
LC	Schedule IV	a, b, c		8B
LC	Schedule IV	a, b, c		8B
LC	Schedule IV	a, b, c		8B
LC	NL	a, b, c		8B

ANNEXURE VI

Amphibian species of the Ganga River

Family	Common Name/s	Scientific Name
Ranidae	Mountain cascade frog	<i>Amolops monticola</i> (Anderson, 1871)
	Dehradun stream frog	<i>Amolops chakrataensis</i> (Ray, 1992)
	Assam cascade frog	<i>Amolops himalayanus</i> (Boulenger, 1988)
	Cascade frog	<i>Amolops formosus</i> (Gunther, 1876)
	Jaunsar stream frog	<i>Amolops jaunsari</i> (Ray, 1992)
	Marbled cascade frog	<i>Amolops marmoratus</i> (Blyth, 1855)
	Tytler's pond frog	<i>Hylarana tytleri</i> (Theobald, 1868)
Rhacophoridae	Dudhwa tree frog	<i>Chiromantis dudhwaensis</i> (Ray, 1992)
	Himalayan tree frog	<i>Polypedates maculatus</i> ssp. <i>maculatus</i> (Gray, 1830)
	Bengal whipping frog	<i>Polypedates taeniatus</i> (Boulenger, 1906)
Bufonidae	Himalayan toad	<i>Duttaphrynus himalayanus</i> (Günther, 1864)
	Black-spectacled toad	<i>Duttaphrynus melanostictus</i> (Schneider, 1799)
	Marbled toad	<i>Duttaphrynus stomaticus</i> (Lütken, 1864)
Dicroglossidae	Indian skipping frog	<i>Euphlyctis cyanophlyctis</i> (Schneider, 1799)
	Indian green frog	<i>Euphlyctis hexadactylus</i> (Lesson, 1834)
	Hill cricket frog	<i>Fejervarya syhadrensis</i> (Annandale, 1919)
	Terai cricket frog	<i>Fejervarya teraiensis</i> (Dubois, 1984)
	Jerdon's bullfrog	<i>Hoplobatrachus crassus</i> (Jerdon, 1854 "1853")
	Indian bullfrog	<i>Hoplobatrachus tigerinus</i> (Daudin, 1802)
	Annandale's paa frog	<i>Nanorana annandalii</i> (Boulenger, 1920)
	Blanford's paa frog	<i>Nanorana blanfordii</i> (Boulenger, 1882)
	Sikkim paa frog	<i>Nanorana liebigii</i> (Günther, 1860)
	Nepal paa frog	<i>Nanorana minica</i> (Dubois, 1975)
	Himalaya paa frog	<i>Nanorana vicina</i> (Stoliczka, 1872)
Microhylidae	Ant frog	<i>Microhyla ornata</i> (Duméril & Bibron, 1841)

^aAnderson (1871); ^bSclater (1892); ^cTilak & Hussain (1977); ^dSarkar (1984); ^eTilak & Roy (1985); ^fSarkar et al. (1992); ^gRay & Tilak (1995); ^hRay (1995); ⁱRay (1999); ^jHussain (2003); ^kMehta & Uniyal (2007); ^lHegde (2009); ^mSingh et al. (2009); ⁿBahuguna & Bhutia (2010); ^oHegde & Roy (2011); ^pDas et al. (2012); ^qKanaujia & Kumar (2013); ^rBahuguna et al. (2014); ^sHussain (2015); ^tBahuguna et al. (2017); ^uBilgrami (1991); ^vPratihara & Deuti (2011); ^wSubba et al. (2017). ^xPatel & Das (2016): Pers. Obs.

IUCN status	IWPA status	Previous studies	Present Study	Ganga specific biogeographic province/s (2B, 7A, 7B, 8B)
LC	Schedule IV	x		2B
DD	Schedule IV	i, n	+	2B
NE	Schedule IV	w		7B
LC	Schedule IV	x	+	2B
DD	Schedule IV	i, n	+	2B
LC	Schedule IV	h, n		7B
LC	NL	v	+	7B
DD	NL	p, q	+	7A
LC	NL	a, d, f, g, h, k, n, o, q, s		7A, 7B
LC	NL	p, q	+	7B
LC	NL	h, i, n, q	+	2B
LC	NL	a, d, f, g, h, j, n, o, p, q, r, s	+	2B,7A,7B,8B
LC	NL	b, d, f, g, h, i, n, o, p, q, s, u	+	7A, 7B
LC	NL	a, d, f, g, h, j, k, n, o, p, q, s, t	+	2B,7A, 7B, 8B
LC	Schedule IV	d, f, u		7B
LC	Schedule IV	n	+	7A, 7B
LC	Schedule IV	l, o, p, q	+	7A,7B
LC	Schedule IV	d, f, g, i, k, n, q, s		7B
LC	Schedule IV	a, d, f, g, h, k, m, n, o, p, q, s, t	+	2B,7A,7B,8B
NT	Schedule IV	n		7B
LC	Schedule IV	h, n		7B
LC	Schedule IV	n		7B
VU	Schedule IV	e, h, n	+	2B
LC	Schedule IV	q	+	2B
LC	NL	d, f, g, h, i, j, n, o, p, q, s	+	2B,7A,7B,8B

REFERENCES

- Anderson, J. (1871). A list of the reptilian accession to the Indian Museum, Calcutta, from 1865 to 1870, with a description of some new species. *Journal of the Asiatic Society of Bengal*, 40(2), 12-39.
- Bahuguna, A., & Butia, P. (2010). Amphibia. In *Fauna of Uttarakhand. State Fauna Series*, 18(1), 505-532.
- Bahuguna, A., & Kumar, V. (2017). Diversity and morphometry of *Fejervarya* (anura: dicroglossidae) in Sonanadi Wildlife Sanctuary, Nainital, Uttarakhand. (23), 391-403.
- Bahuguna, A., Sharma, I., Sharma, G., Nesemann, H., & Parmod, K. (2017). Study of some selected aquatic and semi-aquatic fauna of Baanganga wetlands, Haridwar (Uttarakhand, India). *International Journal of Advanced Research*, 5(3), 2228-2243.
- Das, A., Basu, D., Converse, L., & Suresh, C. C. (2012). Herpetofauna of Katarniaghat Wildlife Sanctuary, Uttar Pradesh, India. *Journal of Threatened Taxa*, 4(5), 2553-2568.
- Hegde, V. D., & Roy, S. (2011). Amphibian fauna of Katarniaghat Wildlife Sanctuary, Uttar Pradesh, India. *Frog leg*, 15.
- Hegde, V. D., Roy, S., & Lal, B. (2009). First record of Terai Cricket Frog *Fejervarya teraiensis* (Dubois 1984) from Uttar Pradesh. *Record of Zoological Survey of India*, 109(Part I), 109-110.
- Husain, A. (2003). Amphibia. In *Fauna of Asan Wetland. Wetland Ecosystem Series. Zoological Survey of India*, (5), 27-28.
- Husain, A. (2015). Amphibians of Doon Valley (Dehra Dun, Uttarakhand) with their Systematics, Distribution, Ecology, Conservation Status and Threats. In *Aquatic Ecosystem: Biodiversity, Ecology and Conservation. Springer*, 217-229.
- Kanaujia, A., & Kumar, A. (2013). Amphibians of Uttar Pradesh and Their Ecological Importance. In *National Conference on Water and Biodiversity on occasion of International Day for Biological diversity*, 148-157.
- Mehta, H. S., & Uniyal, D. P. (2007). In Mehta, H. S., & Gupta, S. K. (Eds). *Faunal diversity western Doon Shiwaliks. Zoological Survey of India, Kolkata*, 41-59.
- Ohler, A., & Mallick, P. K. (2002). *Rana (Hylarana) sensu Dubois (1992) in India and the identity of Hylorana tyleri* Theobald, 1868. *Hamadryad Madras*, 27, 62-70.
- Pratihari, S., & Deuti, K. (2011). Distribution and threats of amphibian species in west Midnapore District, West Bengal, India. *frog leg*, (17), 2.
- Ray, P. (1995) Amphibia. In *Fauna of Western Himalaya, Part 1, Uttar Pradesh. Himalayan ecosystem series. Zoological Survey of India, Kolkata*, 151-157.
- Ray, P. (1999). Systematic studies on the amphibian fauna of the district Dehradun, Uttar Pradesh, India. *Zoological Survey of India, Kolkata*.
- Ray, P., & Tilak, R. (1995). Amphibia. In *Fauna of Rajaji National Park, fauna of conservation areas, Western Himalaya (Uttar Pradesh). Zoological Survey of India, Kolkata*, (5), 54-75.
- Sarkar, A. K. (1984). Taxonomic and ecological studies on the amphibians of Calcutta and its environs. In *Records of Zoological Survey of India*, 81(3-4), 215-236.
- Sarkar, A. K., & Biswas, M. L. (1992). Ray S. Amphibia. In *State Fauna of West Bengal. Zoological Survey of India, State Fauna Series*, 3(2), 67-100.
- Slater, W. L. (1892). On some specimens of frogs in the Indian Museum, Calcutta, with descriptions of several new species. In *Proceedings of the Zoological Society of London*, (1892), 341-348.

- Singh, S. K., Srivastava, S. P., Tandon, P., & Azad, B. S. (2009). Faunal diversity during rainy season in reclaimed sodic land of Uttar Pradesh, India. *Journal of Environment Biology*, 30(4):551-6
- Subba, B., Aravind, N. A., & Ravikanth, G. (2017). Amphibians of the Sikkim Himalaya, India: an annotated checklist. *Check List*, 13(1), 1-14.
- Tilak, R., & Husain, A. (1977). Extension of the range of distribution of a microhylid frog, *Uperodon systoma* (Schneider). *Journal of Bombay Natural History Society*, 73(1), 407.
- Tilak, R., & Roy, P. (1985). Description of a new species of the sub-genus *Rana* (Paa) Dubois from Chakrata Hills, district Dehra Dun, Uttar Pradesh, India (Ranidae, Anura). *Zoologischer Anzeiger*, 215(3-4), 231-239.

ANNEXURE VII



Fish species of the Ganga River

Family	Common Name/s	Scientific Name
Adrianichthyidae	Rice fish	<i>Oryzias melastigma</i> (McClelland, 1839)
Ambassidae	Elongate glass perchlet Highfin glassy perchlet Indian glassy fish Himalayan glassy perchlet	<i>Chanda nama</i> (Hamilton, 1822) <i>Parambassis lala</i> (Hamilton, 1822) <i>Parambassis ranga</i> (Hamilton, 1822) <i>Pseudambassis baculis</i> (Hamilton, 1822)
Amblycipitidae	Biting catfish, Indian torrent catfish	<i>Amblyceps mangois</i> (Hamilton, 1822)
Anabantidae	Gangetic koi Climbing perch Indian mottled eel	<i>Anabas cobojus</i> (Hamilton, 1822) <i>Anabas testudineus</i> (Bloch, 1792) <i>Anguilla bengalensis</i> (Gray, 1831)
Aplocheilidae	Blue panchax	<i>Aplocheilus panchax</i> (Hamilton, 1822)
Ariidae	Threadfin sea catfish Gagora catfish	<i>Arius arius</i> (Hamilton, 1822) <i>Arius gagora</i> (Hamilton, 1822)
Badidae	Badis	<i>Badis badis</i> (Hamilton, 1822)
Bagridae	Batasio, bojori Menoda catfish Day's mystus Gangetic mystus Long-whiskered catfish Tengaramystus Striped dwarf catfish Rita Long-whiskered catfish Giant river-catfish	<i>Batasio batasio</i> (Hamilton, 1822) <i>Hemibagrus menoda</i> (Hamilton, 1822) <i>Mystus bleekeri</i> (Day, 1877) <i>Mystus cavasius</i> (Hamilton, 1822) <i>Mystus gulio</i> (Hamilton, 1822) <i>Mystus tengara</i> (Hamilton, 1822) <i>Mystus vittatus</i> (Bloch, 1794) <i>Rita rita</i> (Hamilton, 1822) <i>Sperata aor</i> (Hamilton, 1822) <i>Sperata seenghala</i> (Sykes, 1839)
Balitoridae	- Mottled loach, Sand loach, Striped loach Doon loach Moorangi Korika, Stone loach Loach Loach Loach Puinya Savon khorka Dari	<i>Aborichthys elongatus</i> (Hora, 1921) <i>Acanthocobitis botia</i> (Hamilton, 1822) <i>Nemacheilus doonensis</i> (Tilak & Husain, 1977) <i>Nemacheilus mooreh</i> (Sykes, 1839) <i>Schisturacorica</i> (Hamilton, 1822) <i>Schistura devdevi</i> (Hora, 1935) <i>Schistura gangetica</i> (Menon, 1987) <i>Schistura multifasciata</i> (Day, 1878) <i>Schistura rupecula</i> (McClelland, 1838) <i>Schistura savona</i> (Hamilton, 1822) <i>Schisturas caturigina</i> (McClelland, 1839)
Belonidae	Banded needlefish Freshwater garfish	<i>Strongylura leiura</i> (Bleeker, 1850) <i>Xenentodon cancila</i> (Hamilton, 1822)
Carcharhinidae	Gangetic shark	<i>Glyphis gangeticus</i> (Müller & Henle, 1839)*

IUCN Status	Previous Studies	Present Study	Ganga specific biogeographic province/s (2B, 7A, 7B, 8B)
LC	d,e		8B
LC	a,b,c,d,e,f,h,i,k	+	7A, 7B, 8B
NT	a,h	+	7A, 7B, 8B
LC	b,c,d,e,f,h,i,j,k	+	7A, 7B, 8B
LC	a,b,c,d,e,i	+	2B , 7A, 7B
LC	a,b,d,f,i	+	2B, 7A
DD	a,b,c,d,e,f,h,i,j,k		7A, 7B, 8B
DD	a,b,c,d,e,f,i	+	7A, 7B, 8B
NT	a,c,d,e,h,i,j,k	+	7A, 7B, 8B
LC	a,b,d,e,f,h,j,k		7B, 8B
LC	a,d		7B, 8B
LC	a,d,e,h	+	7B, 8B
LC	a,b,c,d,e,f,h,j,k	+	7A, 7B, 8B
LC	a,e,h,i		7A, 7B, 8B
LC	a,b,c,d,i		7B
LC	b,c,d,e,f,g,h,i,j	+	7A, 7B, 8B
LC	a,b,c,d,e,f,g,h,i,j,k	+	7A, 7B, 8B
LC	d,e,g,h,j,k		7B, 8B
LC	a,b,d,f,h,i,k	+	7A, 7B, 8B
LC	b,c,d,e,f,g,h,i,j,k	+	7A, 7B, 8B
LC	a,b,c,d,e,f,g,h,i,j,k	+	7A, 7B, 8B
LC	a,b,c,d,e,f,g,h,i,j,k	+	7A, 7B, 8B
LC	b,c,d,e,f,g,h,i,j,k	+	7A, 7B, 8B
LC	h		7B, 8B
LC	a,c,d,e,f,h,i	+	2B, 7A, 7B, 8B
DD	d		2B
LC	i		2B
LC	a,d,f,i		2B
NT	i		2B
NL	d,h		2B
LC	f		2B
LC	d,f,i		2B
LC	a,d,f		2B
LC	f,h		2B
NL	d,h		8B
LC	a,b,c,d,e,f,h,i,j,k	+	2B, 7A, 7B, 8B
CR	d		8B

Family	Common Name/s	Scientific Name
Chacidae	Squarehead catfish	<i>Chaca chaca</i> (Hamilton, 1822)
Channidae	Borna snakehead	<i>Channa amphibeus</i> (McClelland, 1845)
	Barca snakehead	<i>Channa barca</i> (Hamilton, 1822)
	Dwarf snakehead	<i>Channa gachua</i> (Hamilton, 1822)
	Bull's eye snakehead, Great snakehead	<i>Channa marulius</i> (Hamilton, 1822)
	Asiatic snakehead	<i>Channa orientalis</i> (Bloch & Schneider, 1801)
	Spotted snakehead, Goroi	<i>Channa punctata</i> (Bloch, 1793)
	Golden snakehead, Snakehead murrel, Shoal	<i>Channa stewartii</i> (Playfair, 1867) <i>Channa striata</i> (Bloch, 1793)
Cichlidae	Green chromide, Pearlsnail cichlid	<i>Etroplus suratensis</i> (Bloch, 1790)
	Mozambique tilapia	<i>Oreochromis mossambicus</i> (Peters, 1852)
	Tilapia	<i>Oreochromis niloticus</i> (Linnaeus, 1758)
Clariidae	Clarias catfish	<i>Clarias batrachus</i> (Linnaeus, 1758)
	African catfish	<i>Clarias gariepinus</i> (Burchell, 1822)
	Mangur	<i>Clarias magur</i> (Hamilton, 1822)
Clupeidae	Shortnose gizzard shad	<i>Anodontostoma chacunda</i> (Hamilton, 1822)
	The Ganges river sprat	<i>Corica soborna</i> (Hamilton, 1822)
	White sardine	<i>Escualosa thoracata</i> (Valenciennes, 1847)
	Ganges river gizzard shad	<i>Gonialosa manmina</i> (Hamilton, 1822)
	Indian river shad	<i>Gudusia chapra</i> (Hamilton, 1822)
	Hilsa	<i>Tenualosa ilisha</i> (Hamilton, 1822)
Cobitidae	Almorha loach, Pakistani loach, Yo-Yo loach	<i>Botia almorhae</i> (Gray, 1831)
	Bengal loach, Queen loach	<i>Botia dario</i> (Hamilton, 1822)
	Hora loach	<i>Botia dayi</i> (Hora, 1932)
	Y-loach	<i>Botia lohachata</i> (Chaudhuri, 1912)
	Gangetic loach	<i>Botia rostrata</i> (Günther, 1868)
	Peppered loach, Guntea loach, Scavenger loach	<i>Lepidocephalichthys guntea</i> (Hamilton, 1822)
	Pangia	<i>Pangio pangia</i> (Hamilton, 1822)
Cynoglossidae	Gangetic tongue-sole	<i>Cynoglossus cynoglossus</i> (Hamilton, 1822)
	Long tonguesole	<i>Cynoglossus lingua</i> (Hamilton, 1822)
	Speckled tongue-sole	<i>Cynoglossus puncticeps</i> (Richardson, 1846)
Cyprinidae	Indian carplet	<i>Amblypharyngodon microlepis</i> (Bleeker, 1853)
	Molacarpulet, Pale carplet	<i>Amblypharyngodon mola</i> (Hamilton, 1822)
	Jaya, Choto-piali	<i>Aspidoparia jaya</i> (Hamilton, 1822)
	Kalabans	<i>Bangana dero</i> (Hamilton, 1822)
	Common barb, Spotted barb	<i>Barbodes binotatus</i> (Valenciennes, 1842)
	Barred baril	<i>Barilius barila</i> (Hamilton, 1822)
	Barnabaril	<i>Barilius barna</i> (Hamilton, 1822)
	Hamilton's barila, Dudhnea, Gheur	<i>Barilius bendelisis</i> (Hamilton, 1807)

IUCN Status	Previous Studies	Present Study	Ganga specific biogeographic province/s (2B, 7A, 7B, 8B)
LC	a,b,c,d	+	7A, 7B, 8B
LC	a,b,d		7A, 7B, 8B
DD	b,d		7A, 7B, 8B
LC	a,b,c,f,k	+	7A, 7B, 8B
LC	a,b,c,d,f,g,h,i,k	+	7A, 7B, 8B
NL	b,d,e,f,h,i,j,k		7A, 7B, 8B
LC	a,b,c,d,e,f,g,h,i,j,k	+	7A, 7B, 8B
LC	b,d,e,i		7A, 7B, 8B
LC	a,b,c,d,e,f,g,h,i,k	+	7A, 7B, 8B
LC	a,j		8B
NT	i	+	7A, 7B, 8B
NL	h,i	+	7A, 7B, 8B
LC	b,d,e,f,g,h,i,k		7A, 7B, 8B
LC	h,i	+	7A, 7B
EN	a	+	7A, 7B, 8B
LC	a,d,e,h,i,j,k		8B
LC	a,d,f,h,j		7B, 8B
LC	e,h		8B
LC	a,d,h,i	+	7A, 7B, 8B
LC	a,c,d,e,f,g,h,i,j	+	7A, 7B, 8B
LC	a,c,d,e,f,g,h,i,j	+	7B, 8B
LC	b,d,i		2B, 7A, 7B, 8B
LC	a,b,c,d,f,h,i		2B, 7A, 7B, 8B
NL	b,c,d,f,h		2B, 7A
NL	b,c,d,f,h,i	+	2B, 7A, 7B, 8B
VU	b,d,h	+	2B, 7A, 7B, 8B
LC	a,c,d,e,f,h,i,j,k	+	2B, 7A, 7B, 8B
LC	a,f,h	+	7A, 7B
NL	a,d,h		8B
NL	a,d,e		8B
NL	d,e		8B
LC	c,d,i		7B, 8B
LC	a,b,c,f,i,j,k	+	7A, 7B, 8B
LC	a,b,c,d,f,h,i	+	7A, 7B, 8B
LC	a,c,d,e,f,h,i	+	2B, 7A, 7B, 8B
LC	i		7A
LC	a,b,d,f,h,i		2B
LC	a,b,d,e,f,i	+	2B
LC	a,b,c,d,e,f,h,i	+	2B, 7A

Family	Common Name/s	Scientific Name
	Shacra baril	<i>Barilius shacra</i> (Hamilton, 1822)
	Tileo baril	<i>Barilius tileo</i> (Hamilton, 1822)
	Vagra baril	<i>Barilius vagra</i> (Hamilton, 1822)
	Morar, Ray-finned fish	<i>Cabdio morar</i> (Hamilton, 1822)
	Chaguni, Lal puti	<i>Chagunius chagunio</i> (Hamilton, 1822)
	Silver hatchet	<i>Chela cachius</i> (Hamilton, 1822)
	Mrigal, Mirka	<i>Cirrhinus mrigala</i> (Hamilton, 1822)
	Reba carp	<i>Cirrhinus reba</i> (Hamilton, 1822)
	Gangetic latia	<i>Crossocheilus latius</i> (Hamilton, 1822)
	Grass carp	<i>Ctenopharyngodon idella</i> (Valenciennes, 1844)
	Wild common carp	<i>Cyprinus carpio</i> (Linnaeus, 1758)
	Common carp	<i>Cyprinus carpio</i> (Var. <i>specularis</i>) (Linnaeus, 1758)
	Dangila danio	<i>Danio dangila</i> (Hamilton, 1822)
	Zebra fish, Anju	<i>Danio rerio</i> (Hamilton, 1822)
	Debari, Devario	<i>Devario devario</i> (Hamilton, 1822)
	Flying barb	<i>Esomus danrica</i> (Hamilton, 1822)
	Annandale garra	<i>Garra annandalei</i> (Hora, 1921)
	Gotyla	<i>Garra gotyla</i> (Gray, 1830)
	Lamtagarra	<i>Garra lamta</i> (Hamilton, 1822)
	Catla	<i>Gibelion catla</i> (Hamilton, 1822)
	Silver carp	<i>Hypophthalmichthys molitrix</i> (Valenciennes, 1844)
	Bighead carp	<i>Hypophthalmichthys nobilis</i> (Richardson, 1845)
	Angra labeo	<i>Labeo angra</i> (Hamilton, 1822)
	Minor carp	<i>Labeo bata</i> (Hamilton, 1822)
	Boga labeo	<i>Labeo boga</i> (Hamilton, 1822)
	Boggut labeo	<i>Labeo boggut</i> (Sykes, 1839)
	Orange-fin labeo	<i>Labeo calbasu</i> (Hamilton, 1822)
	Kali, Boalla	<i>Labeo dyocheilus</i> (McClelland, 1839)
	Fringed-lipped peninsula carp	<i>Labeo fimbriatus</i> (Bloch, 1795)
	Kuria labeo	<i>Labeo gonius</i> (Hamilton, 1822)
	Pangusia labeo	<i>Labeo pangusia</i> (Hamilton, 1822)
	Rohu	<i>Labeo rohita</i> (Hamilton, 1822)
	Indian hatchetfish	<i>Laubuka laubuca</i> (Hamilton, 1822)
	Bengala barb	<i>Megarasbora elanga</i> (Hamilton, 1822)
	Gila khani, Cotio	<i>Osteobrama cotio</i> (Hamilton, 1822)
	Rosy barb, Red barb	<i>Pethia conchonius</i> (Hamilton, 1822)
	Golden dwarf barb	<i>Pethia gelius</i> (Hamilton, 1822)
	Glass-barb	<i>Pethia guganio</i> (Hamilton, 1822)
	Spottedsail barb	<i>Pethia phutunio</i> (Hamilton, 1822)
	Ticto barb, Firefin barb	<i>Pethia ticto</i> (Hamilton, 1822)
	Chola barb, Green barb	<i>Puntius chola</i> (Hamilton, 1822)

IUCN Status	Previous Studies	Present Study	Ganga specific biogeographic province/s (2B, 7A, 7B, 8B)
LC	a,b,d,f,h		2B
LC	a,d,h,i		2B
LC	a,b,d,f,h,i	+	2B
LC	a,b,c,d,f,h,i	+	7A, 7B, 8B
LC	a,b,d,e,f,h,i,j	+	7A, 7B, 8B
LC	a,b,c,d,e,f,h,i		7A
LC	a,b,c,d,e,f,g,h,i,j	+	7A, 7B, 8B
LC	a,b,c,d,f,i,j	+	7A, 7B, 8B
LC	a,b,c,d,e,f,h,i,	+	7A, 7B
NL	h,i	+	7A
VU	h,i	+	2B, 7A, 7B
NL	h,i	+	2B, 7A
LC	a,b,c,d,f		2B, 7A
LC	a,c,e,h		2B, 7A
LC	a,b,c,d,f,h,i	+	7A, 7B
LC	a,b,c,d,e,f,h,i,j,k		7A, 7B, 8B
LC	b,c,f		2B
LC	b,c,d,i	+	2B
LC	a,b,f,h		2B
LC	a,b,c,d,e,f,g,h,i,j	+	7A, 7B, 8B
NT	h,i,j	+	7A, 7B, 8B
DD	h,i	+	7A, 7B
LC	a,b,d,f,h,i		7A, 7B, 8B
LC	a,b,c,d,e,f,g,h,i,j	+	7A, 7B, 8B
LC	a,d,e,f	+	7A, 7B
LC	d,i		7A, 7B
LC	a,b,c,d,e,f,g,h,i,j	+	7A, 7B, 8B
LC	b,d,f,h,i	+	2B
LC	a,i	+	2B
LC	a,c,d,f,h,i	+	7A, 7B, 8B
NT	a,b,d,f,h,i	+	7A, 7B, 8B
LC	a,b,c,d,e,f,g,h,i,j	+	7A, 7B, 8B
LC	a,b,c,d,f,h,i,j,k	+	7A, 7B
LC	a,d,h,i		7A
LC	a,b,c,d,e,f,h,i,j	+	7A, 7B, 8B
LC	a,b,c,d,e,f,h,i,j,k	+	7A, 7B, 8B
LC	a,b,d,f,k		7A, 7B, 8B
LC	a,b,d,f		7A, 7B, 8B
LC	a,b,d,f,h,i,k		7A, 7B, 8B
LC	a,b,c,d,e,f,h,i,j,k	+	7A, 7B, 8B
LC	a,b,c,d,e,f,h,i	+	7A, 7B, 8B

Family	Common Name/s	Scientific Name
	Puntio barb	<i>Puntius puntio</i> (Hamilton, 1822)
	Spotfin swamp barb	<i>Puntius sophore</i> (Hamilton, 1822)
	Onespot barb, Teri barb	<i>Puntius terio</i> (Hamilton, 1822)
	Indian trout	<i>Raiamas bola</i> (Hamilton, 1822)
	Slender barb, Blackline rasbora	<i>Rasbora daniconius</i> (Hamilton, 1822)
	Gangetic scissortail rasbora	<i>Rasbora rasbora</i> (Hamilton, 1822)
	Large razorbelly minnow	<i>Salmophasia bacaila</i> (Hamilton, 1822)
	Finescalerazorbelly minnow	<i>Salmostoma phulo</i> (Hamilton, 1822)
	Dapeghat snowtrout, Grot snowtrout, Ladakh snowtrout	<i>Schizothorax curvifrons</i> (Heckel, 1838)
	Chirruhs nowtrout	<i>Schizothorax esocinus</i> (Heckel, 1838)
	Kumaon snow trout	<i>Schizothorax kumaonensis</i> (Menon 1971)
	Kunar snowtrout	<i>Schizothorax labiatus</i> (McClelland, 1842)
	Hill trout	<i>Schizothorax plagiosomus</i> (Heckel, 1838)
	Dinnawah snowtrout	<i>Schizothorax progastus</i> (McClelland, 1839)
	Common snowtrout	<i>Schizothorax richardsonii</i> (Gray, 1832)
	Gora-chela	<i>Securicula gora</i> (Hamilton, 1822)
	Olive barb	<i>Systomus sarana</i> (Hamilton, 1822)
	Dark mahasheer	<i>Tor chelynooides</i> (McClelland, 1839)
	Putitor mahseer, Golden mahaseer	<i>Tor putitora</i> (Hamilton, 1822)
	Mahseer	<i>Tor tor</i> (Hamilton, 1822)
Dasyatidae	Gangetic stingray	<i>Himantura fluviatilis</i> (Hamilton, 1822)*
Eleotridae	Brown spinecheek gudgeon, Dusky sleeper	<i>Eleotris fusca</i> (Forster, 1801)
Engraulidae	Anchovy, Frill-tailed anchovy	<i>Coiliaramcarati</i> (Hamilton, 1822)
	Short-hairfin anchovy	<i>Setipinnabrevifilis</i> (Valenciennes, 1848)
	Gangetic anchovy	<i>Setipinna phasa</i> (Hamilton, 1822)
	Scaly hairfin anchovy	<i>Setipinna taty</i> (Valenciennes, 1848)
	Hamilton's thryssa, Phansha	<i>Thryssa hamiltonii</i> (Gray, 1835)
Erethistidae	Giant moth catfish	<i>Erethistes pusillus</i> (Müller & Troschel, 1849)
	Catfish	<i>Erethistoides pipri</i> (Hora, 1950)
	Kosi hara	<i>Hara hara</i> (Hamilton, 1822)
	Sylhet hara	<i>Hara jerdoni</i> (Day, 1870)

IUCN Status	Previous Studies	Present Study	Ganga specific biogeographic province/s (2B, 7A, 7B, 8B)
NL	a,d,i		7A, 7B, 8B
LC	a,b,c,d,e,f,g,h,i,k	+	7A, 7B, 8B
LC	a,b,d,f,h,i,k	+	7A, 7B, 8B
LC	a,b,c,d,f,g,h,i	+	7A
LC	a,b,d,f,h,i,k	+	7A, 7B, 8B
LC	a,b,d,h,i		7A, 7B, 8B
LC	a,b,c,d,e,f,g,h,i,j,k	+	7A, 7B, 8B
LC	a,d,e,f,h,i,j	+	7A, 7B, 8B
NL	f,i		2B
NL	f		2B
DD	b		2B
NL	f		2B
NL	f,i	+	2B
LC	d,i		2B
VU	b,d,f,g,h,i	+	2B
LC	b,c,d,e,h,i	+	7A, 7B, 8B
LC	a,b,c,d,e,f,g,h,i,j,k	+	7A, 7B, 8B
NL	b,f,i	+	2B
EN	a,b,c,d,f,g,h,i	+	2B
NT	a,b,c,d,f,g,h,i	+	2B
NL	a,d		8B
LC	e,h,j,k		8B
NL	a,d,e		
NL	d,h,i	+	7A, 7B, 8B
LC	c,d,e,f,g,h,i,j	+	7B, 8B
LC	d,e		8B
NL	d,e		8B
LC	c,e,h,i	+	7A, 7B
DD		+	7A
LC	a,b,c,d,f		7A, 7B
LC	a,b,d		7A, 7B

Family	Common Name/s	Scientific Name
Gerreidae	Gangetic silvery-biddy, Small bengal silvery-biddy	<i>Gerres setifer</i> (Hamilton, 1822)
Gobiidae	- Cheeng Boddart's goggle-eyed goby Bareye Goby, Belay Eelgoby - -	<i>Amblyeleotris gymnocephala</i> (Bleeker, 1853) <i>Apocryptes bato</i> (Hamilton, 1822) <i>Boleophthalmus boddarti</i> (Pallas, 1770) <i>Glossogobius giuris</i> (Hamilton, 1822) <i>Odontamblyopus rubicundus</i> (Hamilton, 1822) <i>Pseudapocryptes lanceolatus</i> (Cuvier, 1816) <i>Stigmatogobius sadanundio</i> (Hamilton, 1822)
Hemiramphidae	Black-barred halfbeak Congaturi halfbeak	<i>Hemiramphus far</i> (Forsskål, 1775) <i>Hyporhamphus limbatus</i> (Valenciennes, 1847)
Heteropneustidae	Stinging catfish	<i>Heteropneustes fossilis</i> (Bloch, 1794)
Latidae	Barramundi, Bhetki	<i>Lates calcarifer</i> (Bloch, 1790)
Leiognathidae	Deep pugnose ponyfish	<i>Secutor ruconius</i> (Hamilton, 1822)
Loricariidae		<i>Pterygoplichthys anisitsi</i> (Eigemann & Kennedy, 1903)
Mastacembelidae	Goichi Spiny Eel Barred spiny eel or Indian spiny eel Spiny eel	<i>Macrognathus aculeatus</i> (Bloch, 1786) <i>Macrognathus aral</i> (Bloch & Schneider, 1801) <i>Macrognathus pancalus</i> (Hamilton, 1822) <i>Mastacembelus armatus</i> (Lacepède, 1800)
Mugilidae	Grey mullet, Bangon River mullet Largescale mullet Corsula mullet Yellowtail mullet	<i>Liza tade</i> (Forsskål, 1775) <i>Mugilcephalus</i> (Linnaeus, 1758) <i>Planiliza macrolepis</i> (Smith, 1846) <i>Rhinomugil corsula</i> (Hamilton, 1822) <i>Sicamugil cascasia</i> (Hamilton, 1822)
Muraenesocidae	Daggertooth pike-conger	<i>Muraenesox cinereus</i> (Forsskål, 1775)
Nandidae	Gangetic leaf fish, Nanda, Mottled nandus	<i>Nandus nandus</i> (Hamilton, 1822)
Notopteridae	Indian featherback, Kandla Bronze featherback	<i>Chitala chitala</i> (Hamilton, 1822) <i>Notopterus notopterus</i> (Pallas, 1769)
Ophichthidae	Bengal's snake-eel	<i>Pisodonophis boro</i> (Hamilton, 1822)
Osphronemidae	Sunset gourami Banded gourami Thick-lipped gourami Dwarf gourami	<i>Trichogaster chuna</i> (Hamilton, 1822) <i>Trichogaster fasciata</i> (Bloch & Schneider, 1801) <i>Trichogaster labiosus</i> (Day, 1877) <i>Trichogaster lalius</i> (Hamilton, 1822)
Pangasiidae	Pungas	<i>Pangasius pangasius</i> (Hamilton, 1822)
Platycephalidae	Bartail flathead	<i>Platycephalus indicus</i> (Linnaeus, 1758)
Plotosidae	Canine catfish eel	<i>Plotosus canius</i> (Hamilton, 1822)

IUCN Status	Previous Studies	Present Study	Ganga specific biogeographic province/s (2B, 7A, 7B, 8B)
NL	a		8B
NL	d		8B
NL	a,d,f,h,j	+	7B, 8B
LC	a,d,k		8B
LC	a,b,c,d,e,f,h,i,j,k	+	7A, 7B, 8B
NL	a,d,e,h,j,k		8B
LC	e,h,j,k		7B, 8B
NL	a,d,k		8B
NL	j		8B
LC	c,d,i		7A, 7B, 8B
LC	a,b,c,d,e,f,g,h,i,j,k	+	7A, 7B, 8B
NL	a,d,e,g,h,j		8B
NL	a,j		8B
NL	i		7A, 7B, 8B
NL	a,c,f		7A, 7B, 8B
LC	b,d,e,g,h,i,j,k	+	7A, 7B, 8B
LC	b,c,d,e,g,h,i,j,k	+	7A, 7B, 8B
LC	a,b,c,d,e,f,g,h,i,j,k	+	7A, 7B, 8B
DD	d,e		8B
LC	a,d,e		8B
NL	a,d,e,g,h		8B
LC	a,c,d,e,f,g,h,i,j	+	7A, 7B, 8B
LC	a,b,c,d,f,h,i,j	+	7A, 7B, 8B
NL	k		8B
LC	a,b,c,d,e,f,g,h,i,j,k	+	7A, 7B, 8B
NT	a,b,c,d,e,f,g,h,i,j,k	+	7A, 7B, 8B
LC	a,b,c,d,e,f,g,h,i,j,k	+	7A, 7B, 8B
LC	a,d,h,j,k		7B, 8B
LC	a,d,f,h,k	+	7A, 7B
LC	a,c,d,e,f,h,i,j,k	+	7A, 7B, 8B
LC	d		7A, 7B
LC	a,d,h,k	+	7A, 7B
LC	a,b,c,d,e,f,g,h,i,j,k		7B, 8B
DD	d,h,j		8B
NL	a,d,k		8B

Family	Common Name/s	Scientific Name
Polynemidae	Bamin Indian salmon	<i>Eleutheronema tetradactylum</i> (Shaw, 1804) <i>Polynemus paradiseus</i> (Linnaeus, 1758)
Pristigasteridae	Bigeye ilisha, Indian ilisha	<i>Ilisha megaloptera</i> (Swainson, 1839)
Psilorhynchidae	Balitora minnow Sucatio minnow	<i>Psilorhynchus balitora</i> (Hamilton, 1822) <i>Psilorhynchus sucatio</i> (Hamilton, 1822)
Salmonidae	Rainbow trout	<i>Oncorhynchus mykiss</i> (Walbaum, 1792)
Scatophagidae	Spotted Butter Fish, Spotted Scad	<i>Scatophagus argus</i> (Linnaeus, 1766)
Schilbeidae	Gangetic ailia Garua Bachcha Kocha garua Murius vacha Vacha, tunti Indian potasi Silond catfish, Silong catfish	<i>Ailia coila</i> (Hamilton, 1822) <i>Clupisoma garua</i> (Hamilton, 1822) <i>Clupisoma montana</i> (Hora, 1937) <i>Eutropiichthys murius</i> (Hamilton, 1822) <i>Eutropiichthys vacha</i> (Hamilton, 1822) <i>Neotropiusath erinoides</i> (Bloch, 1794) <i>Silonia silondia</i> (Hamilton, 1822)
Sciaenidae	Big-eyed jewfish, Coitor croaker, Ganges croaker Gangetic bola Bronze croaker Pama Panna croaker	<i>Johnius coitor</i> (Hamilton, 1822) <i>Johnius gangeticus</i> (Talwar, 1991) <i>Otolithoides biauritus</i> (Cantor, 1849) <i>Otolithoides pama</i> (Hamilton, 1822) <i>Panna microdon</i> (Bleeker, 1849)
Sillaginidae	Gangetic sillago, Gangetic whiting	<i>Sillaginopsis panijus</i> (Hamilton, 1822)
Siluridae	Indian butter-catfish Pabdah catfish Pabo catfish Parhin, Helicopter catfish	<i>Ompok bimaculatus</i> (Bloch, 1794) <i>Ompok pabda</i> (Hamilton, 1822) <i>Ompok pabo</i> (Hamilton, 1822) <i>Wallago attu</i> (Bloch & Schneider, 1801)
Sisoridae	Devil catfish, Dwarf goonch Giantdevil catfish, Goonch Indian gagata Gangetic gagata Buhani Kapre Kapre - Kani tengra - Capre Catfish - -	<i>Bagarius bagarius</i> (Hamilton, 1822) <i>Bagarius yarrelli</i> (Sykes, 1839) <i>Gagata cenia</i> (Hamilton, 1822) <i>Gagata gata</i> (Hamilton, 1822) <i>Gagata sexualis</i> (Tilak, 1970) <i>Glyptothorax alaknandi</i> (Tilak, 1969) <i>Glyptothorax annandalei</i> (Hora, 1923) <i>Glyptothorax brevipinnis</i> (Hora, 1923) <i>Glyptothorax cavia</i> (Hamilton, 1822) <i>Glyptothorax conirostris</i> (Steindachner 1867) <i>Glyptothorax garhwali</i> (Tilak, 1969) <i>Glyptothorax gracilis</i> (Günther, 1864) <i>Glyptothorax horai</i> (Fowler, 1934) <i>Glyptothorax madraspatanus</i> (Day, 1873)

	IUCN Status	Previous Studies	Present Study	Ganga specific biogeographic province/s (2B, 7A, 7B, 8B)
	NL	d,e		8B
	NL	a,e,g,j,k		8B
	LC	a,c,d,h		8B
	LC	a,b,d,f		7B, 8B
	LC	a,b,d		7B, 8B
	NL	i	+	2B
	LC	a,d,g,h,j		7B, 8B
	NT	a,b,c,d,e,f,h,i,j,k	+	7A, 7B, 8B
	LC	a,b,c,d,e,f,g,h,i,j	+	7A, 7B, 8B
	LC	b,f		7A, 7B, 8B
	LC	a,d,h,i,j	+	7A, 7B, 8B
	LC	a,b,c,d,e,f,g,h,i,j	+	7A, 7B, 8B
	LC	e,h,j	+	7A, 7B, 8B
	LC	a,b,c,d,f,g,h,i,j	+	7A, 7B, 8B
	LC	a,c,d,f,h,i	+	7A, 7B, 8B
	NL	d,h,i,j	+	7A, 7B, 8B
	NL	d,e		8B
	NL	d,f		8B
	NL	d,e,i		7B,8B
	NL	a,d,e,g,h,j		8B
	NT	a,b,c,d,e,f,g,h,i,k	+	7A, 7B, 8B
	NT	a,b,d,e,f,g,h,i,j,k	+	7A, 7B, 8B
	NT	a,b,c,d,e,i		7A, 7B, 8B
	NT	a,b,c,d,e,f,g,h,i,j,k	+	7A, 7B, 8B
	NT	a,b,c,d,e,f,g,h,i,j,k	+	7A, 7B, 8B
	NT	b,h,i	+	7A, 7B, 8B
	LC	a,b,c,d,e,f,h,i,j	+	7A, 7B, 8B
	LC	a,b,d,h,j		7A, 7B, 8B
	LC	b,d,f,h,j	+	7A, 7B, 8B
	LC	b,d		2B
	LC	b,f		2B
	DD	d,f,i		2B
	LC	a,b,d,f,h	+	2B, 7A
	DD	b,f		2B
	LC	b,d		2B
	DD	b,d	+	2B
	LC	d,f		2B
	EN	f		2B

Family	Common Name/s	Scientific Name
	River cat fish, Nayid Telchitta, Dhal magur Three lined catfish Kosi nangra Sisor catfish	<i>Glyptothorax pectinopterus</i> (McClelland, 1842) <i>Glyptothorax telchitta</i> (Hamilton, 1822) <i>Glyptothorax trilineatus</i> (Blyth, 1860) <i>Nangra nangra</i> (Hamilton, 1822) <i>Sisor raddophorus</i> (Hamilton, 1822)
Soleidae	Oriental-sole	<i>Brachirus orientalis</i> (Bloch & Schneider, 1801)
Sparidae	Yellowfin seabream	<i>Acanthopagrus latus</i> (Houttuyn, 1782)
Synbranchidae	Rice swamp eel, Asian swamp eel Gangetic mudeel, Cuchi	<i>Monopterus albus</i> (Zuiew, 1793) <i>Monopterusuchia</i> (Hamilton, 1822)
Syngnathidae	Crocodile-tooth Pipefish	<i>Microphis cuncalus</i> (Hamilton, 1822)
Synodontidae	Bombay duck	<i>Harpadon nehereus</i> (Hamilton, 1822)
Terapontidae	Tiger perch, Crescent grunter	<i>Terapon jarbua</i> (Forsskål, 1775)
Tetraodontidae	Ocellated pufferfish Green pufferfish	<i>Leiodon cutcutia</i> (Hamilton, 1822) <i>Dichotomysctere fluviatilis</i> (Hamilton, 1822)
Toxotidae	Spotted archerfish	<i>Toxotes chatareus</i> (Hamilton, 1822)

^aHamilton (1822); ^bMenon (1962); ^cBilgrami (1991); ^dTalwar (1991); ^eMishra et al (2003); ^fPayne et al (2004); ^gVass et al (2010); ^hDas et al (2010); ⁱSarkar et al (2012); ^jNath&Patra (2015); ^kDubey et al (2015)

*Note: Only two species viz, *Glyphis gangetica* (Gangetic shark) and *Himantura fluviatilis* (Gangetic stingray) have been placed in Schedule I of the Indian Wildlife (Protection) Act, 1972 (IWPA).

IUCN Status	Previous Studies	Present Study	Ganga specific biogeographic province/s (2B, 7A, 7B, 8B)
LC	d,f,i		2B
LC	a,b,f,h,i,j		2B
LC	f		2B
LC	a,b,c,d,f,i		7B
LC	a,b,c,d,h,i	+	7A,7B,8B
NL	h,j		8B
DD	a,d,j		8B
LC	i		7A,7B
LC	a,d,e		7B,8B
LC	d,j		8B
NL	a,d,g		8B
LC	a,d,e		8B
LC	a,c,d,f,h,i,j,k	+	7A,7B,8B
LC	a,c,d,f,i,k		7A
NL	a,d,h		8B

REFERENCES

- Bilgrami, K.S. (1991). Biological profile of the Ganga: Zooplankton, Fish, Birds, and other minor fauna. In Krishna Murti, C.R., Bilgrami, K.S., Das, T.M. & Mathur, R.P. (Eds.). *The Ganga: A Scientific study*. Ganga Project Directorate, Ministry of Environment & Forests, Government of India. Northern Book Centre, New Delhi, 83-94.
- Das, M. K., Sharma, A. P., Tyagi, R. K., Saha, P. K., Pathak, V., Suresh, V. R., De, D. K., Paul, S. K., Sett, P., Chakrabarty, M., & Mondal, K. (2010). Fishes of River Ganga - A Field Identification Manual. *Central Inland Fisheries Research Institute, Barrackpore, West Bengal*, 165, 1-100.
- Dubey, S. K., Trivedi, R. K., & Chand, B. K. (2015). Indigenous freshwater piscine resources of Indian Sundarban biosphere reserve: status and prospects. *World Journal of Fish and Marine Sciences*, 7(1), 21-28.
- Hamilton, F. (1822). *An account of the fishes found in the river Ganges and its branches*. London: Archibald Constable and Company, Edinburgh and Hurst, Robinson, and Co. Cheapside.
- Menon, A. G. K (1962). A distributional list of fishes of the Himalayas. *Journal of Zoological Society of India*, 14(1) 23-32.
- Mishra, S. S., Pradhan, P., Kar S., & Chakraborty, S.K. (2003). Ichthyofaunal diversity of Midnapore, Bankura and Hooghly Districts, South West Bengal. *Recording of Zoological Survey of India*, 220, 1-65.
- Nath, A. K., & Patra, A. (2015). Survey on the present status of fish species diversity in a stretch of Hooghly river of West Bengal, India. *International Journal of Fisheries and Aquatic Studies*, 3(1), 244-250.
- Payne A.I., Sinha R., Singh H.R., & Huq S. (2004). A review of the Ganges Basin; its fish and fisheries. In Welcomme, R.L. & Petr, R.L. (Eds.) *Proceedings of the Second International Symposium on the Management of Large Rivers for Fisheries Volume 1*. Food and Agriculture Organization of the United Nations & Mekong River Commission. FAO Regional Office for Asia and the Pacific, Bangkok. RAP Publication 2004/16, 229- 252
- Sarkar, U. K., Pathak, A. K., Sinha, R. K., Sivakumar, K., Pandian, A. K., Pandey, A., Dubey, V.K., & Lakra, W. S. (2012). Freshwater fish biodiversity in the River Ganga (India): changing pattern, threats and conservation perspectives. *Reviews in Fish Biology and Fisheries*, 22(1), 251-272.
- Talwar, P. K. (1991). Pisces. In Director, Zoological Survey of India (Eds.) *Faunal Resources of Ganga, Part 1. Zoological Survey of India*, Calcutta. 59-145.
- Vass, K.K., Mondal, S.K., Samanta, S., Suresh, V.R., & Katiha, P.K. (2010). The environment and fishery status of the river Ganges. *Aquatic Ecosystem Health and Management Society*, 13 (4), 385-394.



PHOTO CREDITS

Abhijit Das, Arvind Dwivedi, Bivash Pandav, Bhumesh Bharti, Ganesh Chowdhury, Goura Chandra Das, Indraneil Das, Klaus rudloff, Moumita Chakraborty, Narendra Mohan, Niladri Dasgupta, Pariva Dobriyal, Rajeev Basumatary, SA Hussain, Suyash Katdare, Threesh Kapoor and miscellaneous source.



**National Mission
for Clean Ganga**

NMCG

National Mission
for Clean Ganga,
Ministry of Water
Resources, River
Development &
Ganga Rejuvenation



**भारतीय वन्यजीव संस्थान
Wildlife Institute of India**

GACMC

Ganga Aqualife
Conservation
Monitoring
Centre

Wildlife Institute of India

Post Box #18,
Chandrabani
Dehradun - 248001
Uttarakhand, India
t.: +91 135 2640114-15,
+91 135 2646100
f.: +91 135 2640117



*Printed on high
quality FSC,
Recycled, eco
friendly & carbon
neutral fine papers.
WII supports the
responsible and
sustainable use of
our forest
resources.*